



## ENVIRONMENTAL INDICATORS

# 2014


Republic of Macedonia  
Ministry of Environment and Physical Planning  
Macedonian Environmental Information Center

# AIR POLLUTION



## **MK-NI 001**

### **EMISSIONS OF ACIDIFYING SUBSTANCES**



#### **Definition**

The indicator tracks the trends in anthropogenic emissions of acidifying substances, i.e. acidifying processes in the air. These substances include nitrogen oxides, ammonia, and sulphur dioxide, and their acidifying power is weighted by their acidifying potential.

The indicator also provides information on emissions by sectors: energy generation and transformation, road and other transport, industry (processes and energy), fugitive emissions, waste, agriculture and other.

#### **Units**

- kt (acidifying equivalent)

#### **Key policy issue**

**What progress has been made in overall reducing acidifying substances emissions in the air?**

#### **Key message**


##### **Sulfur dioxide (SO<sub>2</sub>)**

Compared to 2005, emissions of sulfur dioxide dropped by 11.28 %, though emissions of sulfur dioxide vary for the whole reviewed period and there were no significant changes due to the fact that no best available techniques were applied in the key sector by energy producing installations, in relation to this pollutant. In addition to this, SO<sub>2</sub> emissions from fuels combustion in industry and administrative facilities owing to use of crude oil with sulfur content up to 1%.

##### **Nitrogen oxide (NO<sub>x</sub>)**

The trend of nitrogen oxide emissions is variable because data used in the period 2008-2012 for electricity and heat production sector was from measured concentrations by installations, while for the preceding years calculations were applied based on emission factor and therefore emissions were lower. With regard to emissions of this pollutant from traffic, there was slight increase in these pollutants during the last three years due to increased number of diesel driven vehicles.

Diagram 1. Emission trends for nitrogen oxides and sulfur dioxide



## Assessment

Under the CORINAIR Programme, inventory of air pollutants emission was established in 2005 by individual sectors, i.e. activities, and in 2012 Inventory including all pollutants was prepared.

Sectors based on EMEP/EEA guideline and SNAP – selective nomenclature are presented in the table below:

SNAP	Sector
1	Combustion in energy and transformation industries (stationary sources)
2	Non-industrial combustion plants (stationary sources)
3	Combustion in manufacturing industry (stationary sources)
4	Production processes (stationary sources)
5	Extraction and distribution of fossil fuels and geothermal energy
6	Solvent and other product use
7	Road transport
8	Other mobile sources and machinery
9	Waste treatment and disposal
10	Agriculture
11	Nature

Besides the Inventory identifying the quantities of air emissions of the main pollutants, data was also used from the updated Cadastre of air polluters and pollutants in the Republic of Macedonia for the period 2008-2009, and it identified air pollutants at the level of facilities.

Here, analysis was made of the trend of emission of the pollutants nitrogen oxides and sulfur dioxide for the period 2004 to 2012 and it was planned to conduct calculation and recalculation of emissions for the period 1990-2003 in the course of 2015. Calculations of emissions for the period 2002-2003 were modified in this indicator report compared to the previous one, because they involve great uncertainty deriving from the use of data from measurement instead of emissions factors specified in EMEP/EEA guideline which was the case for years following 2004.

With reference to the trend of nitrogen oxides, it was variable, but it should be pointed out that data

used for the period 2008-2012 for the electricity and heat production sector was from concentrations measured by installations, while calculations were used for the preceding period based on emission factor and therefore emissions were lower. Inventory taken of ozone precursors indicated that there were no significant reductions in the emissions of these pollutants during the analyzed period.

Nevertheless, compared to emissions in 1990 as baseline year of the Protocols to CLRTAP, there was reduction in ozone precursors and emission ceilings specified in Annex II of the Gothenburg Protocol were not exceeded.


Reduction of SO<sub>2</sub> compared to emissions in 1990 as baseline year by 30% resulted from reduction of sulfur content in oil derivatives. Reduction of sulfur content in fuels was achieved by transposition and implementation of Directive 1999/32/EC on the quality of liquid fuels into the Macedonian legislation. Emissions of nitrogen oxides compared to 1987 (baseline year under the Protocol on NO<sub>x</sub>) were reduced by 23% or by 20% compared to 1990. In 2012, emissions of NO<sub>x</sub> (35 Gg) were below the national emission ceiling. Emission reduction resulted from reduced consumption of crude oil in combustion plants.

The above resulted from reduction of emissions from the traffic due to introduction of catalytic converters and improved composition of fuel composition, as well as improved EURO standards, especially for light freight and heavy vehicles. As far as the implementation of IPPC Directive and introduction of the best available techniques in electricity and heat producing installations are concerned, these techniques have been introduced for NO<sub>x</sub> and emission reduction is expected for 2014-2015.

## Policy specific issue

### Which different sectors and processes contribute to acidifying substances emissions?


Diagram 2. Total SO<sub>x</sub> emissions by sectors on annual level



With regard to contributions of different sectors in 2012, 93.8% of the total quantity of SO<sub>2</sub> was released during fuels combustion in the process of electricity and heat generation, 3% was released from combustion plants in manufacturing industry. Considering that the key sector contributes 93.8%,

reduction of sulfur dioxide emissions can be achieved only through introduction of the best available techniques in the two largest installations for electricity production at the level of Macedonia – MEC Bitola and TPP Oslomej. Under the agreement with the energy community, installations for electricity production should, by the end of 2015, prepare plan for gradual reduction of emissions and achievement of emission limit values by the end of 2017. This is expected to result in significant reduction of sulfur dioxide after 2017.

Diagram 3. Total NOx emissions by sectors on annual level



Data coverage: excel

Source: State Statistical Office, Energy balance of the country – Report by the Government, Cadastre of Air Polluters and Pollutants, Data from measurements in companies – major polluters, Database on motor vehicles of the Ministry of Interior, Project: Introduction of CORINAIR Inventory Methodology of the Ministry of Environment and Physical Planning, Spatial Plan of the Republic of Macedonia.

In 2012, the greatest quantities of nitrogen oxides emissions were released from electricity production (54.97%), road transport (28.41%) and combustion in manufacturing industry with 8.49%.

Slight increase of emissions was recorded in transport sector in the period 2010-2012 which was most probably due to import of 100.000 used vehicles with low Euro standards. With regard to emissions from SNAP 1 sector – Combustion and transformation of energy in electric power facility, different methodology was used in the period 2004-2007 and period 2008-2012 and thus we cannot derive specific conclusion about the trend in this sector. Nevertheless, given the fact that no specific activities were undertaken in this sector before 2012, no significant changes are expected upon alignment of the methodology. With regard to emissions of nitrogen oxides from SNAP 1 sector, measures undertaken so far by the installations MEC Bitola and District Heating (Toplifikacija AD) Skopje, significant emission reduction can be expected in 2013.

## Methodology

- **Methodology for the indicator calculation**

The methodology for this indicator calculation is based on aggregation and calculation of data on SO<sub>2</sub>, NH<sub>3</sub> and NO<sub>x</sub> emissions at annual basis, on national level, as overall and distributed to sectors, i.e. activities.

Calculations are in line with the Guidelines of UNECE/EMEP Convention on Long-Range Transboundary Air Pollution (LRTAP Convention), as well as application of the SNAP – selective nomenclature of air pollution. With regard to this specific indicator, factors have been used in order to express the acidifying property potential. These factors are for NO<sub>x</sub> 0.02174, for SO<sub>2</sub> 0.03125 and for NH<sub>3</sub> 0.05882, respectively. The results are expressed in kilotonnes equivalent acidity

### **Uncertainty**

- **Methodological and data uncertainty**

Use of factors of the acidifying property potential leads to some uncertainty. Furthermore, national emission factors obtained by measurements in the Republic of Macedonia are used only in energy sector. For other sectors, standard emission factors from the EMEP/EEA Guideline are used for the inventory of air pollutants. It has been assumed that the factors are representative for Europe as a whole; different factors may be estimated at local level. Comprehensive discussion on uncertainty of these factors can be found in de Leeuw (2002). As far as activity installment is concerned, uncertainty arises from data taken from Statistical Yearbook and other sources. Definition of activity installment which is not required form, expert estimations containing uncertainty are made.

- **Reference of used methodology**

EMEP/EEA Guideline for inventory of air pollutant emissions 2013, Technical Report no. 12/2013, EEA.

## Policy relevance of the indicator

### **List of relevant policy documents**

Action Plan for European Partnership, as well as National Plan for approximation of the national legislation with European regulations specifying bylaws that need to be prepared has been adopted.

The National Environmental Action Plan (NEAP II) has been adopted. It contains the measures that need to be taken to improve the overall status of air quality, including the reduction of emissions of acidifying substances. The National Plan for Ambient Air Protection for the period 2012 to 2017 and the National Programme for gradual air emissions reduction by 2020 have been adopted in order to define and implement measures on national level concerning introduction of renewable energy sources, application of the code of good agricultural practice, technical control of vehicles at registration, application of the best available techniques in industrial facilities, etc. At the same time, for the purpose of air quality improvement in certain local self-government units (LSGUs) with action plans, pilot programme was prepared for the City of Bitola, serving as basis for preparation of local planning documents in other cities.

All 8 Protocols to the Convention on Long-Range Transboundary Air Pollution (CLRTAP) were ratified in 2010. With regard to the last three Protocols, i.e. Protocol on heavy metals, Protocol on POPs and Gothenburg Protocol, National Action Plan for Ratification and Implementation was adopted at their request. Due to the requirement for modifications in contributions related to emissions in baseline year (1990) and national emission ceilings for 2010, the Gothenburg Protocol and Protocol on sulfur of 1995 entered into force for the Republic of Macedonia in 2014, upon the adoption of the values

specified in Annex II to these Protocols.

In order to fulfill one of the basic requirements under the Protocols to CLRTAP, Inventory of Air Pollution is prepared on regular annual level by EMEP/EEA Methodology and the prepared inventory is reported to UNECE and CLRTAP and European Environmental Agency.

With regard to fulfillment of the requirements of the Stockholm Convention covering the same pollutants specified in the Protocol on POPs, the National Implementation Plan for old and new organic pollutants was updated and Inventory of old and new POPs chemicals was prepared.

## **Legal grounds**

The Law on Ambient Air Quality was adopted in August 2004 and amended several times afterwards (Official Gazette of RM no. 67/2004, 92/2007, 83/2009, 35/10, 47/11, 100/12, 163/2013) as framework law in the area of air. The goals of this Law include avoiding, prevention and reduction of harmful effects on human health and environment as a whole, prevention and abatement of pollutions leading to climate change, as well as provision of appropriate information on the quality of ambient air.

On the basis of the Law on Ambient Air Quality, 16 bylaws were prepared and adopted to introduce limit values for air quality and air emissions, methodology of air quality and air emissions monitoring, manner of preparation of planning documents for air protection against pollution, manner of informing the citizens and international organizations, etc.

With reference to air standards transposed in part of the mentioned rulebooks, 86 ISO and 48 CEN standards in the area of air emissions and air quality were adopted by means of endorsement method.

Other legislation related to the regulation of air quality and air emissions regulation includes the Law on Vehicles, Law on Standardization, Rulebook on liquid fuels quality with national standards for liquid fuels quality, etc.

## **Targets**

### **Does any of the national documents set targets or targets set under international documents should be achieved?**

National documents listed as references in the above text provide guidelines and specify actions that should be undertaken as a matter of priority. It is important to mention that the transposition of Directives 96/61/EC, 2000/81/EC, 2000/76/EC, 99/13/EC and 2001/81/EC into the national laws and bylaws ranges between 90-100%, while activities towards their implementation are in progress.

In accordance with the requirements of the UNECE Convention on Long-Range Transboundary Air Pollution, inventory based on the CORINAIR Programme has been introduced, setting the target of regular inventory of pollutants in tons per year following the n-2 principle, where n is the current year.

Also, in accordance with Directive 2001/81/EC, as well as Gothenburg Protocol, the ceilings of the amounts of emissions have been set at the level of the Republic of Macedonia for 2010 that shall not be exceeded at annual level. The Executive Body of the Convention on Long-Range Transboundary Air Pollution, upon submission of the values of national ceilings in order to enrol the Republic of Macedonia in Annex II of the Gothenburg Protocol requested correction of the values considering the reported data on air emissions of the pollutants sulfur dioxide and ammonia at national level. Changes in the values of these pollutants were incorporated in the Rulebook amending the Rulebook on upper limits – emission ceilings of pollutants for the purpose of setting projections for certain period concerning reduction of the quantities of pollutant emissions at annual level published in July 2014.



In 2012, national upper limits – ceilings for SO<sub>2</sub> and NO<sub>x</sub> were not exceeded.

	Upper limit - ceiling	Total emissions-2012
SO <sub>2</sub>	110 Gg	89.3906 Gg
NO <sub>x</sub>	39 Gg	35.0683 Gg

With regard to targets – emission projections for SO<sub>2</sub> and NO<sub>x</sub> for 2020, they amount 84.63 Gg, and for nitrogen oxides under the scenario with measures, emission level of 23.8 Gg should be achieved.

## Reporting obligation


- Reporting obligations under multilateral agreements – UNECE/CLRTAP and EEA
- Annual report of processed data on environment

## General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 001</b>	<b>Emissions of acidifying substances</b>	CSI 001	Emissions of acidifying substances	<b>P</b>	<b>B</b>	<ul style="list-style-type: none"> <li>▪ acidification</li> <li>▪ air</li> </ul>	annually

## MK - NI 002

### EMISSIONS OF OZONE PRECURSORS



#### Definition

This indicator tracks trends in emissions of ozone precursors: nitrogen oxides, carbon monoxide, methane and non-methane volatile organic compounds, caused by anthropogenic activities, and each precursor is weighted by its tropospheric ozone-forming potential.

The indicator also provides information on emissions by sectors: energy industries; road and other transport modes; industry (processes and energy); other (energy); fugitive emissions; waste; agriculture and other (non energy).

#### Units

- kt (NMVOC - equivalent)


#### Key policy issue

##### ***What progress has been made in overall ozone precursors emissions reduction in Europe?***

- Emissions of ozone precursors in 2012 compared to 2004 increased for nitrogen oxides (NO<sub>x</sub>) by 4.33%, they decreased for non-methane volatile organic compounds (NMVOC) by 22.34% and carbon monoxide (CO) by 23.44%, and increase was recorded for methane (CH<sub>4</sub>) by 27.47%.
- Increase in emissions of nitrogen oxides during the last several years has resulted from the application of measured instead of estimated emissions in energy sector. For the period 2008-2011, measured values were taken for NO<sub>x</sub> instead of estimated (determined through emission factors) as those became available due to introduction of stricter national legislation in terms of their submission by operators. Increase in nitrogen oxides from traffic during the last several years resulted from increased consumption of diesel fuels related to increased number of this type of vehicles. With regard to CO, there was reduction in the sector of combustion in households, and reduction in the emission for NMVOC resulted from the solvent use sector. For methane emissions, significant increase in the emission of this substance resulted from the key sector, i.e. increased amount of solid waste.

The Diagram below shows annual trend of carbon monoxide, methane, non-methane organic compounds and nitrogen oxides emissions presented as ozone precursors.

Diagram 1. Total ozone precursors emissions



## Assessment

In 2005, in the frames of the CORINAIR Programme, the Inventory of Air Emissions was established in the country, presenting emissions by individual sectors, i.e. activities, and in 2012 Inventory including all pollutants was prepared.

Sectors based on the above stated Methodology and SNAP – selective nomenclature are given in the table below:

SNAP	
1	Combustion in energy and transformation industries (stationary sources)
2	Non-industrial combustion plants (stationary sources)
3	Combustion in manufacturing industry (stationary sources)
4	Production processes (stationary sources)
5	Extraction and distribution of fossil fuels and geothermal energy
6	Solvent and other product use
7	Road transport
8	Other mobile sources and machinery
9	Waste treatment and disposal
10	Agriculture
11	Nature

Besides the Inventory identifying the quantities of air emissions of the main pollutants, data was also used from the updated Cadastre of air polluters and pollutants in the Republic of Macedonia and it identified air pollutants at the level of facilities. The latest updating of data was made in 2008-2009,

Inventory of pollutants is also made by the main sectors of relevance for climate change caused phenomena, based on the United Nations Framework Convention on Climate Change – UNFCCC. Inventory of methane as one of ozone precursors is made by IPCC methodology. In the frames of the Third National Communication on Climate Change, GHG Inventory was made for the period 1990-2012.

Completed inventory of ozone precursors indicates variable trend during the analyzed period.

Changes in the quantities of NOx and CO emissions in transport sector resulted from the change in consumed quantities of diesel fuels and petrols by passenger cars. As far as reduction in NOx emissions from energy sector is concerned, the implementation of IPPC Directive and introduction of the best available techniques in electricity and heat production installations in 2013, reduction is expected in emissions of this pollutant from energy sector upon 2013.

Directive 1999/13/EC regarding NMVOC emissions from solvents use in installations and certain activities has been partially transposed (only for limit values) in the national legislation, and reduction schemes for these pollutants have not been introduced. On the other side, transposition of Directives 1994/63/EC and 2009/126/EC concerning emissions from petrol filling and distribution to petrol stations is in final stage and implementation of the Law on Control of Volatile Organic Compound Emissions at petrol use will commence in the course of 2015. Namely, it is necessary to initiate procedure for registration of installations for storage, installations for filling and emptying of mobile containers and petrol stations in accordance with the adopted rulebook.

With regard to greenhouse gas methane, overall emissions have increased due to emission increase in waste sector owing to the trend of growth in the quantity of solid waste. Emission of methane from agriculture sector was dropping by 2004 and then followed a stable trend.

## Specific policy issue

**Which different sectors and processes contribute to ozone precursors emissions?**

Diagram 2. Total NOx emissions by sectors/year




Diagram 3. Total CO emissions by sectors/year




Diagram 4. Total NMVOC emissions by sectors/year





Diagram 5. Total CH4 emissions by sectors/year



Data coverage: [excel](#)

**Source:** State Statistical Office, Energy balance of the country – Report by the Government, Cadastre of Air Polluters and Pollutants, Data from measurements in companies – major polluters, Database on motor vehicles of the Ministry of Interior, Project: Introduction of CORINAIR Inventory Methodology of the Ministry of Environment and Physical Planning, Spatial Plan of the Republic of Macedonia.

### Nitrogen oxides (NOx)

Diagram 3 indicates that the biggest quantities of nitrogen oxides emissions are released from electricity production, followed by road transport. As mentioned above, introduction of measures for reduction of emissions from electricity and heat production installations and reduced operation regime in OKTA Refinery will result in reduction of NOx emissions in this sector, but for 2014.

### Carbon monoxide (CO)

During the period 2010-2012, emissions of nitrogen oxide and carbon monoxide from transport increased and reduced due to change in fuels consumption in traffic. The next key source of carbon monoxide emission is the sector of non-industrial combustion plants (households and administrative facilities). As far as CO emissions from non-industrial combustion plants are concerned, slight drop in emissions has been recorded during the last year in this sector. However, major reduction in this sector has not been observed because of incomplete combustion of fuels used by households and administrative facilities, especially when heating wood is used.

### Nonmethane volatile organic compounds (NMVOCs)

Nonmethane volatile organic compounds are associated with several sources of emission with contribution of 15-20%. This indicates that there are no predominant sectors in the case of these substances. In the period 2004-2012, there was drop in the emissions of these substances only from the sector of solvents use.

### Methane (CH<sub>4</sub>)

The most significant is the percentage of emission increase in waste sector in 2010 relative to 2009 amounting 40% (followed by stable trend in methane emissions from this sector) which

induces rising trend in the total emissions of this pollutant.

## Methodology

- Methodology for the indicator calculation

The methodology for this indicator calculation is based on aggregation and calculation of data on CO, NMVOC, CH<sub>4</sub> and NO<sub>x</sub> emissions at annual basis, on national level, as overall and distributed to sectors, i.e. activities.

Calculations are in line with EMEP/EEA Guidelines and methodology for inventory and application of the SNAP – selective nomenclature of air pollution.

With regard to this specific indicator, factors have been used in order to express the property of ozone precursors. These factors are specific to each pollutant, namely for NO<sub>x</sub> it is 1.22, for NMVOC it is 1, for CO it is 0.11 and for CH<sub>4</sub> this value is 0.014. The results are expressed in kilotons NMVOC equivalent.

- Reference of used methodology

Methodology applied in the calculation and presentation of this indicator has been taken from the EMEP/EEA Guideline on air pollutant emission inventory 2013, Technical Report no. 12/2013, EEA. Set of emission indicators of long-range transboundary air pollution, Environmental science and policy.

## Policy relevance of the indicator

The European Partnership Action Plan has been adopted, as well as the National plan for approximation of the national legislation with the European regulations stating the bylaws that need to be prepared.

NEAP II has been prepared, specifying the measures that need to be taken to improve the overall status of air and in that sense to reduce emissions leading to acidification. The National Ambient Air Protection Plan 2012-2017 has been adopted, and it contains measures for air protection on national level and the National program for gradual reduction of emissions until 2020 in order to define and implement measures on national level concerning introduction of renewable sources, application of the code of good agricultural practice, technical control of vehicles at registration, application of the best available techniques in industrial facilities, etc., as well as technical control and on road checks. At the same time, for the purpose of air quality improvement in certain local self-government units (LSGUs) with action plans, pilot programme was prepared for the City of Bitola, serving as basis for preparation of local planning documents in other cities.

All 8 Protocols to the Convention on Long-Range Transboundary Air Pollution (CLRTAP) were ratified in 2010. With regard to the last three Protocols, i.e. Protocol on heavy metals, Protocol on POPs and Gothenburg Protocol, National Action Plan for Ratification and Implementation was adopted at their request. Due to the requirement for modifications in contributions related to emissions in baseline year (1990) and national emission ceilings for 2010, the Gothenburg Protocol and Protocol on sulfur of 1995 entered into force for the Republic of Macedonia in 2014, upon the adoption of the values specified in Annex II to these Protocols.

In order to fulfill one of the basic requirements under the Protocols to CLRTAP, Inventory of Air Pollution is prepared on regular level by EMEP/EEA Methodology and the prepared inventory is reported to UNECE and CLRTAP and European Environmental Agency.

With regard to fulfillment of the requirements of the Stockholm Convention covering the same pollutants specified in the Protocol on POPs, the National Implementation Plan for old and new organic pollutants was updated and Inventory of old and new POPs chemicals was prepared.

## Legal grounds

The Law on Ambient Air Quality was adopted in August 2004 and amended several times afterwards (Official Gazette of RM no. 67/2004, 92/2007, 83/2009, 35/10, 47/11, 100/12, 163/2013) as framework law in the area of air. The goals of this Law include avoiding, prevention and reduction of harmful effects on human health and environment as a whole, prevention and abatement of pollutions leading to climate change, as well as provision of appropriate information on the quality of ambient air.

On the basis of the Law on Ambient Air Quality, 16 bylaws were prepared and adopted to introduce limit values for air quality and air emissions, methodology of air quality and air emissions monitoring, manner of preparation of planning documents for air protection against pollution, manner of informing the citizens and international organizations, etc.

With reference to air standards transposed in part of the mentioned rulebooks, 86 ISO and 48 CEN standards in the area of air emissions and air quality were adopted by means of endorsement method.

Other legislation related to the regulation of air quality and air emissions regulation includes the Law on Vehicles, Rulebook on liquid fuels quality with national standards for liquid fuels quality, etc.

## Targets

### **Does any of the national documents set targets or targets set under international documents should be achieved?**

National strategic documents listed as references in the above text provide guidelines and specify actions that should be undertaken as a matter of priority. It is important to mention that the new regulations in the area of air emissions have been adopted transposing the following Directives into the national legislation: 96/61/EC, 2000/81/EC, 2000/76/EC, 99/13/EC and 2001/81/EC and the level of transposition is 90-100%.

In accordance with the requirements of the UNECE Convention on Long-Range Transboundary Air Pollution, inventory taking was introduced based on the EMEP/EEA Guideline on air pollutant emission inventory aimed at regular inventory of pollutants in tonnes per year following n-2 principle where n is the current year.

Also in accordance with Directive 2001/81/EC and Gothenburg Protocol, the upper limits-ceilings of the amounts of emissions have been set at the level of the Republic of Macedonia for 2010 that shall not be exceeded at 2010 annual level. The Executive Body of the Convention on Long-Range Transboundary Air Pollution, upon submission of the values of national ceilings in order to enroll the Republic of Macedonia in Annex II of the Gothenburg Protocol requested correction of the values considering the reported data on air emissions of the pollutants sulfur dioxide and ammonia at national level. Changes in the values of these pollutants were incorporated in the Rulebook amending the Rulebook on upper limits – emission ceilings of pollutants for the purpose of setting projections for certain period concerning reduction of the quantities of pollutant emissions at annual level published in July 2014. In 2012, national upper limits – ceilings for SO<sub>2</sub> and NO<sub>x</sub> were not exceeded.

	Upper limit - ceiling	Total emissions-2012
NM VOC	30 Gg	27.4783
NO <sub>x</sub>	39 Gg	35.0683



With regard to targets – emission projections for NO<sub>x</sub> for 2020, under the scenario with measures, emission level of 23.8 Gg should be achieved. Under the scenario with measures, emission increase is expected for NMVOC of up to 36.67 Gg, which indicates the need to introduce additional measures to reduce this ozone precursor.

These documents set the basis for achievement of the targets for reduction of emissions of ozone precursors thus resulting in reduced degradation of environment and negative effect on human health.

## Reporting obligation

- Reporting obligations under multilateral agreements – UNECE transboundary air pollution transfer, as well as to EEA
- Annual report of processed data on air emissions

## General meta-data

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 002</b>	<b>Emissions of ozone precursors</b>	CSI 002	Emissions of ozone precursors	Π	<b>A</b>	<ul style="list-style-type: none"> <li>▪ air</li> <li>▪ air quality</li> </ul>	annually

## MK - NI 004

# EXCEEDANCE OF AIR QUALITY LIMIT VALUES IN URBAN AREAS – By monitoring station



## Definition

This indicator shows:

- Number of days during year when the level of air pollution (for the pollutants: suspended particulate matters sized up to 10 micrometers (PM<sub>10</sub>), sulphur dioxide (SO<sub>2</sub>), nitrogen dioxide (NO<sub>2</sub>), and ground ozone (O<sub>3</sub>)) exceeds the prescribed limit values (maximum permissible annual and short-term concentrations in urban areas where regular observation of air quality is performed).
- Portion of urban population living in urban areas with at least one monitoring station) in the country exposed at air pollution above the set limit values.
- Absolute values of the concentration of pollutants in the air.

## Units

Number of days

Ambient air concentrations of sulphur dioxide (SO<sub>2</sub>), particulate matter sized up to 10 micrometer (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) are expressed in microgram/m<sup>3</sup> (µg/m<sup>3</sup>).

## Key policy issue

*What progress has been achieved in reducing the concentrations of pollutants in urban areas in order to achieve the limit values (for SO<sub>2</sub>, PM<sub>10</sub>, NO<sub>2</sub>) and target values (for O<sub>3</sub>) set in the Decree?*

## Key message

### **Suspended particulate matters sized up to 10 micrometers (PM10)**

Concentrations of suspended particulate matters sized up to 10 micrometers exceed limit values specified in the Decree at the analyzed measuring points. High concentrations of suspended particulate matters sized up to 10 micrometers occur frequently, especially during winter period. Highest concentrations of PM10 have been measured in Skopje.

### **Nitrogen dioxide (NO<sub>2</sub>)**

In the period 2004 to 2013, decreasing trend was observed for the concentrations of nitrogen dioxide. Measured concentrations of this pollutant exceed specified limit value only in Skopje.


### **Sulphur dioxide (SO<sub>2</sub>)**

In the period 2004 to 2013, decreasing trend was observed for the mean annual concentration at all measuring points. Also, the mean daily limit value for sulphur dioxide was not exceeded, i.e. population was not exposed at concentrations of sulphur dioxide above the limit value at all analyzed measuring points.


### **Ozone (O<sub>3</sub>)**

In the period 2004 to 2013, the highest concentrations of ozone were recorded in Bitola, due to the fact that the city is situated in southwestern part of the country with higher number of sunny days during the year. Ozone concentrations measured in Skopje throughout the analyzed period were significantly lower compared to ozone concentrations measured in Bitola and Veles.


**Diagram 1: Mean annual concentration of PM10**




**Diagram 2: Mean annual concentration of NO<sub>2</sub>**



**Diagram 3: Mean annual concentration of SO<sub>2</sub>**



**Diagram 4: Mean annual concentration of O<sub>3</sub>**



**Data coverage:** excel

**Source:** Ministry of Environment and Physical Planning

## Assessment

### *Suspended particulate matters (PM<sub>10</sub>)*

Suspended particulate matters of size up to 10 micrometers are particles able to pass through an opening conducting selection by size, with 50% loss in efficiency at aerodynamic diameter of size less than ten micrometers (10 µm). These particles of size not exceeding 10 micrometers are the so called fine particles or aerosols. Their retention time in the air is long and they originate from natural and anthropogenic sources. Among natural sources, the more prominent include yellow rains, present also with us, forest fires and chemical reactions going on in nature. Combustion of coal, wood and oil,

industrial processes, transport and waste burning are the most significant anthropogenic sources.

Increased concentrations of suspended particulate matters can be recorded in urban areas, especially in autumn-winter seasons, which is most probably due to increased frequency in traffic, fossil fuels combustion and meteorological conditions.

Based on data processed for the period 2004-2013, we may conclude that throughout the period, the population was exposed at concentrations of suspended particulate matters exceeding limit values (mean limit value of  $50 \mu\text{g}/\text{m}^3$  which should not be exceeded in more than 35 days during a calendar year and annual limit value of  $40 \mu\text{g}/\text{m}^3$ ). Highest concentrations of this pollutant were recorded in Skopje, which was most probably due to human lifestyle, dense population, intensive use of solid fuel for households heating during winter, as well as impacts of industry.

### ***Nitrogen dioxide (NO<sub>2</sub>)***

Investigations have testified the presence of several nitrogen oxides in the air, but the most significant among them are nitrogen dioxide and nitrogen monoxide. These pollutants most often originate from natural sources. However, in urban environments, the main source is the traffic, and industry is minor source. The most toxic of all nitrogen oxides is the nitrogen dioxide, the concentrations of which are dependent on season and meteorological conditions. Namely, concentration of NO is higher in morning hours when the traffic is more frequent, while the intensification of solar radiation during the day leads to transformation of NO into NO<sub>2</sub> resulting in increased concentration of NO<sub>2</sub>. Nitrogen oxides influence the content of ozone and other photochemical oxidants in the air. During the spring-summer period, the concentration of NO<sub>2</sub> is higher, while in autumn-winter period, the concentration of NO is higher. The quantity of NO<sub>x</sub> increases in winter period due to the higher frequency of traffic.

Data processed indicate that the mean annual concentration of nitrogen dioxide was exceeded only in Skopje, which was most probably due to high frequency of traffic and operation of industrial facilities.

### ***Sulphur dioxide (SO<sub>2</sub>)***

Sulphur dioxide in the air most frequently originates from major thermal power plants, as well as from small and medium size boilers for coal combustion in urban environments. The main anthropogenic sources include coal and oil combustion. This pollutant is also released in the air from industrial processes (production of cellulose and paper, sulphuric acid, lead and zinc ores smelting).

In the period 2004 to 2013, there was no exceeding of the mean daily limit value for sulphur dioxide, i.e. population was not exceeded at concentrations of sulphur dioxide above the limit value at all analyzed measuring points.

### ***Ozone - O<sub>3</sub>***

Ozone layer is positioned at height of 10 km to 15 km from Earth and it plays the role of a filter for UV radiation and climate stabilizer.

Automatic monitoring stations measure the ground-level ozone formed as a result of photochemical reactions involving nitrogen oxides, volatile organic compounds (most frequently hydrocarbons), etc. However, its content is also dependent on solar radiation and annual seasons. Thus, higher ozone concentrations are observed in warmer days, especially during summer.

In the period 2004 to 2013, the highest concentrations of ozone were recorded in Bitola, due to the fact that the city is situated in southwestern part of the country with higher number of sunny days during the year. Ozone concentrations measured in Skopje throughout the analyzed period were significantly lower compared to ozone concentrations measured in Bitola and Veles.

## Methodology

- Methodology for the indicator calculation

Calculation of the indicator takes into account data from conducted measurements of the quality of the air in the following three cities: Skopje, Bitola and Veles, as the most representative sites reflecting the state of air quality in the Republic of Macedonia. Skopje was selected as capitol and the largest urban area in the country and big industrial centre. Bitola is second in size city and the largest thermal power plant REK Bitola is located in its vicinity and Veles as city with great historical industrial pollution.

For each selected urban station, the type of station is specified (urban background, station measuring air pollution from traffic, station measuring air pollution from industry). Then, the following is calculated for each pollutant from the available hourly data: mean annual concentration, highest mean daily concentration and number of days with exceeded mean daily limit value.

### Uncertainty

- Methodological uncertainty and data uncertainty

Calculations of the indicator have also taken into consideration stations where data coverage is below 75% per calendar year.

## Policy relevance of the indicator

### List of relevant policy documents

The National Plan for Air Protection presents the state of air quality, defines the measures for ambient air quality protection and improvement in the Republic of Macedonia and all relevant institutions responsible for their implementation within 5 year period, namely from 2013 to 2018 (Official Gazette of the Republic of Macedonia no.170/2012).

### Legal grounds

The Law on Ambient Air Quality was adopted in August 2004 and later amended on several occasions in line with the requirements of the relevant EU legislation (Official Gazette of the Republic of Macedonia Nos. 67/2004, 92/2007, 83/2009, 35/2010, 47/2011 and 59/2012) and it is framework law in the area of air. The main goals of this Law are: avoidance, prevention and reduction of harmful effects on human health and environment as a whole, prevention and reduction of pollution resulting in climate change, as well as provision of the relevant information on the quality of ambient air. This Law establishes the legal grounds for adoption of a number of bylaws in line with the requirements of the relevant *Acquis Communautaire*. So far, 12 bylaws have been adopted. Calculations for this indicator are based on the provisions of the Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term ozone targets (Official Gazette of the Republic of Macedonia No. 50/05, 4/2013).

## Targets

The Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term targets, defines the limit values for SO<sub>2</sub>, PM<sub>10</sub>, NO<sub>2</sub> and target values for O<sub>3</sub>.

### Limit values for concentrations of sulphur dioxide in ambient air

In accordance with the said Decree, two limit values are specified for sulphur dioxide for the purpose of

human health protection.

- Mean daily limit value of 125 µg/m<sup>3</sup> which shall not be exceeded by more than three times during one calendar year
- Hourly limit value of 350 µg/m<sup>3</sup>, which shall not be exceeded by more than 24 times during one calendar year.

#### Limit values for concentrations of nitrogen dioxide in ambient air

In accordance with the said Decree, two limit values are specified for nitrogen dioxide for the purpose of human health protection.

- Hourly mean concentration of nitrogen dioxide shall not exceed the limit value of 200 µg/m<sup>3</sup> by more than 18 times during one calendar year.
- The mean annual concentration shall not exceed 40 µg/m<sup>3</sup>.

#### Limit values for concentrations of suspended particulate matter of size up to 10 micrometers in the ambient air

The said Decree specifies two limit values for suspended particulate matter of size up to 10 micrometers, for the purpose of human health protection.

- 24-hourly limit value is 50 µg/m<sup>3</sup>, and it shall not be exceeded by more than 35 times during one calendar year
- The mean annual concentration shall not exceed 40 µg/m<sup>3</sup>.

#### Target values for ozone concentrations in ambient air

The said Decree, with regard to ozone, specifies target value for the purpose of human health protection and long-term target for the purpose of human health protection.

- The target value for ozone, for the purpose of human health protection, is specified so that 8-hourly mean value is calculated from the hourly concentrations in each day. The maximum daily 8-hourly mean value of ozone shall not exceed the value of 120 µg/m<sup>3</sup> in more than 25 days in the course of the year (calculated as an average value for three years). This target value should be achieved by 2010.
- The Decree also defines long-term target for the purpose of human health protection, set at 120 µg/m<sup>3</sup>, as maximum daily 8-hourly mean value during a calendar year.

## Reporting obligation

European Environmental Agency


- Air quality data exchange in accordance with implementing Decision containing the rules of Directives 2004/10/EC and 2008/50/EC of the European Parliament and of the Council concerning reciprocal exchange of information I reporting on ambient air quality (Decision 2011/850/EC).

## General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 004	Exceedance of air quality limit values in urban areas – by monitoring station	CSI 004	Exceedance of air quality limit values in urban areas	S	A	air Air quality	annual

## **MK - NI 004**

# EXCEEDANCE OF AIR QUALITY LIMIT VALUES IN URBAN AREAS - SO<sub>2</sub>



## Definition

The indicator shows the portion of urban population potentially exposed at ambient air concentrations of pollutants in excess of the limit value set for human health protection.

Urban population taken into account is actually the total number of inhabitants living in cities with at least one monitoring station. These cities include the capital and other major cities of the Republic of Macedonia. The number of inhabitants is based on the last census carried out by the State Statistical Office in 2002.

Exceedance of air quality limit values occurs when the concentration of air pollutants exceeds the limit values for SO<sub>2</sub>, PM<sub>10</sub>, NO<sub>2</sub> and the target values for O<sub>3</sub> as specified in the Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term ozone targets (Official Gazette of the Republic of Macedonia No. 50/05, 4/2013), wherein the requirements of the Directive on Ambient Air Quality and Cleaner Air in Europe 2008/50 EC and Heavy Metals Directive 2004/107/EC have been transposed. Where there are multiple limit values (see section on Policy Targets), the indicator uses the most stringent case:

- Sulphur dioxide (SO<sub>2</sub>): the daily mean limit value
- Nitrogen dioxide (NO<sub>2</sub>): the annual limit value
- Particulate matter of a size up to 10 micrometer (PM<sub>10</sub>): the daily mean limit value
- Ozone (O<sub>3</sub>): the short term objective

## Units

The percentage of urban population potentially exposed at ambient air concentrations of sulphur dioxide (SO<sub>2</sub>), particulate matter sized up to 10 micrometer (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) above limit values set for human health protection. Ambient air concentrations of sulphur dioxide (SO<sub>2</sub>), particulate matter sized up to 10 micrometer (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) are expressed in microgram/m<sup>3</sup>(µg/m<sup>3</sup>).

## Key policy issue

***What progress has been achieved in reducing the concentrations of pollutants in urban areas in order to achieve the limit values for SO<sub>2</sub> in the Decree?***

## Key message

No excess of mean daily concentrations of sulphur dioxide was recorded in the period from 2004 to 2013, i.e. the population was not exposed at sulphur dioxide concentrations above limit value, except in 2006 when out of the allowed 3 days, exceedance of the limit value was recorded in the course of 8 days in Skopje, which was not seen as significant problem.



Diagram 1: Percentage of urban population exposed at air pollution in areas where concentrations of pollutants are in excess of limit/target values




Diagram 2: Percentage of urban population exposed at concentrations of SO2 above the daily mean limit value expressed as number of days in the course of a calendar year





Diagram 3: 4th highest average mean daily concentration of SO<sub>2</sub>



Data coverage: [excel](#)

Source: Ministry of Environment and Physical Planning

## Assessment

Sulphur dioxide in the air most frequently originates from major thermal power plants, as well as from small and medium size boilers for coal combustion in urban environments. The main anthropogenic sources include coal and oil combustion. This pollutant is also released in the air from industrial processes (production of cellulose and paper, sulphuric acid, lead and zinc ores smelting).

In the period 2004 to 2013, there have been no concentrations above the daily mean limit value for sulphur dioxide, i.e. population was not exposed at sulphur dioxide concentrations above the limit value, except in 2006 when out of the allowed 3 days, exceedance of the limit value was recorded in the course of 8 days in Skopje, which was not seen as significant problem.

In 2006, 53% of the population was exposed at sulphur dioxide concentration above 125 µg/m<sup>3</sup> for more than 6 days in the course of the year. In 2005 and 2007, there was higher percentage (around 50%) of the population exposed at sulphur dioxide concentration above 125 µg/m<sup>3</sup> for 1 to 3 days in the course of the year, while in 2008 and 2009, this percentage of population exposure was very low (3%).

## Methodology

- Methodology for the indicator calculation

For each measuring station located in urban environment, the number of days with mean daily concentration higher than the limit value (daily mean value of 125 µg/m<sup>3</sup>) is calculated from the available hourly data. Selected urban stations include stations of the following types: stations measuring traffic pollution, stations measuring industrial pollution and urban background stations. The number of days with excess in a city is obtained by averaging the results of all stations located in that city.

## **Uncertainty**

- **Methodological uncertainty and data uncertainty**

In general, data is not representative for all urban environments in the Republic of Macedonia. Compared to the methodology of the European Environmental Agency, where the calculation of the indicator is based only on data produced by the urban background stations, in our calculations we used data from all measuring stations located in urban environments. Also, due to the minimum number of monitoring stations, the calculation of the indicator also took into account the stations where data coverage is below 75% per calendar year. We can also point out as uncertainty in the indicator calculation the fact that the number of inhabitants in cities is based on the census of the population conducted by the State Statistical Office in 2002, instead of estimated number of population for each year.

## **Policy relevance of the indicator**

### **List of relevant policy documents**

The National Plan for Air Protection presents the state of air quality, defines the measures for ambient air quality protection and improvement in the Republic of Macedonia and all relevant institutions responsible for their implementation within 5 year period, namely from 2013 to 2018 (Official Gazette of the Republic of Macedonia no.170/2012).

### **Legal grounds**

The Law on Ambient Air Quality was adopted in August 2004 and later amended on several occasions in line with the requirements of the relevant EU legislation (Official Gazette of the Republic of Macedonia Nos. 67/2004, 92/2007, 83/2009, 35/2010, 47/2011 and 59/2012) and it is framework law in the area of air. The main goals of this Law are: avoidance, prevention and reduction of harmful effects on human health and environment as a whole, prevention and reduction of pollution resulting in climate change, as well as provision of the relevant information on the quality of ambient air. This Law establishes the legal grounds for adoption of a number of bylaws in line with the requirements of the relevant *Acquis Communautaire*. So far, 12 bylaws have been adopted. Calculations for this indicator are based on the provisions of the Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term ozone targets (Official Gazette of the Republic of Macedonia No. 50/05, 4/2013).

## **Targets**

The Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term targets, defines the limit values for SO<sub>2</sub>, PM<sub>10</sub>, NO<sub>2</sub> and target values for O<sub>3</sub>.

### **Limit values for concentrations of sulphur dioxide in ambient air**

In accordance with the said Decree, two limit values are specified for sulphur dioxide for the purpose of human health protection.

- Mean daily limit value of 125 µg/m<sup>3</sup> which shall not be exceeded by more than three times during one calendar year
- Hourly limit value of 350 µg/m<sup>3</sup>, which shall not be exceeded by more than 24 times during one calendar year.

## Reporting obligation

European Environmental Agency


- Air quality data exchange in accordance with implementing Decision containing the rules of Directives 2004/10/EC and 2008/50/EC of the European Parliament and of the Council concerning reciprocal exchange of information I reporting on ambient air quality (Decision 2011/850/EC).

## General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 004</b>	<b>Exceedance of air quality limit values in urban areas</b>	CSI 004	Exceedance of air quality limit values in urban areas	<b>S</b>	<b>A</b>	air Air quality	annual

## MK - NI 004

# EXCEEDANCE OF AIR QUALITY LIMIT VALUES IN URBAN AREAS - NO<sub>2</sub>



## Definition

The indicator shows the portion of urban population potentially exposed at ambient air concentrations of pollutants in excess of the limit value set for human health protection.

Urban population taken into account is actually the total number of inhabitants living in cities with at least one monitoring station. These cities include the capital and other major cities of the Republic of Macedonia. The number of inhabitants is based on the last census carried out by the State Statistical Office in 2002.

Exceedance of air quality limit values occurs when the concentration of air pollutants exceeds the limit values for SO<sub>2</sub>, PM<sub>10</sub>, NO<sub>2</sub> and the target values for O<sub>3</sub> as specified in the Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term ozone targets (Official Gazette of the Republic of Macedonia No. 50/05, 4/2013), wherein the requirements of the Directive on Ambient Air Quality and Cleaner Air in Europe 2008/50 EC and Heavy Metals Directive 2004/107/EC have been transposed. Where there are multiple limit values (see section on Policy Targets), the indicator uses the most stringent case:

- Sulphur dioxide (SO<sub>2</sub>): the daily mean limit value
- Nitrogen dioxide (NO<sub>2</sub>): the annual limit value
- Particulate matter of a size up to 10 micrometer (PM<sub>10</sub>): the daily mean limit value
- Ozone (O<sub>3</sub>): the short term objective

## Units

The percentage of urban population potentially exposed at ambient air concentrations of sulphur dioxide (SO<sub>2</sub>), particulate matter sized up to 10 micrometer (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) above limit values set for human health protection. Ambient air concentrations of sulphur dioxide (SO<sub>2</sub>), particulate matter sized up to 10 micrometer (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) are expressed in microgram/m<sup>3</sup>(µg/m<sup>3</sup>).

## Key policy issue

***What progress has been achieved in reducing the concentrations of pollutants in urban areas in order to achieve the limit values, NO<sub>2</sub> set in the Decree?***

## Key message

In the period from 2004 to 2013, the portion of the population exposed at concentrations of nitrogen dioxide above the limit values set for human health protection ranged between 0 and 69%. The highest percentage of population exposure of 69% was recorded in 2011.

Diagram 1: Percentage of urban population exposed at air pollution in areas where concentrations of pollutants are in excess of limit/target values




Diagram 2: Percentage of population exposed at NO2 annual concentrations in urban areas





Diagram 3: Average annual concentration of NO<sub>2</sub>



Data coverage: [excel](#)

Source: Ministry of Environment and Physical Planning

## Assessment

Investigations have testified the presence of several nitrogen oxides in the air, but the most significant among them are nitrogen dioxide and nitrogen monoxide. These pollutants most often originate from natural sources. However, in urban environments, the main source is the traffic, and industry is minor source. The most toxic of all nitrogen oxides is the nitrogen dioxide, the concentrations of which are dependent on season and meteorological conditions. Namely, concentration of NO is higher in morning hours when the traffic is more frequent, while the intensification of solar radiation during the day leads to transformation of NO into NO<sub>2</sub> resulting in increased concentration of NO<sub>2</sub>. Nitrogen oxides influence the content of ozone and other photochemical oxidants in the air. During the spring-summer period, the concentration of NO<sub>2</sub> is higher, while in autumn-winter period, the concentration of NO is higher. The quantity of NO<sub>x</sub> increases in winter period due to the higher frequency of traffic.

In the period 2004 to 2013, the portion of population exposed at nitrogen dioxide concentrations above the limit value for human health protection (40 µg/m<sup>3</sup> mean annual limit value) ranged from 0 to 69%. The highest percentage of 69% population exposure was recorded in 2011.

In 2004, 2006 and 2011, significant portion of the population (55-67%) of the population was exposed at concentrations above 40 µg/m<sup>3</sup>.

## Methodology

### ▪ Methodology for the indicator calculation

The mean annual concentration in a city is calculated as an average of the mean annual value measured in all monitoring stations located in urban areas. Selected urban stations include stations of the following types: stations measuring traffic pollution, stations measuring industrial pollution and urban background stations.

## **Uncertainty**

- **Methodological uncertainty and data uncertainty**

In general, data is not representative for all urban environments in the Republic of Macedonia. Compared to the methodology of the European Environmental Agency, where the calculation of the indicator is based only on data produced by the urban background stations, in our calculations we used data from all measuring stations located in urban environments. Also, due to the minimum number of monitoring stations, the calculation of the indicator also took into account the stations where data coverage is below 75% per calendar year. We can also point out as uncertainty in the indicator calculation the fact that the number of inhabitants in cities is based on the census of the population conducted by the State Statistical Office in 2002, instead of estimated number of population for each year.

## **Policy relevance of the indicator**

### **List of relevant policy documents**

The National Plan for Air Protection presents the state of air quality, defines the measures for ambient air quality protection and improvement in the Republic of Macedonia and all relevant institutions responsible for their implementation within 5 year period, namely from 2013 to 2018 (Official Gazette of the Republic of Macedonia no.170/2012).

### **Legal grounds**

The Law on Ambient Air Quality was adopted in August 2004 and later amended on several occasions in line with the requirements of the relevant EU legislation (Official Gazette of the Republic of Macedonia Nos. 67/2004, 92/2007, 83/2009, 35/2010, 47/2011 and 59/2012) and it is framework law in the area of air. The main goals of this Law are: avoidance, prevention and reduction of harmful effects on human health and environment as a whole, prevention and reduction of pollution resulting in climate change, as well as provision of the relevant information on the quality of ambient air. This Law establishes the legal grounds for adoption of a number of bylaws in line with the requirements of the relevant *Acquis Communautaire*. So far, 12 bylaws have been adopted. Calculations for this indicator are based on the provisions of the Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term ozone targets (Official Gazette of the Republic of Macedonia No. 50/05, 4/2013).

## **Targets**

The Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term targets, defines the limit values for NO<sub>2</sub>.

### **Limit values for concentrations of nitrogen dioxide in ambient air**

In accordance with the said Decree, two limit values are specified for nitrogen dioxide for the purpose of human health protection.

- Hourly mean concentration of nitrogen dioxide shall not exceed the limit value of 200 µg/m<sup>3</sup> by more than 18 times during one calendar year.
- The mean annual concentration shall not exceed 40 µg/m<sup>3</sup>.



## Reporting obligation

European Environmental Agency

- Air quality data exchange in accordance with implementing Decision containing the rules of Directives 2004/10/EC and 2008/50/EC of the European Parliament and of the Council concerning reciprocal exchange of information I reporting on ambient air quality (Decision 2011/850/EC).

## General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 004</b>	<b>Exceedance of air quality limit values in urban areas</b>	CSI 004	Exceedance of air quality limit values in urban areas	<b>S</b>	<b>A</b>	air Air quality	annual

## **MK - NI 004**

# EXCEEDANCE OF AIR QUALITY LIMIT VALUES IN URBAN AREAS - PM<sub>10</sub>



## Definition

The indicator shows the portion of urban population potentially exposed at ambient air concentrations of pollutants in excess of the limit value set for human health protection.

Urban population taken into account is actually the total number of inhabitants living in cities with at least one monitoring station. These cities include the capital and other major cities of the Republic of Macedonia. The number of inhabitants is based on the last census carried out by the State Statistical Office in 2002.

Exceedance of air quality limit values occurs when the concentration of air pollutants exceeds the limit values for SO<sub>2</sub>, PM<sub>10</sub>, NO<sub>2</sub> and the target values for O<sub>3</sub> as specified in the Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term ozone targets (Official Gazette of the Republic of Macedonia No. 50/05, 4/2013), wherein the requirements of the Directive on Ambient Air Quality and Cleaner Air in Europe 2008/50 EC and Heavy Metals Directive 2004/107/EC have been transposed. Where there are multiple limit values (see section on Policy Targets), the indicator uses the most stringent case:

- Sulphur dioxide (SO<sub>2</sub>): the daily mean limit value
- Nitrogen dioxide (NO<sub>2</sub>): the annual limit value
- Particulate matter of a size up to 10 micrometer (PM<sub>10</sub>): the daily mean limit value
- Ozone (O<sub>3</sub>): the short term objective

## Units

The percentage of urban population potentially exposed at ambient air concentrations of sulphur dioxide (SO<sub>2</sub>), particulate matter sized up to 10 micrometer (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) above limit values set for human health protection. Ambient air concentrations of sulphur dioxide (SO<sub>2</sub>), particulate matter sized up to 10 micrometer (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) are expressed in microgram/m<sup>3</sup>(µg/m<sup>3</sup>).

## Key policy issue

***What progress has been achieved in reducing the concentrations of pollutants in urban areas in order to achieve the limit values for PM<sub>10</sub> set in the Decree?***

## Key message

In the period from 2004 to 2013, 100% of the population has been exposed at concentrations of suspended particulate matter in excess of the limit values specified in the Decree. Significantly higher concentrations of PM<sub>10</sub> are measured during winter period.

Diagram 1: Percentage of urban population exposed at air pollution in areas where concentrations of pollutants are in excess of limit/target values




Diagram 2: Percentage of urban population exposed at concentrations of PM10 above the daily mean limit value, expressed as number of days in the course of a calendar year





Diagram 3: 36th highest mean daily concentration of PM10



Data coverage: [excel](#)

Source: Ministry of Environment and Physical Planning

## Assessment

Suspended particulate matters of size up to 10 micrometers are particles able to pass through an opening conducting selection by size, with 50% loss in efficiency at aerodynamic diameter of size less than ten micrometers (10  $\mu\text{m}$ ). These particles of size not exceeding 10 micrometers are the so called fine particles or aerosols. Their retention time in the air is long and they originate from natural and anthropogenic sources. Among natural sources, the more prominent include yellow rains, present also with us, forest fires and chemical reactions going on in nature. Combustion of coal, wood and oil, industrial processes, transport and waste burning are the most significant anthropogenic sources.

Increased concentrations of suspended particulate matters can be recorded in urban areas, especially in autumn-winter seasons, which is most probably due to increased frequency in traffic, fossil fuels combustion and meteorological conditions.

The processed data for the period 2004 to 2013 show that during the entire period, 100% of the population was exposed at concentrations of suspended particulate matters are in excess of the limit value (mean daily limit value of 50  $\mu\text{g}/\text{m}^3$  that shall not be exceeded in more than 35 days in the course of a calendar year). 100% of the population was exposed at concentrations above the limit value in more than 35 days in a calendar year.

## Methodology

### ▪ Methodology for the indicator calculation

For each urban measuring station, the number of days with daily mean concentration above the limit value (daily mean limit value is 50  $\mu\text{g}/\text{m}^3$ ) is calculated from the available hourly data. Selected urban stations include stations of the following types: stations measuring traffic pollution, stations measuring industrial pollution and urban background stations. The number of days with exceedance in a city is obtained by averaging the mean values of the results from all stations located in that city.

## Uncertainty

- Methodological uncertainty and data uncertainty

In general, data is not representative for all urban environments in the Republic of Macedonia. Compared to the methodology of the European Environmental Agency, where the calculation of the indicator is based only on data produced by the urban background stations, in our calculations we used data from all measuring stations located in urban environments. Also, due to the minimum number of monitoring stations, the calculation of the indicator also took into account the stations where data coverage is below 75% per calendar year. We can also point out as uncertainty in the indicator calculation the fact that the number of inhabitants in cities is based on the census of the population conducted by the State Statistical Office in 2002, instead of estimated number of population for each year.

## Policy relevance of the indicator

### List of relevant policy documents

The National Plan for Air Protection presents the state of air quality, defines the measures for ambient air quality protection and improvement in the Republic of Macedonia and all relevant institutions responsible for their implementation within 5 year period, namely from 2013 to 2018 (Official Gazette of the Republic of Macedonia no.170/2012).

### Legal grounds

The Law on Ambient Air Quality was adopted in August 2004 and later amended on several occasions in line with the requirements of the relevant EU legislation (Official Gazette of the Republic of Macedonia Nos. 67/2004, 92/2007, 83/2009, 35/2010, 47/2011 and 59/2012) and it is framework law in the area of air. The main goals of this Law are: avoidance, prevention and reduction of harmful effects on human health and environment as a whole, prevention and reduction of pollution resulting in climate change, as well as provision of the relevant information on the quality of ambient air. This Law establishes the legal grounds for adoption of a number of bylaws in line with the requirements of the relevant *Acquis Communautaire*. So far, 12 bylaws have been adopted. Calculations for this indicator are based on the provisions of the Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term ozone targets (Official Gazette of the Republic of Macedonia No. 50/05, 4/2013).

## Targets

The Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term targets, defines the limit values for SO<sub>2</sub>, PM<sub>10</sub>, NO<sub>2</sub> and target values for O<sub>3</sub>.

### Limit values for concentrations of suspended particulate matter of size up to 10 micrometers in the ambient air

The said Decree specifies two limit values for suspended particulate matter of size up to 10 micrometers, for the purpose of human health protection.

- 24-hourly limit value is 50 µg/m<sup>3</sup>, and it shall not be exceeded by more than 35 times during one calendar year
- The mean annual concentration shall not exceed 40 µg/m<sup>3</sup>.

## Reporting obligation

European Environmental Agency


- Air quality data exchange in accordance with implementing Decision containing the rules of Directives 2004/10/EC and 2008/50/EC of the European Parliament and of the Council concerning reciprocal exchange of information I reporting on ambient air quality (Decision 2011/850/EC).

## General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 004</b>	<b>Exceedance of air quality limit values in urban areas</b>	CSI 004	Exceedance of air quality limit values in urban areas	<b>S</b>	<b>A</b>	air Air quality	annual

## MK - NI 004

# EXCEEDANCE OF AIR QUALITY LIMIT VALUES IN URBAN AREAS - O<sub>3</sub>



## Definition

The indicator shows the portion of urban population potentially exposed at ambient air concentrations of pollutants in excess of the limit value set for human health protection.

Urban population taken into account is actually the total number of inhabitants living in cities with at least one monitoring station. These cities include the capital and other major cities of the Republic of Macedonia. The number of inhabitants is based on the last census carried out by the State Statistical Office in 2002.

Exceedance of air quality limit values occurs when the concentration of air pollutants exceeds the limit values for SO<sub>2</sub>, PM<sub>10</sub>, NO<sub>2</sub> and the target values for O<sub>3</sub> as specified in the Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term ozone targets (Official Gazette of the Republic of Macedonia No. 50/05, 4/2013), wherein the requirements of the Directive on Ambient Air Quality and Cleaner Air in Europe 2008/50 EC and Heavy Metals Directive 2004/107/EC have been transposed. Where there are multiple limit values (see section on Policy Targets), the indicator uses the most stringent case:

- Sulphur dioxide (SO<sub>2</sub>): the daily mean limit value
- Nitrogen dioxide (NO<sub>2</sub>): the annual limit value
- Particulate matter of a size up to 10 micrometer (PM<sub>10</sub>): the daily mean limit value
- Ozone (O<sub>3</sub>): the short term objective

## Units

The percentage of urban population potentially exposed at ambient air concentrations of sulphur dioxide (SO<sub>2</sub>), particulate matter sized up to 10 micrometer (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) above limit values set for human health protection. Ambient air concentrations of sulphur dioxide (SO<sub>2</sub>), particulate matter sized up to 10 micrometer (PM<sub>10</sub>), nitrogen dioxide (NO<sub>2</sub>) and ozone (O<sub>3</sub>) are expressed in microgram/m<sup>3</sup>(µg/m<sup>3</sup>).

## Key policy issue

***What progress has been achieved in reducing the concentrations of pollutants in urban areas in order to achieve the target values for O<sub>3</sub> set in the Decree?***

## Key message

In the period from 2004 to 2013, the portion of the population exposed at concentrations of ozone higher than the target value set for human health protection ranged from 12 to 43%. The highest percentage of population exposure of 43% was recorded in 2007.

Diagram 1: Percentage of urban population exposed at air pollution in areas where concentrations of pollutants are in excess of limit/target values




Diagram 2: Percentage of urban population exposed at concentrations of O3 above the long-term target value for human health protection, expressed as number of days in the course of a calendar year







Diagram 3: 26th highest maximum 8-hourly mean concentration of O<sub>3</sub>



Data coverage: [excel](#)

Source: Ministry of Environment and Physical Planning

## Assessment

Ozone layer is positioned at height of 10 km to 15 km from Earth and it plays the role of a filter for UV radiation and climate stabilizer.

Automatic monitoring stations measure the ground-level ozone formed as a result of photochemical reactions involving nitrogen oxides, volatile organic compounds (most frequently hydrocarbons), etc. However, its content is also dependent on solar radiation and annual seasons. Thus, higher ozone concentrations are observed in warmer days, especially during summer.

In the period 2004 to 2013, the portion of population exposed at ozone concentrations above the target value for human health protection ranged from 0 to 43%. The highest percentage of 43% population exposure was recorded in 2007, followed by falling trend to reach 0% in 2012, while the percentage of exposure in 2013 was 15%.

In the period 2004 to 2009, population was exposed at concentrations above the target value of 120 µg/m<sup>3</sup> in more than 25 to 50 days in a calendar year. Percentage of exposure at concentrations above the target value of 120 µg/m<sup>3</sup> in more than 25 to 50 days in a calendar year was recorded during the entire reporting period, except in 2005 and 2012.

## Methodology

For each measuring station located in urban environment, the number of days in which the maximum daily 8-hourly mean concentration of ozone is in excess of ozone target value for human health protection - 120 µg/m<sup>3</sup> is calculated. Selected urban stations include stations of the following types: stations measuring traffic pollution, stations measuring industrial pollution and urban background stations. The number of days with excess in a city is obtained by averaging the results of all stations located in that city.

## **Uncertainty**

- **Methodological uncertainty and data uncertainty**

In general, data is not representative for all urban environments in the Republic of Macedonia. Compared to the methodology of the European Environmental Agency, where the calculation of the indicator is based only on data produced by the urban background stations, in our calculations we used data from all measuring stations located in urban environments. Also, due to the minimum number of monitoring stations, the calculation of the indicator also took into account the stations where data coverage is below 75% per calendar year. We can also point out as uncertainty in the indicator calculation the fact that the number of inhabitants in cities is based on the census of the population conducted by the State Statistical Office in 2002, instead of estimated number of population for each year.

## **Policy relevance of the indicator**

### **List of relevant policy documents**

The National Plan for Air Protection presents the state of air quality, defines the measures for ambient air quality protection and improvement in the Republic of Macedonia and all relevant institutions responsible for their implementation within 5 year period, namely from 2013 to 2018 (Official Gazette of the Republic of Macedonia no.170/2012).

### **Legal grounds**

The Law on Ambient Air Quality was adopted in August 2004 and later amended on several occasions in line with the requirements of the relevant EU legislation (Official Gazette of the Republic of Macedonia Nos. 67/2004, 92/2007, 83/2009, 35/2010, 47/2011 and 59/2012) and it is framework law in the area of air. The main goals of this Law are: avoidance, prevention and reduction of harmful effects on human health and environment as a whole, prevention and reduction of pollution resulting in climate change, as well as provision of the relevant information on the quality of ambient air. This Law establishes the legal grounds for adoption of a number of bylaws in line with the requirements of the relevant *Acquis Communautaire*. So far, 12 bylaws have been adopted. Calculations for this indicator are based on the provisions of the Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term ozone targets (Official Gazette of the Republic of Macedonia No. 50/05, 4/2013).

## **Targets**

The Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term targets, defines the limit values for SO<sub>2</sub>, PM<sub>10</sub>, NO<sub>2</sub> and target values for O<sub>3</sub>.

### **Target values for ozone concentrations in ambient air**

The said Decree, with regard to ozone, specifies target value for the purpose of human health protection and long-term target for the purpose of human health protection.

- The target value for ozone, for the purpose of human health protection, is specified so that 8-hourly mean value is calculated from the hourly concentrations in each day. The maximum daily 8-hourly mean value of ozone shall not exceed the value of 120 µg/m<sup>3</sup> in more than 25 days in the course of the year (calculated as an average value for three years). This target value should be achieved by 2010.
- The Decree also defines long-term target for the purpose of human health protection, set at 120 µg/m<sup>3</sup>, as maximum daily 8-hourly mean value during a calendar year.

## Reporting obligation

European Environmental Agency


- Air quality data exchange in accordance with implementing Decision containing the rules of Directives 2004/10/EC and 2008/50/EC of the European Parliament and of the Council concerning reciprocal exchange of information I reporting on ambient air quality (Decision 2011/850/EC).

## General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 004</b>	<b>Exceedance of air quality limit values in urban areas</b>	CSI 004	Exceedance of air quality limit values in urban areas	<b>S</b>	<b>A</b>	air Air quality	annual

## MK – NI 050

### EMISSION OF AMMONIA



#### Definition

The indicator tracks the trends in ammonia.

The indicator also provides information on emissions by subsectors in the key sector - agriculture.

#### Units

kt (kilotons per year)


#### Key policy issue

#### What progress has been made in overall reduction of emissions of polluting substances ozone precursors in Europe?

- Emission of ozone precursors in 2012 compared to 2004 dropped by around 18%, and compared to 2011 it dropped by 8%. In 2012, ammonia emissions amounted 10.82 Gg and thus the national ceiling for ammonia emission was not exceeded, which under Annex II to Gothenburg Protocol was 12 Gg.
- **Dropping trend in ammonia emissions during the analyzed period derived from reduction** in the number of bred livestock. Furthermore, the use of nitrogen artificial fertilizers contributing to emissions of this polluting substance was reduced. Other sectors, like passenger transport and vehicles recorded increase in ammonia emissions, but due to low share they had no significant effect on overall emissions on national level.

The Diagram below shows annual trend in ammonia emissions for the period 2004 to 2012.

Diagram 1. Trend in ammonia emissions



#### Assessment


Under the CORINAIR Programme, inventory of air pollutants emission in the country was established in 2005 by individual sectors, i.e. activities, and in 2012 Inventory including all pollutants was prepared.

Sectors based on the above mentioned methodology and SNAP – selective nomenclature are presented in the table below:

SNAP	Sector
1	Combustion in energy and transformation industries
2	Non-industrial combustion plants
3	Combustion in manufacturing industry
4	Production processes
5	Extraction and distribution of fossil fuels and geothermal energy
6	Solvent and other product use
7	Road transport
8	Other mobile sources and machinery
9	Waste treatment and disposal
10	Agriculture
11	Nature

In the overall emissions of ammonia in 2012, minimum share belonged to sectors 2, 7 and 9 with 0.22%, 0.24% and 2.53% respectively, while agriculture was the key sector with 97%.

Diagram 2. Emissions of NH<sub>3</sub> by SNAP sectors per year



## Policy specific issue


**Which different sectors and processes contribute to ammonia emissions?**

Agriculture is the key sector for ammonia emissions. Therefore, Diagram 2 shows emissions from agricultural sector by subsectors, where the main subsectors giving rise to the highest ammonia emissions, such as breeding of poultry, sheep, pigs and cattle are presented separately, while 'other'

summarizes emissions originating from breeding of horses, goats and other poultry, as well as emissions from SNAP sectors 2,7 and 9.

Emissions from livestock breeding originate from urea decomposition in the feces of animals and decomposition of urinary acid in poultry. Ammonia emissions depend on species of animals, their age, manner of breeding, and waste management and disposal. The main reason for emission reduction is the reduction in the number of animals.

Diagram 3. Emissions of NH<sub>3</sub> from agriculture by NFR sectors per year



Data coverage: [excel](#)

**Sources of data:** State Statistical Office, Energy Balance – Report of the Government, Cadastre of air polluters and pollutants, Data from the measurements of companies – major polluters, Database of motor vehicles of the Ministry of Interior, Inventory by EMEP/EEA Methodology of the Ministry of Environment and Physical Planning, Ministry of Agriculture, Forestry and Water Economy.

## Methodology

- Methodology for indicator calculation

The methodology for this indicator calculation is based on aggregation and calculation of data on ammonia emissions at annual basis, on national level, as overall and distributed to sectors, i.e. activities.

Calculations are in line with the Guidelines of UNECE/EMEP, i.e. methodology for inventory taking, as well as use of SNAP – selective nomenclature of air pollution.

## Reference of used methodology

Methodology used for calculation and presentation of this indicator is given in EMEP/EEA Guideline for inventory of air pollutant emissions 2013, Technical Report no. 12/2013, EEA and de Leeuw, F. (2002). Group of emission indicators of long-range transboundary air pollution. Science and policy of environment.

## Policy relevance of the indicator

Action Plan for European Partnership, as well as National Plan for approximation of the national legislation with European regulations specifying bylaws that need to be prepared has been adopted.

The National Environmental Action Plan (NEAP II) was adopted. It contains the measures that need to be taken to improve the overall status of air quality, including the reduction of emissions of acidifying substances. The National Plan for Ambient Air Protection for the period 2012 to 2017 and the National Programme for gradual air emissions reduction by 2020 have been adopted in order to define and implement measures on national level concerning reduction of emissions from agricultural sector which has the highest share in ammonia emissions.

In 2007, the Government of the Republic of Macedonia adopted the National Strategy for Agriculture and Rural Development, establishing the principles of the policies for support and measures adapted to expected changes in legislation, institutions and control systems. The Strategy defines strategic goal as basis for development of agricultural and rural sector in the Republic of Macedonia in the period from 2007 to 2013 reading: "to strengthen agriculture to be competitive on integrated regional markets of the European Union and Southeastern Europe through measures aimed at enhancing the efficiency of agricultural production, processing and placement and building of relevant effective public and private institutions; to improve revenues on farm; ensure that consumers have access to safe, healthy food; to optimize the benefit of limited soil, forest and water resources in environmentally favourable manner; and build vital rural communities through sustainable rural development". In 2007, the Government of the Republic of Macedonia adopted the National Strategy for Organic Agricultural Production (2008 - 2011) setting the grounds for introduction and development of organic agricultural production. This Strategy is accompanied with Action Plan of measures and activities for strategy implementation, which have been to a great extent implemented.

National Plan for Organic Production 2013-2020 was adopted. The purpose of the National Plan for Organic Production for the period 2013-2020 is the instrument providing basis for further development of the organic production in the Republic of Macedonia. This National Plan also sets the directions, activities and measures, namely policies to be implemented by MAFWE during the period 2013 -2020 concerning future development of organic production in the Republic of Macedonia and also provides basis for planning and implementation of financial support in this sector.

The Law on Agricultural Land stipulates measures for agricultural land fertility improvement through undertaking agritechnical measures, and one of these measures is application of mineral fertilizers. It is stated that protection of agricultural land against pollution and infection is carried out by prohibition, restriction and prevention of direct inlet of harmful matters in soil, water and air and undertaking of other measures to maintain and improve productivity. It is specified that protection of agricultural land from contamination and infection are subject of application of regulations for environment and nature protection and improvement.

The Law on Fertilizers regulates the manner of fertilizers use. This is especially important for ammonia emissions reduction from the use of nitrogen artificial; fertilizers.

As far as ammonia emissions from traffic are concerned, in accordance with the Strategy for Energy, renewal of vehicle fleet has been envisaged with expectation for use of vehicles with secondary catalysts with lower NH<sub>3</sub> emission levels in future.

## Targets

### **Does any of the national documents set target or target should be achieved in accordance with other international documents?**

National strategic documents listed as references in the above text provide guidelines and specify actions that should be undertaken as a matter of priority. It is important to mention that bylaws have been

prepared in the area of air emissions transposing Directives 96/61/EC, 2000/81/EC, 2000/76/EC, 99/13/EC and 2001/81/EC, ranging between 90 and 100%.

In accordance with the requirements of the UNECE Convention on Long-Range Transboundary Air Pollution, inventory based on EMEP/EEA Guideline for inventory of polluting substances into the air, setting the target of regular inventory of pollutants in tons per year following the n-2 principle, where n is the current year.

Also, in accordance with Directive 2001/81/EC, as well as Gothenburg Protocol, the ceilings of the amounts of emissions have been set at the level of the Republic of Macedonia for 2010 that shall not be exceeded at the annual level of 2010. The Executive Body of the Convention on Long-Range Transboundary Air Pollution, upon submission of the values of national ceilings in order to enrol the Republic of Macedonia in Annex II of the Gothenburg Protocol requested correction of the values considering the reported data on air emissions of the pollutants sulfur dioxide and ammonia at national level. Changes in the values of these pollutants were incorporated in the Rulebook amending the Rulebook on upper limits – emission ceilings of pollutants for the purpose of setting projections for certain period concerning reduction of the quantities of pollutant emissions at annual level published in July 2014. In 2012, national upper limits – ceiling for NO<sub>3</sub> was not exceeded

	Upper limit - ceiling	Total emissions-2012
NH <sub>3</sub>	12 Gg	10,81 Gg

With regard to targets – emission projections for NO<sub>3</sub> for 2020, it should be pointed out that they do not take into account emissions from nitrogen artificial fertilizers and therefore review of projections has been planned for the period 2015-2016 and these are not included in this report.

### Legal grounds

The Law on Ambient Air Quality adopted in August 2004 and amended several times afterwards (Official Gazette of RM no. 67/2004, 92/2007, 83/2009, 35/10, 47/11, 100/12, 163/2013) is framework law in the area of air. The goals of this Law include avoiding, prevention and reduction of harmful effects on human health and environment as a whole, prevention and abatement of pollutions leading to climate change, as well as provision of appropriate information on the quality of ambient air.

On the basis of the Law on Ambient Air Quality, 16 bylaws were prepared and adopted to introduce limit values for air quality and air emissions, methodology of air quality and air emissions monitoring, manner of preparation of planning documents for air protection against pollution, manner of informing the citizens and international organizations, etc.

With reference to air standards transposed in part of the mentioned rulebooks, 86 ISO and 48 CEN standards in the area of air emissions and air quality were adopted by means of endorsement method.

Other legislation related to the regulation of air quality and air emissions regulation includes the Law on Vehicles, Law on Standardization, Rulebook on liquid fuels quality with national standards for liquid fuels quality, etc

In 2007, the Law on Agriculture and Rural Development was adopted as basic horizontal act regulating areas of planning and implementation of agricultural and rural development policy, as well as other aspects of agricultural and rural development policy.

Amendments of the Law on Agriculture and Rural Development in 2010 further harmonized and integrated the principles of programming, monitoring and implementation of the policy of the Republic of Macedonia for agriculture and rural development with the European Common Agricultural Policy.



The Law includes provisions for programming and implementing the policy of state aid, as well as provisions for application of the measures for rural development.

The framework of organic agricultural production is defined in the Law on Organic Agricultural Production. A number of bylaws were adopted under this Law in 2010, regulating the areas of production, processing, labeling of organic products, authorization and certification, as well as control systems. Also, the Law on Agriculture Development Fostering, the Law on Environment and the Law on Nature Protection are of relevance.

The above documents provide basis for achievement of the targets for emission reduction of pollutants that are ozone precursors, accompanied with reduced degradation of environment and negative effects on human health.

## Reporting obligation

- Reporting obligation towards UNECE-CLRTAP and EEA
- Annual Report of Processed Data on Air Emissions

## General metadata


Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 050</b>	<b>Emissions of ammonia</b>	APE 003	Ammonia (NH <sub>3</sub> ) emissions	<b>P</b>	<b>A</b>	<ul style="list-style-type: none"> <li>▪ air</li> <li>▪ quality of air</li> </ul>	annually

# NATURE AND BIODIVERSITY



## MK – NI 007

### THREATENED AND PROTECTED SPECIES



#### Definition

This indicator shows the number of species present in the Republic of Macedonia and assessed as globally threatened and/or protected by European instruments, such as: Rio Convention on Biological Diversity, Bonn Convention on the Conservation of Migratory Species of Wild Animals, Hague Agreement on the Conservation of Migratory Waterbirds, London Agreement on Bats Protection, EU Directives on habitats and wild birds and Bern Convention on the Conservation of European Wildlife and Natural Habitats, protected at national level.

At present, the indicator shows the status of the number of endemic and threatened species at national level, identified in accordance with the relevant international documents and the national legislation:

- Number of endemic and threatened wild species of plants (flora)
- Number of endemic and threatened wild species of native fungi
- Number of endemic and threatened wild species of animals (fauna)

#### Units

- Number of species

#### Key policy issue

***How many species of global/ European significance are protected by national instruments?***

#### Key message

Abundance and variety of ecosystems, types of natural habitats and wild species, as well as genetic resources, are the main features of biological diversity in the Republic of Macedonia. According to the available information, this wealth comprises the imposing number of around 17.604 species, out of which 976 species are endemic species.

Considering the fact that national Red Lists of animals, plants and fungi are under establishment, the analysis of threatened and protected wild species has been made in accordance with international criteria contained in a number of multilateral documents (conventions, agreements, Global Red List, European Red List, EU Directives).

Thus, the IUCN World Red List contains 72 taxa of higher plants from the Republic of Macedonia, 19 of which are local endemic taxa.

The Annexes of the Bern Convention include 12 species of higher plants.

The European List of vertebrate animals includes 113 species, of which: 30 fish species, 66 bird species, 16 mammals and 1 reptile species. Out of the total of 20 endemic fish species from the Republic of Macedonia, 17 have been enrolled in the category of globally threatened species. The total number of identified “Emerald” species (under Resolution No.6 to the Bern Convention) on the territory of the Republic of Macedonia is 165 species.

Figure 1. Number of protected species of animals




Figure 2. Number of endemic species of animals




Figure 3. Number of threatened species of animals




Figure 4. Number of protected species of plants




Figure 5. Number of endemic species of plants




Figure 6. Number of threatened species of plants





Figure 7. Number of Emerald species in Europe and in the Republic of Macedonia



coverage: **excel**

**Sources of data:** Study of Biological Diversity in the Republic of Macedonia, Strategy and Action Plan for Biological Diversity Protection in the Republic of Macedonia, Report on the establishment of the National Emerald Network, Analysis and valorization of species diversity in the Republic of Macedonia.

## Assessment

The **flora** of the Republic of Macedonia is very rich and diverse and represented by 5.843 species, of which 2.169 algae and 3.674 species of plants. Recent flora of higher plants is a mosaic of various floral elements: Tertiary relicts, Mediterranean, Greek-Asia Minor, Illyric, Caucasian, Middle European, Skardo-Pindian, Eurasian, Arctic-Alpine, cosmopolitan, among which 228 species are endemic (Balkan, South Balkan, Macedonian, local, etc.). The highest number of endemic plant species (114) has been recorded with *Angiosperms*.

The National Red List of threatened wild plant species is under establishment. The number of threatened wild plant species is shown in Table 1. The assessment of threat of species by taxonomic groups has been carried out in accordance with international criteria contained in several international documents (conventions, agreements, Global Red List, European Red List, EU Directives).

The IUCN Global Red List contains 72 taxa from the Republic of Macedonia, 19 of which are local endemic species. The Annexes of the Bern Convention list 12 plant species having their area of spreading on the territory of the Republic of Macedonia.

**Lichens** are represented by 354 species. Considering lichens studied in the Republic of Macedonia, no lichen species have been included in Annexes I and II of EU Habitat Directive. On national level, 12 lichen species have been identified to have status of being threatened.

**Fungi** compose exceptionally heterogeneous group of organisms and explorations so far have focused on the phyla of *Ascomycota* and *Basidiomycota*, while others are poorly studied. The total number of studied and recorded native fungi on the territory of the Republic of Macedonia is 1245 species. Most of those belong to the phyla *Myxomicota* (10), *Oomycota* (20), *Zygomycota* (35), *Ascomycota* (130) and *Basidiomycota* (1050).

From among studied native fungi in the Republic of Macedonia, no species has been included in Annexes II and IV of the Habitat Directive. The Preliminary National Red List of threatened fungi species includes 67 species belonging to the phylum of *Basidiomycota*.

**Fauna diversity** is characterized with high extent of taxonomic diversity, represented by 10.354 species and 228 subspecies or 10.582 taxa in total.

**Invertebrate animals** (Invertebrata) are represented by 9.819 species, 635 of which are endemic. Nevertheless, detailed analyses of threat to species by taxonomic groups have not been completed yet, and therefore the number of preliminary listed species is only 25 invertebrate animal species.

**Vertebrate wild animals** (Vertebrata) are represented by 535 species, 30 of which are endemic species. The class of fish includes 78 species, 27 (34.5%) of which are endemic species. No endemic species have been recorded among the classes of amphibians, reptiles and birds, while three endemic species have been recorded with the class of mammals. With regard to the extent of threat to populations of vertebrate animals, the class of fish has 18 species included in the category of globally threatened species.

It is of particular importance to point out that the fauna of vertebrate animals includes 113 species that have been enrolled on the European Red List, namely: 30 species of fish, 66 birds, 16 mammals and 1 reptile species. The National Red List of threatened fauna species is under development.

Within the species diversity, particular significance is attributed to the identified "Emerald" species. Namely, total of 165 species have been identified, as follows: 6 species of invertebrate animals, 154 vertebrate animals (12 species of fish, 3 species of amphibians, 7 species of reptiles, 115 species of birds, 17 species of mammals and 5 species of plants).

Two lists of strictly protected and protected species were adopted on national level (2011).

List 1: Strictly protected wild species in the Republic of Macedonia includes: 9 species of native fungi, 51 plant species, 36 invertebrate species and 98 vertebrate species.

List 2: Protected wild species in the Republic of Macedonia includes: 63 species of native fungi, 12 species of lichen, 151 plant species, 507 invertebrate species and 87 vertebrate species.

## Policy relevance of the indicator

### List of relevant policy documents:

The Second National Environmental Action Plan, in its Chapter on Nature, emphasizes the goal of the establishment of integrated system for nature and biological diversity protection, in line with multilateral agreements and EU standards, through the measure for application of mechanisms for further implementation of the National Strategy for Biological Diversity Protection with Action Plan and the National Capacity Self-Assessment (NCSA), the Law on Nature Protection and creation of appropriate conditions for Natura 2000 network establishment. It envisages action towards development of National Red Lists and Red Book of the Republic of Macedonia.

The National Strategy for Biological Diversity Protection with Action Plan (NSBAP) defines integrated approach to the protection and sustainable use of components of biological diversity. The Strategic Action Plan (SAP) outlines the specific actions to be taken to achieve the targets. One of the measures in this document is the Protection of Species, through several actions concerning



elaboration of National Red Lists and Red Book of the Republic of Macedonia, and protection of globally endangered wild species. Particularly important conservation activities concern protection of butterflies, Pelister, Pelagonia and Lokven dragonfly, Macedonian crayfish, From among vertebrate animals, critically endangered and endangered endemic and other fish species (lamprey, trout, eel), amphibians (sand-hoppers, frogs), reptiles (turtles, lizards, snakes), birds of prey (vultures, eagles, falcons, wild species of migratory waterbirds, other wild bird species, small size mammals (rodents, bats), wild carnivores: bear, lynx, wolf, golden jackal, otter, marten, badger, souslik, Balkan snow vole, Balkan chamois, red deer, deer, etc.

From among plants, the following have been identified: endemic and relict species Ema's *Thymus oehmianus*, Alshar's *Thymus alcharensis*, Macedonian ramonda (*Ramonda macedonica*), Balkan ramonda (*Ramonda serbica*), Mariana's tulip (*Tulipa marianae*), Sharplanina tulip (*Tulipa scardica*), Mayer's nepeta (*Nepeta ernesti-mayeri*), Alshar's violet (*Viola allchariensis*), Arsen's violet (*Viola arsenica*), Koshaninova violet (*Viola kosaninii*), spring Adonis flower (*Adonis vernalis*), orchids (Orchidaceae), Macedonian dwarf pine (*Pinus mugo macedonicus*), Macedonian oak (*Quercus macedonicus*), etc.

## Legal grounds

The Law on Nature Protection provides for elaboration of National Red Lists and Red Book of the Republic of Macedonia, as well as proclamation of strictly protected wild species and protected wild species, by which they shall acquire the status of natural heritage. It is also legal obligation to establish Cadastre of protected areas and Registry of natural heritage including also strictly protected and protected wild species in the Republic of Macedonia.

## Targets

Identification of the extent of threat for certain species of plants, lichens, fungi and animals found in the Republic of Macedonia, which are of global, European and national significance and definition of measures for their protection and conservation.

## Reporting obligation


- UNEP/CBD Secretariat
- Secretariat of UNEP/CMS, AEWA, EUROBATS
- Secretariat of BC/CE

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 007</b>	<b>Threatened and protected species</b>	CSI 007	Threatened and protected species	<b>S/I</b>		▪ Biological diversity	5 - annually

## MK – NI 008

### DESIGNATED AREAS



### Definition

The indicator shows the proportion of a country designated total area that is protected under national instruments, or under the EU Birds and/or Habitats Directives (Natura 2000 sites), or under the Bern Convention (Emerald sites) and other multilateral agreements.

- Total (cumulative) designated area of sites protected under national instruments, or under the EU Birds and/or Habitats Directives and under multilateral agreements over the time.

The indicator is also broken down to show the different trends of surface area in km<sup>2</sup> designated under international conventions and initiatives, under EU Directives and under national legislation:

- Number of protected areas under the national categorization
- Percentile representation of individual national categories of protected areas out of the total protected area
- Changes over time in cumulative surface area of Emerald sites (designated under the Bern Convention).
- International instruments

### Units

- Number of sites, ha, km<sup>2</sup> and %.

### Key policy issue

***What is the progress in designation of areas (km<sup>2</sup>, %) under the national legislation, EU Directives and multilateral agreements?***

### Key message

As of 1948, when the First National Park “Pelister” was designated in the Republic of Macedonia, the number and the total surface area of different categories of protected areas have noted permanent growth on national level.

At this moment, due to the new categorization of designated areas established under the new Law on Nature Protection, in accordance with the IUCN categorization, the designated area system is in transition and includes designated areas under the old and under the new categorization. The analysis (in relation to the number and the area) includes all designated areas in the Republic of Macedonia designated under the old and also under the new categorization. In doing so, the areas designated in accordance with the old categorization have been processed according to the appropriate/corresponding IUCN category. The analysis of the area of the designated areas has been made by rendering the borders of the areas in GIS.

Presently, the network of protected areas includes 86 areas in total and these occupy area of 226.809.54 ha or 8.82 % of the territory of the Republic of Macedonia. The largest portion is occupied by national parks and nature monuments.

Total of 35 sites have been identified in the National Emerald network of areas of special conservation interest. They occupy an area of 752.223 ha or 29 % of its territory.

Figure 1. Total area of designated areas




Figure 2. Area of designated areas by category of conservation




Figure 3. Share of individual national categories of designated areas in the overall territory of the Republic of Macedonia




Figure 4. Total number of designated areas





Figure 5. Area of identified Emerald areas



Data coverage: [excel](#)

Source: MoEPP - CDDA, Emerald database

## Assessment

### 1. Designated areas at national level

Under the 2004 Law on Nature Protection, new categorization of designated area is introduced, aligned with the International Union for Conservation of Nature (IUCN), enabling inclusion of the national designated areas in the world network of designated areas. The Law stipulates a responsibility that, within 6 years, all designated areas (nominated before 2004) to be re-evaluated and designated accordance with the new categorization. Because of the current transitional period, the network of designated areas (areas designated according to the new categorization redesignated areas) the analysis (regarding the number and area they occupy) includes all designated areas in the Republic of Macedonia, designated under the old and also the new categorization. In doing so, the areas designated in accordance with the old categorization have been processed according to the appropriate/corresponding IUCN category. The analysis of the area of the designated areas has been made by rendering the borders of the areas in GIS (according to the data from the acts of designation or redesignation of areas, the Spatial Plan of the Republic of Macedonia, and where precise data in the Spatial Plan were missing, the area of the designated areas was rendered in accordance with the experts opinion).<sup>1</sup>


In the analyzed period, the area of designated areas has grown, i.e. the share of designated areas in the overall area of Macedonia in 1990 was 7.14% and in 2013 it grew to 8.82%. Also, the number of designated areas recorded increase from 67 in 1990 to 86 areas in 2013, most of which – 67 areas – belong to natural monuments, followed by nature park with 12 areas.

Thus, currently the designated area network comprises 86 areas, with total area of 226.809,54 ha or 8.82 % of the territory of Macedonia.

Most of it falls into the category national parks with around 4.5 %, then natural monuments with 2.78 % and the multipurpose area Jasen with 1,05 % of the national territory.

<sup>1</sup> The analysis of the number and area of designated areas, i.e. the rendering of the borders in GIS has been done during 2010-2011, within the UNDP and GEF project „Strengthening the environmental, institutional and financial sustainability of the system of designated areas in the Republic of Macedonia “.

Map 1. National designated areas



## 2. Designated areas with internationally recognized status


- Monument of Nature “Ohrid Lake” – World Natural Heritage (UNESCO);
- Monument of Nature “Prespa Lake” – Ramsar Site (1995);
- Monument of Nature “Dojran Lake” – Ramsar Site (2008);
- Monument of Nature “Markovi Kuli” (*King Marko’s Towers*) - World Natural Heritage (UNESCO’s Tentative List); and
- Monument of Nature “Slatinski izvor” (*The Springs of Slatino*) - World Natural Heritage (UNESCO’s Tentative List).
- Biosphere Ohrid-Prespa Reserve – World Natural Heritage (UNESCO) (2014).

## 3. Emerald network

4 phase process was implemented in the Republic of Macedonia from 2002 to 2008 to identify the areas of special conservation interest (ASCI) and establish the Emerald Network. This network is established on the territory of the countries Parties to the Bern Convention and important part in the preparation of candidate countries for EU membership for the implementation of the EU Birds and/or Habitats Directives, or an additional tool in the process of establishing the coherent European Environmental Network Natura 2000.

35 locations have been identified in the National Emerald network. Under the first project, implemented in 2002-2003, 3 areas have been identified: SNR Ezerani, NP Galichica and SR Dojran Lake, with a total area of 27.660 ha (3,6%). In 2004 a second project was implemented, identifying another 3 areas: SNR Tikves, NP Pelister, SR Demir Kapija, with a total area of 28.000 ha (3,8%). Under the third project implemented in 2005-2006, ten areas with a total area of 144.783 ha (19,1%) were identified, while with the fourth project (implemented in 2008), another 19 areas were identified with a total area of 556.447 ha (73,5%). The National Emerald Network of the Republic of Macedonia identifies 35 areas covering an area of 752.223 ha or around 29 % of its territory.

Map 2. National Emerald network of the Republic of Macedonia



## Methodology

- Methodology for the indicator calculation

The procedure for identification and designation of different categories of protected areas derives from the Law on Nature Protection, EU Directives and provisions of multilateral agreements.

## Policy relevance of the indicator

### List of relevant policy documents

Spatial Plan of the Republic of Macedonia

The Second National Environmental Action Plan, in its Chapter on Nature, emphasizes the goal of the establishment of integrated system for nature and biological diversity protection, in line with the EU standards and multilateral agreements, through the measure for application of mechanisms for further implementation of the National Strategy for Biological Diversity Protection with Action Plan and the National Capacity Self-Assessment (NCSA), the Law on Nature Protection and creation of appropriate conditions for Natura 2000 network establishment.

Draft National Strategy for Biological Diversity with Action Plan has been prepared for the period 2014 to 2020 and its adoption by the Government of the Republic of Macedonia is in process.

### Legal grounds

The Law on Nature Protection provides for introduction of a system of designated areas aimed at protecting biological diversity in natural environments, natural processes, as well as abiotic characteristics and biological diversity of the area. The Law adopts new categorization of the designated areas which is in accordance with the International Union for Conservation of Nature (IUCN) model, establishing 6 categories of protected natural heritage. The law stipulates a responsibility that, within 6 years, all designated areas (nominated before 2004) to be re-evaluated and designated according with the new categorization. These areas are foreseen with the Spatial Plan of the Republic of Macedonia for the period until 2020.

### Targets

Expansion of the network of national designated areas up to around 12% in relation to the territory of the Republic of Macedonia is foreseen under the Spatial Plan of the Republic of Macedonia, the Spatial Plans of the Regions and the National Strategy for Biological Diversity Protection

The 35 areas identified by the National Emerald network will be used to form the Natura 2000 network; therefore it is necessary to strengthen the capacities at national and local level to start the process of identification of all areas in accordance with Natura 2000 and the EU Birds and Habitats Directive.

### Reporting obligation

- Annually, to the European Environmental Agency
- Annually, to the Secretariat of the Bern Convention with the Council of Europe

### General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 008	Designated areas	CSI 008	Designated areas	R	A	<ul style="list-style-type: none"><li>▪ Biological diversity</li><li>▪ nature</li><li>▪ policies</li></ul>	Annually



# ***MK – NI 009***

## **SPECIES DIVERSITY**

### ***Definition***

The indicator will show the trend in the number and distribution of selected species or species groups at national level, as relative assessment compared to the baseline year of the monitoring commencement.

At the moment, species groups considered are the birds.

The indicator will show the trend of common birds species (developing) and the trend of certain selected species of birds of prey.

### ***Units***

- Number of species, estimated number of individuals for certain species.

### ***Policy relevance of the indicator***

#### **List of relevant policy documents**

The Second National Environmental Action Plan, in its Chapter on Nature, emphasizes the goal of the establishment of integrated system for nature and biological diversity protection, in line with the EU standards and multilateral agreements. One of the actions envisaged for the goal achievement is development of national monitoring programme for biological diversity components and elaboration of national biodiversity indicators.

The National Strategy for Biological Diversity Protection with Action Plan defines integrated approach to the protection and sustainable use of components of biological diversity. The Action Plan outlines the specific actions to be taken to achieve the goals. The strategic commitment "Research and monitoring" includes action for national biodiversity indicators development. (D.1.11).

### **Legal grounds**

The Law on Nature Protection provides for organization of monitoring of nature state. The monitoring methodology needs to be specified in a regulation. The monitoring over the state of nature is carried out through: measurement, observation, assessment and control of the state of species, their habitats, habitat types, environmentally important areas, ecosystems, landscape types, monitoring and assessment of geological values and monitoring of the state of natural heritage.

### ***Targets***

Identification of the trend in populations for selected bird species and establishment of the reasons leading to reduction in their number and development and implementation of measures for the negative trend halting (contribution to the achievement of the Target 2020 for biodiversity loss prevention/reduction by 2020).

### ***Key policy issue***

What is the trend in the populations of selected bird species and what are the reasons leading to their number reduction?

## Key message

According to available data, there is a trend of populations number reduction.

Figure 1. Trend in the number of Griffon Vulture in Macedonia (by colonies)





Figure 2. Trend in the number of Griffon Vulture in Macedonia (by couples)



## Assessment

Abundance and variety of species and ecosystems are the main features of biological diversity in the Republic of Macedonia. According to the available information, this wealth comprises the imposing number of 17.604 species, out of which 976 are endemic species. According to the analysis of the richness of biodiversity in the countries of the European continent, the Republic of Macedonia is positioned at the very top of the list of states known as "European Hotspot".

Ornithofauna of the Republic of Macedonia is represented by 338 taxa (309 species and 19 subspecies) of birds.

66 bird species found in the Republic of Macedonia have been included in the European Red List. From among “Emerald” species identified in the Republic of Macedonia (under the Resolution No. 6 of the Bern Convention), 115 species are birds.

In the period from January 2003 until October 2010, within the Project for vultures conservation, monitoring of the populations of two species of birds of prey, namely Griffon Vulture and Egyptian Vulture has been carried out, accompanied also by monitoring of the Imperial Eagle (there is also older data on the said species, as well as on the Golden Eagle, Mediterranean Falcon and long-legged buzzard; however, this data should be updated and analyzed again).

The implementation of the Project for common bird species monitoring was initiated in 2007 and data on the trend in certain indicative bird species has been obtained in the course of the past four years.

## ***Methodology***

### ▪ Methodology for indicator calculation

Sampling quadrants (10 – 15) with an area of 1 km<sup>2</sup> are taken by random and common bird species were counted there by the method of linear transect. Counting took place in the period of nesting (from 15 April to 15 June) with the intensity of two searches per year. Data was statistically processed resulting in the established trend in the populations of species at national level.

With reference to birds of prey, full census of couples in nest and determination of their success in nest have been envisaged.

### ▪ Source of used methodology

We applied methodology used by the organization British Trust for Ornithology in carrying out the census of birds in nests on the territory of the United Kingdom.

### **Indicator: Number of vulture couples**

The indicator shows the estimate of the size of populations (in couples) of Griffon Vulture and Egyptian Vulture in Macedonia, based on detailed monitoring of nesting sites of the two species on the whole territory of Macedonia.

## **Data specification**

Title of the indicator	Source	Reporting obligation
Species diversity	<ul style="list-style-type: none"> <li>• Grubac, B. &amp; VELEVSKI, M. (2004-2010): Survey and monitoring of the status, breeding success and threats to the Egyptian Vulture in Macedonia- Report to BVCF/FZS.</li> <li>• Grubac B., Veleviski M., Lisicanec T., Lisicanec E., Roleviski, D.&amp;Andevski, J. (2007): Decrease of population size of the Griffon vulture <i>Gypsfulvus</i> in Macedonia and assessment of conservation measures. III Congress of Ecologists of Republic of Macedonia, 06-09.10.2007, Struga. Abstract Book, 101-102. Macedonian Ecological Society.</li> <li>• Macedonian Ecological Society and Wild Flora and Fauna Fund.</li> </ul>	

## Data coverage:

Table 1. Trend in the number of Griffon Vulture in Macedonia (by colonies)

Griffon Vulture - couples						
	Demir Kapija	Mariovo	Tikvesh Lake	Osogovo	Matka	total
2003	4	4	9	6	1	24
2004	9	3	9	4	1	26
2005	7	4	4	0	1	16
2006	3	4	5	0	0	12
2007	3	5	5	0	1	14
2008	5	8	8	0	0	21
2009	7	4	6	0	0	17
2010	6	8	6	0	0	20

Table 2. Trend in the number of Egyptian Vulture in Macedonia (in couples)


Egyptian Vulture	
year	couples
2004	36
2005	34
2006	32
2007	31
2008	28
2009	27
2010	25

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 009</b>	<b>Species diversity</b>	CSI 009	Species diversity	<b>S</b>		▪ Biological diversity	

## MK – NI 052

### FORESTS AND OTHER WOODED LAND



#### Definition

The indicator shows the overall forest and other wooded land area, the share of this area in country area, as well as share of forest landscapes naturally recovered and planted forests, share of production forest area and forests intended for soil, water, ecosystem and biodiversity protection.

The term *forest* means forest ecosystem on wooded land grown over with forest tree species and shrubs, bareland adjacent to forest, as well as other barelands and meadows within forest, forest nurseries, forest roads, seeds plantations, fire prevention passes in forest, wind protection belts on areas exceeding two ares in size, as well as forest within protected areas.

*Wooded land* is land under forest or which due to its natural properties is the most suitable for forest production, as well as land with facilities intended for forest production.

*Production forests* are primarily intended for permanent production of wood assortments and other wood products and services.

*Forests within protected areas* are designated under the Law on Nature Protection.

#### Units

- km<sup>2</sup>, ha and %.

#### Key policy issue

***What is the trend of overall forest and other woodland area?***

#### Key message

The share of overall forest and other wooded land area in overall country area ranges between 35.5% and 38.8%; forest area was the largest in 2001, and the smallest in 1990. In 2013, there was decline in the overall forest area by 0.98% compared to 2001 when the forest area was the largest.

With regard to share of wooded land areas in country area, data is available only for the period 2010 to 2013 and ranges between 3.8% and 4.4%, reflecting increase by 17.56%.

The share of planted forest area has variable trend of decrease and increase in area.

Figure 1. Share of forests and other wooded land in country area




Figure 2. Share of planted forest in country area





Figure 3. Share of production, multiple use forest and forest area within protected areas in country area



Data coverage: [excel](#)

Source of data: State Statistical Office

## Assessment

Besides production, forests also have role of nature protection through protection of biodiversity and protection against erosion. They are of great importance for sustainable development owing to influence they make on climate, water regime, oxygen production and quality of environment.

During the reporting period, the share of overall forest and other wooded land area in country area ranges between 35.5% and 38.8%; forest area was the largest in 2001, and the smallest in 1990. In 2013, there was decline in the overall forest area by 0.98% compared to 2001 when the forest area was the largest. Slight decline by 0.13% was recorded in forest area in 2013 compared to 2012 due to forest fires.

With regard to share of wooded land areas in country area, data is available only for the period 2010 to 2013 and ranges between 3.8% and 4.4%, reflecting increase by 17.56%.

The share of planted forest area has variable trend of decrease and increase in area. Significant decline by 52.8% was recorded in the period 1990 to 2001, followed by increase of 53.2% by 2003, and then decrease again of 37.3% by 2007. The greatest increase was recorded from 2007 to 2009 reaching 120.23%, to be followed by the greatest decline of 62.2% by 2013.

The share of production forests in country area, in the period 2010 to 2013 recorded increase by 4.1%; the share of multiuse forests noted decline by 11.45%, and the share of forest areas within protected areas noted drop by 11.53% in country area.

## Methodology

- Methodology for the indicator calculation

Parameters are obtained from reporting units delivered with regular statistical reports, book-keeping records, based on specific forest management plans.

## Policy relevance of the indicator

### Legal grounds

- Law on Nature Protection (Official Gazette of RM no. 67/04, 14/06, 84/07, 35/10, 47/11, 148/11, 52/12, 13/13, 163/13 and 41/14)
- Law on Forests (Official Gazette of RM no.64/09, 24/11, 53/11, 25/13, 79/13, 147/13, 43/14 and 160/14)
- Law on State Statistics (Official Gazette of RM no.54/97, 21/07 and 51/11) and Programme for statistical survey for the period 2013-2017 (Official Gazette of RM no.20/13)

### Targets

The Spatial Plan of the Republic of Macedonia specifies enlargement of forest and wooded land areas, namely by 2020 the share in country area should be 48.8%.

### Reporting obligation

- UNECE/FAO
- FOREST EUROPE Report on “State of Forests and Sustainable Forest Management in Europe”
- FAO Global Forest Resources Assessments (FRA)

### General metadata

Code	Title of the indicator	Compliance with CSI/ EEA or other indicators	Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 052</b>	<b>Forests and wooded land</b>		<b>S</b>	A	▪ Biodiversity	Annually



# CLIMATE CHANGE



# MK - NI 010

## GREENHOUSE GAS EMISSIONS



### Definition

The indicator shows the quantities of greenhouse gas emissions into atmosphere on national level. The emissions are presented by greenhouse gas type. The indicator provides information on emissions in the following sectors: energy, industrial processes and products use, land use and land use change and forestry (LULUCF) and waste.

### Units

- Kilotons CO<sub>2</sub>-equivalent

### Key message

Total emissions throughout the inventory taking period have undergone slight growth from 0.4% compared to 1990. Total national emissions in 2012 amounted 12223.37 Gg of CO<sub>2</sub>-eq. The five biggest categories with highest sources of emissions in Macedonia are as follows:

- Emissions of CO<sub>2</sub> from energy industries (coal, lignite) (49.5%);
- Emissions of CH<sub>4</sub> from solid waste landfills (11.7%);
- Emissions of CO<sub>2</sub> from mobile sources, including road motor vehicles (11.6%);
- Production and construction industries (8.8%); and
- Emissions of CH<sub>4</sub> from enteric fermentation of livestock (3.9%).

Figure 1. Total GHG emission in kilotons CO<sub>2</sub>-equivalent (baseline 2000)




Figure 2. Share by sector of GHG emissions in kilotons per year




Figure 3. Share by sector of GHG emissions in % per year





Figure 4. Share of individual pollutants in the total GHG emissions in % per year



Data coverage: [excel](#)

Source: Macedonia’s Third National Communication to UN Framework Convention on Climate Change (UNFCCC), Ministry of Environment and Physical Planning, [www.unfccc.org.mk](http://www.unfccc.org.mk)

## Assessment

In general, energy sector has the greatest contribution to national GHG emissions, with an average share of 77% in the period 1990 – 2012. Waste sector was the second sector contributor with an average share of 9%, followed by agriculture, forestry and land use with an average share of 8%. Industrial processes were the last sector contributor with an average share of 6% in the national GHG emissions in the period 1990-2012.

With regard to the assessment of key sources by sub-categories, sub-sector energy industries was the predominant source of emissions throughout the period, with an average share of 50.2% in 1990 and 49.5% in 2012.

Sub-sector production and construction industries was the second national contributor of emissions in 1990 with an average share of 13.6%. In the last year of the inventory, 2012, this sector was responsible for 8.82% GHG emissions and was therefore regarded the fourth contributor. Thus, emissions from the subsector of production noted falling trend owing to reduced industrial activity in the country, partially explained by closure of industrial plants for aluminum, lead and zinc in 2003.

Contrary to the above, emissions from the sub-sector road transport experienced significant increase of their contribution to overall national emissions in the period 1990 - 2012; the average share of emissions from this sub-sector in 1990 was 6.2% of the overall national emissions or 760.85 Gg of CO<sub>2</sub>-eq., while in 2012 this sub-sector was responsible for 11.6% of the overall national emissions or 1415.14 Gg of CO<sub>2</sub>-eq.

Similarly to this, emissions from the sub-sector of waste disposal increased significantly in the period 1990 - 2012 due to population growth which caused greater consumption and waste generation.

Emissions in the category of land use and land use change and forestry remained relatively unchanged, except in 2007, 2008 and 2012 when carbon sinks were significantly reduced because of great forest fires. The greatest portion of CH<sub>4</sub> emissions in agricultural sector (89%) was caused by enteric livestock fermentation and these emissions decreased constantly in parallel with the reduction of livestock scale. Emissions from manure amounted 8% of GHG emissions, while other emissions came from rice fields and burning of plant residues.

## Methodology

- Methodology for the indicator calculation

To calculate GHG emissions as well as GHG inventories, the methodology provided by UNFCCC/IPCC is used.

Methodology is based on the calculation of GHGs as a product from the rate of activity for individual sectors and emission factors.

In the frames of the Third Communication on Climate Change, GHGs inventory has been prepared. Identification of the key categories of sources relied on methods Tier 1 and Tier 2. Tier 1 method is used to identify categorization of key sector and establishment of the trend in emissions from the national emission inventories. Given the fact that the inventory has data for several years, estimates of shares of each category in emission level and trends were made.

Tier 1 level is also applied for certain categories of sources, by using analysis of uncertainty. Application of Tier 1 method is useful, as it enables additional identification of the reasons for which certain categories of emission are key and enables prioritization of activities in order to improve the quality of inventory and reduce overall uncertainty.

## Policy relevance

The Greenhouse Gases Inventory establishes the basis for the analysis of the GHG reduction.

## Legal grounds

Republic of Macedonia is a Party to the United Nations Framework Convention on Climate Change and to the Kyoto Protocol. Climate change issues have been incorporated in the Law on Environment, including the requirements for preparation of GHG emission inventories and GHG removal via sinks, as well as development of action plan with measures and activities aimed at GHG emissions abatement and climate change impacts mitigation. In addition to this, by means of amendment of the Law on Environment, provision has been made for Designated National Authority to approve the projects under the Kyoto Protocol Clean Development Mechanism.

## Reporting obligation

UNFCCC

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 010</b>	<b>Greenhouse gases emissions and removals</b>	CSI 010	<b>Greenhouse gases emissions and removals</b>	<b>P</b>	B	<ul style="list-style-type: none"><li>▪ air</li><li>▪ air quality</li><li>▪ climate change</li></ul>	Annually

## MK - NI 011

# PROJECTIONS OF GREENHOUSE GAS EMISSIONS



## Definition

The indicator illustrates projected trends in anthropogenic greenhouse gas (GHG) emissions by means of application of the existing policies and measures and/or additional policies and measures. Projected trends are presented by sector types: energy, industrial processes, agriculture, land use change and forestry and waste.

## Units

- Tons CO<sub>2</sub>-equivalent.

## Key message

According to specific emissions (kt CO<sub>2</sub>-eq per capita), Macedonia remains among countries with relatively high emissions per capita, mainly due to the use of fossil fuels in electricity production. Compared to the baseline scenario, this parameter notes gradual decline along with the introduction of gas in mitigation scenarios. Considering the close interaction between GHG emissions and the manner of energy production and consumption, the national policy for energy efficiency (EE) and renewable energy sources (RES) will by itself contribute to the climate change mitigation, as achievement of the objectives set in these policies will at the same time reduce the GHG emissions.

Figure 1: Projections of total GHG emissions [kt CO<sub>2</sub>-eq]- Baseline scenario




Figure 2: Projections of total GHG emissions [kt CO2-eq]- First mitigation scenario




Figure 3: Projections of total GHG emissions [kt CO2-eq]- Second mitigation scenario




Figure 4: Effectiveness of the three scenarios expressed as absolute emissions growth in 2025 relative to emissions in 2008




Figure 5: Effectiveness of the three scenarios expressed as relative emissions growth in 2025 relative to emissions in 2008







Figure 6: Projections of total GHG emissions for the three scenarios [kt CO<sub>2</sub>-eq]



Data coverage: [excel](#)

Source: First National Communication on Climate Change to UNFCCC, MEPP, UNDP, p. 47-84, 2003, Second National Communication on Climate Change to UNFCCC, MEPP, [www.unfccc.org.mk](http://www.unfccc.org.mk)

## Assessment

This Chapter integrates sectoral emissions in order to project total GHG emissions for the period 2008 - 2025 based on adopted scenarios: baseline, first and second mitigation scenarios. It should be noted that the first and the second mitigations scenarios differ only in electric power sector, where the second mitigation scenario includes additional measures for emission abatement, which will, as shown below, contribute significantly to the overall emission abatement. Total emissions at the beginning and at the end of the period by all scenarios are summarized on Table 1. Further on, Table 2 (and Figure 1), Table 3 (and Figure 2) and Table 4 (and Figure 3) present sectoral and total GHG emissions by years, for each scenario, respectively.

**Analysis of the baseline scenario:** According to projections presented in Table 2 and on Figure 1, there will be significant increase of GHG emissions by 2025 compared to values estimated for 2008 (around 9.900 kt CO<sub>2</sub>-eq in absolute value, or around 71% in relative terms), provided that business-as-usual is preserved (Figure 4 and Figure 5, the last group of columns). This increase is mainly connected to the growth in electric power sector (absolute difference of 6.400 kt CO<sub>2</sub>-eq and 78% relative growth of the value in 2008), which reflects the so called dark scenario, i.e. development scenario of the national energy sector based on lignite (Figure 4 and Figure 5, first group of columns). Other sectors also show significant growth in GHG emissions, so that values in 2025 compared to values in 2008 are higher by 75% - transport, 71% - heating and industry, 60% - agriculture and 6% - waste (Figure 4 and Figure 5).

**Analysis of scenarios of emission abatement:** The state could improve if development paths include activities/measures leading to GHG emissions reduction. As a result, the first mitigation scenario (as defined in the analysis by sectors) will lead to increase in total emissions of 46% of the values in 2025 compared to the value in 2008 or absolute difference of around 6.400 kt CO<sub>2</sub>-eq. (Table 3 and Figure 2; also Figure 4 and Figure 5, last group of columns). This increase in total emissions decreases further by 32% (absolute difference of around 4.000 kt CO<sub>2</sub>-eq) if development paths observe the second mitigation scenario (Table 4 and Figure 3; also Figure 4 and Figure 5, last group of columns).

With regard to emission projections by sectors for the three scenarios, comparison of emissions in 2025 with those in 2008 shows highest rise of emissions in electric energy sector. Namely, the relative increase of 78% in the baseline scenario falls at 41% in the first mitigation scenario due to the introduction of two plants on natural gas for combined electricity and heat production (the first one in 2009 and the second in 2015). The relative increase drops at 14% under the second mitigation scenario, as a result of reduced consumption by major consumers, introduction of renewable energy sources and termination of the thermal power plant (TPP) in Negotino upon the establishment of the new gas power plant (Figure 4 and Figure 5, last group of columns). As far as the sectors are concerned, there is a notable result in the waste sector, where the relative increase of 6% in the baseline scenario reaches negative relative increase (-13%) under both mitigation scenarios. This means that, under the mitigation scenario, the values of emissions in 2025 will be by 13 % lower compared to their values in 2008 (Figure 4 and Figure 5, fifth group of columns), owing to the introduction of technology for landfill gas combustion at several landfills in the country. Other sectors note minor contribution to the reduction of overall emissions, considering the fact that the relative difference between baseline and mitigation scenarios ranges within 2 - 4%. (Figure 5).

Summary of the projections of total GHG emissions by years, in line with the adopted scenarios, is presented in Table 5 and on Figure 6.

According to its country specific emissions (kt CO<sub>2</sub>-eq/capita), Macedonia remains among countries with relatively high emissions per capita, owing mainly to the use of fossil fuels in electricity production. Compared to the baseline scenario, this parameter notes gradual decrease with the introduction of gas under mitigation scenarios. The calculated specific emissions for the three scenarios are presented in Table 6.

## Methodology

- Methodology for the indicator calculation

The electric energy production sector (which contributes more than 50% to the total GHG emissions) has been modeled by application of the WASP software (a tool used in the energy sector development planning), while emissions estimates for other sectors have been made by using the software tool GACMO and expert judgments. Projections of GHG emissions have been made by downscaling analysis made under the national reports.

## Policy relevance

This indicator is of vital importance for the national climate change mitigation policy. It is also related to future implementation of projects based on the Clean Development Mechanism (CDM) of the Kyoto Protocol.

## Legal grounds

Republic of Macedonia is a Party to the United Nations Framework Convention on Climate Change and to the Kyoto Protocol. Climate change issues have been incorporated in the Law on Environment, including the requirements for preparation of GHG emission inventories and GHG removal via sinks, as well as development of action plan with measures and activities aimed at GHG emissions abatement and climate change impacts mitigation. In addition to this, by means of amendment of the Law on Environment, provision has been made for Designated National Authority (DNA) to approve the projects under the Clean Development Mechanism (CDM) of the Kyoto Protocol.

## Reporting obligation


- UNFCCC

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 011</b>	<b>Projections of GHG emissions</b>	CSI 011	Projections of GHG emissions and removals	<b>P</b>	A	<ul style="list-style-type: none"> <li>▪ air</li> <li>▪ air quality</li> <li>▪ climate change</li> </ul>	Annually

## MK - NI 012

### AIR TEMPERATURE



### Definition

The indicator shows the annual average temperature of the air, its development over a given period of time, and deviations from a long-term average in the country as a whole and in particular regions.

### Units

- Degrees Celsius (°C)

### Key question

What is the trend in annual temperature and average annual deviation from the long-term average temperature?

### Key message

During the reporting period, increase in annual average temperature of the air was recorded in all four populated places, and deviations from a long-term average temperature in the period 1961 - 1990 ranged from -0.3 to 2.2 °C.

Figure 1. Trend in annual temperature at the selected monitoring stations





Figure 2. Average annual deviation from the long-term average temperature at the selected monitoring

stations



Data coverage: [excel](#)

Source: Hydrometeorological Administration

## Assessment

Data is provided for the capital city (Skopje), the second major city (Bitola), and the areas or regions with the highest (Gevgelija) and lowest (Lazaropole) long-term average temperature 1961-1990. During the reporting period, increase in annual average temperature of the air was recorded in all four populated places considered, as follows: by 4.5% in Skopje, 13.39% in Bitola, 10.88% in Gevgelija and 5.1% in Lazaropole. Annual average deviations from long-term average temperature in the period 1961 - 1990 ranged between -0.3 and 2.2 °C.

## Methodology

### Methodology for the indicator calculation

Air temperature means temperature of ambient air measured in the shade (inside meteorological house) at a height of 2 meters. It is measured by standard glass thermometer (mercury or alcohol) and electronic sensor. The main parameters of air temperature for a given location are: daily average temperature, maximum daily and minimum daily temperature.

Daily average temperature is obtained by air temperatures measured at 07, 14 and 21 hours (local time) using the formula:  $T_{avg} = (T_7 + T_{14} + 2 * T_{21}) / 4$

Maximum daily temperature is the highest temperature of the air between 21 hours (local time) preceding and 21 o'clock (local time) current day.

Minimum daily temperature is the lowest temperature of the air between 21 hours (local time) preceding and 21 o'clock (local time) current day.

Processing of collected data includes calculation of average temperatures for a given period and/or finding the maximum and the minimum values.

## Policy relevance

## Legal grounds

Law on Hydrometeorological Activity (Official Gazette of RM no.103/08 and 53/11).

## Target

The European Union proposed in its climate policy that the global average temperature increase should be limited to not more than 2°C above “pre-industrial” levels

## Reporting obligation


World Meteorological Organization

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 012</b>	<b>Air temperature</b>	CSI 012 CLIM 001	<b>Global and European temperature</b>	<b>S</b>	<b>B</b>	<ul style="list-style-type: none"><li>▪ air</li><li>▪ air quality</li><li>▪ climate change</li></ul>	<b>Annually</b>

## MK - NI 051

### ATMOSPHERIC PRECIPITATIONS



#### Definition

Indicator shows annual precipitations or the overall quantity of wet deposit fallen on a given area for a given period of time, in liquid or solid state and deviations from long-term average precipitations on the whole national territory or in certain parts.

#### Units

- millimeters (mm), percentage %, litre per square meter

#### Key policy issue

What is the trend in precipitations?

#### Key message

During the reporting period, variable trend was recorded in all four populated places, while the annual average deviations from long-term average precipitations in the period 1961 - 1990 ranged between 61% and 163%.

Diagram 1. Trend in precipitations in selected populated places





Diagram 2. Deviations from long-term average precipitations in selected populated places



Data coverage: [excel](#)

Source of data: Hydrometeorological Administration

## Assessment

Data is provided for the capital city (Skopje), the second major city (Bitola), and the areas or regions with the highest (Lazaropole) and lowest (Shtip) long-term average precipitation 1961-1990. During the reporting period, variable trend in annual average precipitations was recorded in all four populated places considered. The highest level of annual average precipitations in Skopje, Bitola and Shtip was recorded in 2002, and in Lazaropole in 2010. Annual average deviations from long-term average precipitations in the period 1961 - 1990 ranged between 61% and 163%.

## Methodology

- Methodology for the indicator calculation

Wet precipitations are measured with: rain gauge, ombrometer (pluviometer) and precipitation sensor (with automatic meteorological stations). Instruments are positioned outdoors at 1 meter height. Rain gauge measures the overall precipitations accumulated during certain period. Time resolution can be 6 hours, 12 hours and 24 hours. In practice, daily sum of precipitations is usually operated with. Daily sum of precipitation means the quantity of wet deposit fallen during the period from 07 o'clock yesterday till 07 o'clock today.

Ombrometer is mechanical registration instrument recording the fallen quantity of wet deposit during time. Its minimum time resolution is between 1 and 10 minutes, depending on the type of the instrument.

Automatic sensor is electronic sensor. It has alternating time resolution, starting from 1 minute.

Based on daily sum of precipitation, monthly, annual and long-term average sums are calculated.

Data from ombrometer and automatic sensor enable calculation of precipitation intensity (quantity of wet deposit fallen in unit of time).



## Policy relevance of the indicator

### Legal grounds

Law on Hydrometeorological Activity (Official Gazette of RM no.103/08 and 53/11)

### Targets

There are no targets defined

### General metadata


Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 051</b>	<b>Precipitations</b>	CLIM 002	<b>Mean precipitation</b>	<b>I</b>	A	<ul style="list-style-type: none"><li>▪ water</li><li>▪ climate change</li></ul>	Annually

# SOIL



## MK – NI 014

### LAND TAKE



### Definition

Changes in and current status of agriculture, forest and other semi-natural land taken by urban and other artificial land development. It includes areas sealed by construction and urban infrastructure as well as urban green areas and sport and leisure facilities. The main drivers of land take are grouped in processes resulting in the extension of:

- housing, services and recreation,,
- industrial and commercial sites,
- transport networks & infrastructures,
- mines, quarries and waste dumpsites.

### Units

Units of measurement for changes and current status recording and mapping are hectares. For data presentation, the unit in km<sup>2</sup> can be used as well.

Results are presented as:

- current status of land cover based on the nomenclature adopted at European level, at five-year intervals;
- changes in land cover, at five-year intervals, presented in % of the total area of the country and % of the various land cover types.

Note: Particular attention is paid to areas changing as a result of urban systems extension leading to negative impact on the environment.


### Key policy question
































***How much and in what proportions is agricultural, forest and other semi-natural and natural land being taken for urban and other artificial land development?***

### Key message

Based on the CORINE Land COVER methodology, the largest portion of the land in the Republic of Macedonia is under forest and semi-natural areas, covering 1.548.855 ha or 59,8% of the total area. The category of agricultural land area covers 939.013 ha or 36.9% of the total area, the category of water bodies covers 56.444 ha or 2.2% of the total area, the category of artificial areas covers 41.480 ha or 1.6% of the total area, and the smallest area of 2.000 ha or 0.1% of the total area is wetlands.

Map 1. CORINE Land COVER 2000 (data of 1996)



	Airports		Vineyards		Pastures
	Water courses		Mixed forest		Road and rail networks and Associated land
	Water bodies		Complex cultivation patterns		Beaches, dunes, sands
	Land principally occupied by agriculture, With significant areas of natural vegetation		Moors and heathland		Continuous urban fabric
	Annual crops associated with Permanent crops		Mineral extraction sites		Natural grassland
	Bare rock		Discontinuous urban fabric		Sclerophyllous vegetation
	Dump sites		Sparsely vegetated areas		Sport and leisure facilities
	Green urban areas		Non-irrigated arable land		Transitional woodland/shrub
	Permanently irrigated land		Fruit trees and berry plantations		Coniferous forest
	Industrial or commercial units		Rice fields		Broad-leaved forest
	Inland marshes				

Map 2. CORINE LandCover overall changes 2000-2006





Figure 1. Changes in areas according to CORINE LandCover nomenclature, 2000 relative to 2006



Data coverage: [excel](#)

Source: CORINE Land Cover

## Assessment

Owing to characteristics of land cover of the territory of the Republic of Macedonia, out of 44 possible classifications under the CORINE Land Cover Nomenclature, 31 classifications have been identified up to the third level of the Nomenclature.

CORINE Land COVER changes between 2000 and 2006 cover territory of around 35.565 ha which is around 1.4% of the total territory of the country.

Concerns rising fact is that the biggest change occurs in area of class 311 (broad-leaved forest) into class 324 (transitional woodland with shrubs) covering total area of 18.171 ha, or 44.57% of the total changes. This change is most probably due to wood cut and forest fires.

Opposite process of changes from class 324 (transitional woodland with shrubs) into class 311 (broad-leaved forest) concerns total area of 4.275 ha, or 14.13% of the total changes is due to new forest growth.

## Methodology

- Methodology for the indicator calculation

The assessment of CORINE Land Cover in 2000 and 2006 was based on data from satellite images.

Owing to characteristics of the land cover of the Republic of Macedonia, out of the possible 44 classifications, 31 were identified. In addition to this and for the same reason, the minimum spatial unit treated within the project was reduced at 20 hectares instead of 25 hectares.

The substance of the process is photo-interpretation of satellite images consisting of:

- Delineation of boundaries of areas representing unique land area units at images with "false" colours;
- Application of interpretation keys, supporting documentation and satellite/aeroplane images for marking with identification number - class in nomenclature;
- Extrapolation of this marking and identification of all segments of the image exhibiting similar characteristics: colour, structure and composition.

Technical Guideline for CORINE Land Cover development was prepared by the European Environmental Agency.

## Policy relevance of the indicator

### Legal grounds

Under the Law on Environment, every citizen is entitled to have an access to environmental state information. This indicator provides not only data on the state of the environment (land cover), but it also facilitates uniform access thereto, both at national and European levels.

Based on the Law on Land Survey and Registration, by means of regular land survey information is provided on the types of land cover. Although these parameters do not correspond with the CORINE land cover nomenclature, there is a possibility for unique integration of land cover elements.

Law on Urban and Spatial Planning.

## Targets

Tracking the changes in land cover and mapping of current status. Changes are monitored over five-year intervals. Methodology and nomenclature have been additionally harmonized at European level, thus enabling integrated monitoring of changes at regional and European levels.

## Reporting obligation


- EEA

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MK NI 014</b>	<b>Land take</b>	CSI 014	Land take	<b>P</b>	A	<ul style="list-style-type: none"><li>▪ management</li><li>▪ nature</li><li>▪ other</li><li>▪ population</li><li>▪ soil</li><li>▪ tourism</li><li>▪ transport</li><li>▪ urbanization</li></ul>	10 - annually

## **MK - NI 015**

### **PROGRESS IN MANAGEMENT OF CONTAMINATED SITES**



#### **Definition**

The term 'contaminated site' refers to a well-delimited area where the presence of soil contamination has been confirmed and the severity of possible impacts to ecosystems and human health are such that remediation is needed, specifically in relation to the current or planned use of the site. The remediation or clean-up of contaminated sites can result in a full elimination or in a reduction of these impacts.

The term 'potentially contaminated site' includes any site where soil contamination is suspected but not verified and further investigations need to be carried out.

The progress in the management of contaminated sites has been designed to show reduction and elimination of negative effects to ecosystems and human health where environment degradation has been confirmed.

The management of contaminated sites starts with investigation that can further lead to rehabilitation or treatment of contaminated site, measures for its conservation and maintenance and revitalization of contaminated sites.

The indicator shows progress in five main steps:

- 1) site identification/ preliminary study;
- 2) preliminary investigation;
- 3) main site investigation;
- 4) implementation of remediation measures;
- 5) measure completed.

#### **Units**

- Number of sites managed up to a certain step out of the five main steps of the indicator.
- Share of economic activities in soil contamination as percentage of sites where the activity is present compared to the total number of sites.

#### **Key policy issue**

***What progress has been made in contaminated sites management and what is the share of economic activities contributing to soil contamination?***

#### **Key message**

The management of contaminated sites in the period from 2005 to 2011 showed progress with regard to the main site investigation, as well as implementation of remediation measures. With regard to completion of remediation measures, no progress has been recorded, i.e. completion of remediation measures has not been recorded in none of the identified hot-spots.

With regard to economic activities contributing to soil contamination expressed in percentage, the highest share belongs to mining and metallurgy with 31.25%, and oil refining and leather manufacturing industry with 6.25%.



Figure 1. Progress in contaminated sites management





Figure 2. Share of economic activities in soil contamination



Data coverage: excel

Data source: Ministry of Environment and Physical Planning

### Assessment

In the Republic of Macedonia, identification and preliminary investigations have been carried out with 16 sites, where soil contamination has been detected and those sites have been marked as hot-spots. Main investigations have been carried out with eight sites. Remediation measures have been implemented with three sites, while completion of measures has not been recorded with none of the sites.

With regard to economic activities contributing to soil contamination expressed in percentage, the highest share belongs to mining and metallurgy with 31.25%, followed by organic chemical industry with 12.5% and oil refining and leather manufacturing industry with 6.25%.

## Reporting obligation

Soil contamination (TE-2)

### Methodology

- Methodology for the indicator calculation

Data for the indicator calculation was taken from the National Waste Management Plan of the Republic of Macedonia or Special Study E, and from CARDS 2006 Project concerning development of remediation plans with financial requirements for elimination of industrial hot-spots, as well as Feasibility assessment and development of main technical design for water protection measures in the mine Buchim – UNDP Macedonia.

The estimated shares of economic activities contributing to soil contamination are calculated as e.g.  $[\text{number of mines contributing to soil contamination}]/[\text{total number of sites or sites where soil contamination has been confirmed}] \times 100$ .

- Source of applied methodology

According to European Environmental Agency.

### Uncertainty

- Methodological uncertainty

Although there is a definition of contaminated site, because of the lack of limit values for the concentration of certain toxic chemicals in the soil, it is difficult to determine the exact number of sites where soil contamination has been confirmed.

The assessment of contaminated site depends to a great extent on the individual expert assessment.

- Uncertainty of data set

All sites where certain industrial/economic activity is performed have not been accounted as sites with determined contamination, although such activities generate chemical substances.

## Policy relevance of the indicator

### List of relevant policy documents

The Second National Environmental Action Plan of the Republic of Macedonia.

### Legal grounds

Our country lacks legally prescribed limit values for concentrations of certain contaminating substances in soil and standards for their detection in soil. Generally, the existing legislation is intended to prevent new contaminations.

Soil protection is regulated by several laws, including those concerning environment, nature protection, agricultural land, etc., but there is no soil specific law, with clearly defined institutional responsibilities.

### Targets

Remediation of tailings, stabilization and re-cultivation of industrial landfills.

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 015</b>	<b>Progress in the management of contaminated sites</b>	CSI 015	Progress in management of contaminated sites	<b>R</b>	A	<ul style="list-style-type: none"> <li>▪ chemicals</li> <li>▪ industry</li> <li>▪ management</li> <li>▪ nature</li> <li>▪ soil</li> <li>▪ transport</li> <li>▪ urbanization</li> <li>▪ waste</li> <li>▪ water</li> </ul>	annually

## MK - NI 038 FOREST FIRES



### Definition

The indicator provides information on the number of forest fires on the territory of the Republic of Macedonia. It also provides information on the magnitude of forest fires presenting the area subjected to fire and the type of wood mass seized by fire, as well as the total damage caused by fire.

### Units

The area seized by fire is expressed in ha (hectares), while wood mass seized by fire is expressed in m<sup>3</sup>. The total damage from forest fires is expressed in denars, as well as number of forest fires.

### Key policy issue

***What is the status of forest fires in the Republic of Macedonia? What is the number of forest fires, what is the area and wood mass affected by fire?***

### Key message

In the Republic of Macedonia, rapid increase in the number of fires, area and mass affected by fire was tracked during the analyzed period, reaching the maximum of 652 fires in 2007. Then, there was a trend of gradual fall by 2009 with 61 fire events, and then the number of fires started to rise again in the following years.

The number of fires in 2013 compared to the number of fires in 2009 noted rise by 162%.

Figure 1. Number of forest fires




Figure 2. Area under fire




Figure 3. Wood mass subjected to fire





Figure 4. Total damage from fires expressed in denars



Data coverage: **excel**

Source: Public Enterprise for Forests Management - "Macedonian Forests"

## Assessment

Forest fires are among the most severe problems in forestry, as well as environment as a whole in the Republic of Macedonia. Great quantity of wood mass is destroyed by forest fires and this is an economic problem. Forest fires cause pollution of air, soil and water. Burnt woods are source of pathogens and pests. There is also increase in erosive processes in burnt areas, disturbed balance of water regime, loss of vegetation and desertification. Almost 95% of forest fires are caused by man. At an average, forest fires destroyed around 9.908.20 ha forest per year during the analyzed period. In the period from 1999 to 2013, the average number of forests per year was 214. In 2007, due to extreme draughts and human factor, the number of forest fires reached 652 fires at an area of around 35.000 ha.

In proportion with the parameters discussed above, the overall damage from fires by year expressed in denars was highest in 2007 reaching the value of 1.311.167.721,95 denars. In 2013, this value was 109.500.306.00 denars. The overall average damage from fires in the period from 1999 to 2013 amounted 285.613.377.74 denars per year.

## Methodology

- Methodology for the indicator calculation

Data and the indicator calculation were made by the Public Enterprise for Forests Management of the Republic of Macedonia - "Macedonian Forests".

## Policy relevance of the indicator

### List of relevant policy documents:

The Second National Environmental Action Plan (NEAP 2) defines measures for improved protection against forest fires, instructions on the need for capacity strengthening for sustainable forest management, as well as development of strategy for forest protection against fires.

Strategy for Sustainable Development of Forestry in the Republic of Macedonia.

### Legal grounds

- Law on Forests, which regulates forests and forest resources management and protection. Protection of forests is integrated and indivisible part of the overall forest management. In the context of forests protection against fires and regulation of measures in this area, we should also mention the 2001 Rulebook on specific measures for forest protection against fires.
- Law on Natural Rarities Protection
- Law on National Parks Protection
- Law on Fire Prevention.

## Targets

Compliance with the legislation concerning forests and forest resources protection. Reduction of forest fires number, reduction of wood mass and forest area affected by forest fires. Reduction of costs and damages resulting from forest fires. Increase of the public awareness in relation to fire prevention and undertaking all possible measures to reduce human factor as forest fires cause.

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MK NI 038</b>	<b>Forest fires</b>			<b>P</b>		<ul style="list-style-type: none"> <li>▪ Soil</li> <li>▪ Forestry</li> <li>▪ Agriculture</li> <li>▪ Nature</li> <li>▪ Urbanization</li> </ul>	annually

# MK – NI 053

## SOIL EROSION



### Definition

The indicator shows soil erosion which is natural process and is important for soil formation from geological point of view. With reference to soil erosion, particular attention is paid to accelerated erosion where natural speed of erosion is intensified due to anthropogenic factor, through application of inadequate agricultural practices, forest decrease, forest fires, construction activities, etc. Physical factors like climate, topography and soil properties are also relevant for soil erosion. Erosion of soil under the force of water is the most frequent and most severe problem both in Europe and our country. The indicator shows values for soil erosion, based on which we may identify regions affected by certain degree of erosion and develop measures for problem control.

### Units

- km<sup>2</sup> and %.

### Key policy issue

*To what extent is the soil in Macedonia affected by erosion?*

### Key message

Macedonia is one of erosion most affected territories on the Balkans. Water erosion is dominant in our country including erosion caused by rainfalls and running waters.

According to the Report of the European Environment Agency (Europe's Environment the Dobris Assessment, 1995), Macedonia belongs to the so called red zone of water erosion in Europe.

According to the Map of Erosion of Macedonia, 96.5% of the total area is under erosion process and 3.5% of the total national territory is not affected by erosion.

Diagram 1. Distribution of soil erosion







Figure 1. Map of Soil Erosion



Data coverage: [excel](#)

Source of data: Ministry of Environment and Physical Planning

## Assessment

Water erosion is dominant in our country including erosion caused by rainfalls and running waters. Aeolian erosion is represented on high mountain peaks, but damages resulting from it are negligible compared to water erosion. This is also the case of abrasive erosion which is actually caused by lake water waves in our country.

Macedonia is one of erosion most affected territories on the Balkans. There are several reasons for this, such as: long-term destructive impact by man (destruction of natural vegetation, absence of measures for agricultural soil conservation, inadequate processing resulting in deterioration of physical soil properties); relief in which mountains alternate with valleys with occurrence of steep and long inclination; erodibility of certain sediments, rocks and soils; climate conditions as torrent nature of precipitations, aridity of climate due to which natural vegetation cover on soil is weaker, and destroyed vegetation is harder to recover.

According to the Report of the European Environment Agency (Europe's Environment the Dobris Assessment, 1995), Macedonia belongs to the so called red zone of water erosion in Europe.

According to the Map of Erosion of Macedonia, 96.5% of the total area is under erosion process. 36.65% of the national area is affected by the first three categories of erosion (extreme – 2.7%, high – 7.1% and medium – 26.8%). 59.9% of the national area is affected by low and very low erosion.

Annual loss of soil is average annual loss of arable soil layer of 20 cm in thickness over an area of 8.500 ha, which is loss of 17.000.000 m<sup>3</sup> soi, every year.

The Map of Erosion of Macedonia was completed in 1992, but it was digitalized and published in 2002. It was prepared by empirical model of Gavrilovich, according to whom there are five categories and 12 sub-categories of erosion intensity.

## Methodology

### ▪ Methodology for the indicator calculation

Different methods can be used to assess erosion risk, divided mainly in expertise based assessments and model based assessments.

1. Example for expertise based assessment is Global Assessment of Soil Degradation–GLASOD). This methodology is based on obtaining answers by certain experts in all countries and its main deficiency is the issue of control of objectivity of answers given and application of different standards by different experts in different regions. This methods identifies subjectively the regions with similar intensity of soil erosion, while not taking conditions leading to that into account
2. Example for model based assessment is the Pan-European Soil Erosion Risk Assessment (PESERA). This model is appropriate for anticipation of soil erosion degree, which takes into account precipitations and eroded sediments, daily precipitations accumulated during month, monthly vegetation cover, certain climate information, etc.

Homogenous and comprehensible data on soil erosion degree, including soils in our country, is obtained through the process of calculation and modeling, which use:

- Pedological maps
- Land cover maps (Corine Land Cover 2000)
- Topographic maps
- Climate maps with temperature data

Under the recommendation of the European Commission, soil erosion may be determined through appropriate recommended methods of modeling and on the basis of methods using expert assessment.

## Policy relevance of the indicator

### Legal grounds

- Law on Environment (Official Gazette of RM no. 53/05, 81/05, 24/07, 159/08, 83/09, 48/10, 124/10, 51/11, 123/12, 93/13, 187/13 and 42/14)
- Law on Agricultural Land (Official Gazette of RM no. 135/07, 18/11, 42/11, 148/11, 95/12, 87/13 , 106/13, 164/13 and 39/14)
- Second National Environmental Action Plan 2006
- Thematic Soil Strategy (COM (2006)231)
- Draft Framework Directive on Soils (COM (2006)232)
- To Thematic Strategy for Soil Protection (CEC, 2002)

### Targets

According to the Spatial Plan of the Republic of Macedonia, counter-erosion protection of space should be conducted by combined measures, such as:

- Targeted use of areas susceptible to erosion;
- Terracing, intensive afforestation of erosion active areas and forests tending;
- Regulation of torrent watercourses with structural facilities;
- Special measures for protection against ruining, rock falls and land slides on steep slopes and river banks.

Regulation of torrent watercourses of I and II category of destructiveness which include 10.15 % of overall areas affected by erosive processes will contribute to remediation of erosive areas and regulation of torrent watercourses.



## Reporting obligation

- JRS (Joint Research Centre)/EEA (European Environment Agency)/EIONET

## General metadata

Code	Title of the indicator	Compliance with CSI/ EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 053</b>	<b>Soil erosion</b>	CLIM 028	Soil erosion	<b>P</b>	A	<ul style="list-style-type: none"><li>▪ Soil</li><li>▪ Climate change</li></ul>	Every ten years

# WASTE



# MK - NI 016

## MUNICIPAL WASTE GENERATION



### Definition

The indicator presents generation of the total amount of waste as a whole, as well as the waste generated by economic activities and municipal waste.

### Units

- It is expressed in tons per capita per year.

### Key question

Is the amount of generated municipal waste decreasing?

### Key message

In 2012, full data on the amount of the total waste generated by all activities was obtained for the first time and in future it will be possible to observe the trend of generation of waste as a whole and by economic activities.

With reference to municipal waste, we may say that generation of municipal waste went hand in hand with the economic growth by 2012, while 2012 was marked by fall in the economic growth, but increase in generated municipal waste. This situation might be a result of improved process of collection of data and information on municipal waste, namely receipt of comprehensive and precise data on the quantity of generated municipal waste.

Figure 1. Total generated waste per capita




Figure 2. Generation of municipal waste in kg per capita per year




Figure 3. Total generated waste





Figure 4. Total generated waste by economic activities



Data coverage: [excel](#)

**Source of data:** Releases of the State Statistical Office, Strategy for Waste Management in the Republic of Macedonia (2008-2020), MEPP, National Waste Management Plan (2009-2015) of the Republic of Macedonia, MEPP

## Assessment

The generated waste as a whole assumes loss of resources in a form of matter and energy. Treatment and removal of generated waste may cause pollution of the environment and expose people to harmful substances and infective agents. The generated waste is closely related to the level of economic activity in the country and indicates the consumption and production scheme in a society. The amount of generated waste as indicator may show direction of movement of the use and the consumption of products and goods and it indirectly reflects the potential environmental burden if municipal waste fails to be recognized as potential resource. Increased generation of municipal waste indicates increased use and consumption of products and goods.

Calculation of the total generated waste from 2003 to 2008 was based only on data on municipal waste, while in 2008 and 2010 data on generated waste by individual sectors of economic activities, though not all, was calculated as well. In 2012 calculations of the total generated waste took into consideration the amounts of generated waste by all sectors of economic activities. According to data from 2012, the biggest amount of waste was generated in the sector Other economic activities, followed by waste generated by the sector Manufacture and mining.

Important message contained in the relevant policy documents in the area of waste is to decouple waste generation from economic growth in the Republic of Macedonia. Estimates and data indicate that around 3% of the overall waste generated in the Republic of Macedonia is municipal waste. The observation of the period from 2003 to 2007 showed that the GDP real growth rate noted slight growth of 2.8% to 6.5%, to fall in 2009 by -0.4%. Then, in 2010 and 2011 there was growth by 3.4 and 2,3, respectively. In 2012, fall in GDP real growth rate was noted amounting -0.5%.

In the same period from 2003 to 2007, municipal waste recorded increase by more than 3% at annual level. In the period between 2008 and 2012, generated municipal waste first noted increase by 1.4%, then

decrease by 0.85% and again slight increase by 1.7%, as well as increase by 7% for 2012. . This leads to the conclusion that generation of municipal waste went hand in hand with the economic growth by 2012, while 2012 was marked by fall in the economic growth, but increase in generated municipal waste. This situation might be a result of improved process of collection of data and information on municipal waste, namely receipt of comprehensive and precise data on the quantity of generated municipal waste.

The amount of generated municipal waste expressed in persons was 348 kg/person/year or 0.9 kg/person/day for 2008, for 2009 it was 354 kg/person/year, while for 2010, 2011 it was 351 kg/person/year and 357 kg/ person/year, respectively. For 2012, the amount of generated municipal waste expressed in persons was 382 kg/person/year. These figures also indicate lower generation of municipal waste per person on annual level in the Republic of Macedonia compared to European Union Member States.

## Methodology

- Methodology for the indicator calculation

Certain types of generated waste are grouped by the main economic activities based on the National Classification of Activities (NCA Rev.2), which has been harmonized with the International Standards for Industrial Classification of All Economic Activities (ISIC). Types of wastes are determined by the 2005 List of waste types.

Data collection is performed mainly through surveys, estimates, administrative data. Reports are in a form of Relases of the State Statistical Office (2009, 2010, 2011, 2012), and annual reports by municipalities.

## Policy relevance

### **List of relevant policy documents:**

Second National Environmental Action Plan of the Republic of Macedonia (2006)

Strategy for Waste Management in the Republic of Macedonia (2008-2020)

National Waste Management Plan (2009-2015) of the Republic of Macedonia

### **Legal grounds**

Law on Waste Management (2004)

National Classification of Activities NCA Rev.2 2006)

## Reporting obligation

- EUROSTAT

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 016	Municipal waste generation	CSI 016	Municipal waste generation	P	A	<ul style="list-style-type: none"> <li>Households</li> <li>Economy</li> <li>Population</li> <li>Urbanization</li> </ul>	yearly



Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
						▪ Waste	

# MK - NI 056 MANAGEMENT OF HAZARDOUS WASTE



## Definition

This indicator shows the amounts of generated hazardous waste, amounts of imported and exported hazardous waste, amount of temporarily stored hazardous waste, as well as treated hazardous waste as a whole and individually – recycled, incinerated, landfilled, etc.

## Units

- The amounts of hazardous waste are presented in tons and share of each manner of waste treatment (%).

## Key question

*What is the manner of hazardous waste management, i.e. what is the trend of hazardous waste generation, import and export and the manner of its treatment?*

## Key message

Prevailing manner of hazardous waste management is its landfilling, followed by other manners of disposal, then recycling and hazardous waste incineration as the last option.

Data that we have on hazardous waste management is insufficient to establish and track the trend of hazardous waste management for each manner of waste treatment.

Figure 1. Overview of the total hazardous waste generated and treated or landfilled presented in tons in the period 2008-2012




Figure 2. Overview of hazardous waste treatment in 2008 in percentage




Figure 3. Overview of hazardous waste treatment in 2010 in percentage





Figure 4. Overview of hazardous waste treatment in 2012 in percentage



**Data coverage:** excel

**Source of data:** State Statistical Office

## Assessment

Surveys by the State Statistical Office have been carried out every two years starting in 2008. The survey in 2008 was conducted only in the sector of manufacturing industry; in 2010 it was extended to cover the sectors mining and electricity production, supply of hot water, steam and gas. Only in 2012, the survey was conducted in all sectors of economic activity according to the National Classification of Activities NCA Rev. 2.

Prevailing manner of hazardous waste management in 2010 and 2012 was its disposal as the least favourable manner of waste management with regard to consequences on environment and human health. The rate of hazardous waste incineration and recycling was below 1% in 2010 and 2012, while in 2008 when only the production activities sector was taken into consideration, the rate of hazardous waste recycling was 56.25 %, and incineration was again below 1%.

The waste that cannot be treated or disposed appropriately is temporarily stored. Temporarily stored hazardous waste is the waste pending disposal or treatment. Treatment can be carried out in the country of generation or in another country. Uncontrolled transboundary movement of hazardous waste and its disposal or inappropriate handling may cause heavy health problems in people and may contaminate water and soil. Recycling, appropriate incineration and appropriate disposal of hazardous waste in the country of its disposal reduces the demand for hazardous waste transboundary movement and thus reduces the risk for human health and environment. In certain cases, international transport of hazardous waste is necessary and justified in terms of the waste proper disposal and treatment with no consequences for human health and environment, and such is its reuse as secondary raw material or for energy production. Data on generated, imported, exported and disposed hazardous waste helps to control its movement and removal. Data on hazardous waste import, export and transit on the territory of the Republic of Macedonia is not available.

## Methodology

Certain types of hazardous waste are grouped according the main economic activities based on the National Classification of Activities NCA Rev. 2 harmonized with the International Standards for Industrial Classification of All economic activities (ISIC). Types of hazardous waste are determined by the List of waste types. Appropriate treatment and disposal of waste are in accordance with definitions and conditions set in the Law on Waste Management. Collection of data is acquired mainly through surveys, estimates, and administrative data. Reports are in a form of releases of the State Statistical Office (2008, 2010, and 2012).

## Policy relevance

### *List of relevant policy documents:*

Second National Environmental Action Plan of the Republic of Macedonia (2006)

Strategy for Waste Management in the Republic of Macedonia (2008-2020)

National Waste Management Plan (2009-2015) of the Republic of Macedonia

### Legal grounds

- Law on Waste Management (2004)
- National Classification of Activities NCA Rev.2 2006)
- List of waste types (2005)
- Ratified Basel Convention

## Targets

Establishment of integrated waste management and financially self-sustainable waste management system.

## Reporting obligation

- EUROSTAT

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 056	Management of hazardous waste	UNECE	I-2: Management of hazardous waste	P R	A	<ul style="list-style-type: none"><li>▪ economy</li><li>▪ population</li><li>▪ urbanization</li><li>▪ waste</li></ul>	2 - year

## MK - NI 057

### FINAL WASTE MANAGEMENT



### Definition

This indicator shows the final management of the overall amount of waste divided by sectors (industrial and municipal waste), as well as divided by the hazard it possesses (hazardous waste) through the processes of:

- Incineration (with and without energy recovery)
- Landfilling (controlled or uncontrolled landfills)
- Composting
- Reuse or recycling
- Other manner of management.

### Units

- Tons/year, percentage (%).


### Key question

*In which way or through which processes the final waste management is performed?*

### Key message

Waste landfilling on controlled landfills is the prevailing process in the final waste management in the Republic of Macedonia, followed by waste disposal on uncontrolled landfills. The processes of composting, reuse, recycling of municipal waste, as well as waste incineration with energy recovery are almost not represented in the country.

Figure 1. Overview of % of municipal waste landfilling on controlled and uncontrolled landfills



Data coverage: **excel**

Source of data: State Statistical Office

## Assessment

The way in which the waste is managed in the country has great impact on environment, economy and human health and wellbeing. Adequate waste management assumes that the Government is aware of the risks of waste for human health and environment and it supports and promotes appropriate measures to prevent waste generation or reduce it, as well as to manage it properly. Reduction in the amount of generated waste, as well as reuse and recycling of generated waste are ecologically the most favourable processes of waste management, as it also accomplishes reduction in raw materials and resources extraction. Another option of management of the waste that cannot be reused or recycled is incineration with energy recovery. The last option in waste management is its disposal in landfills and technically appropriately managed and controlled landfills are recommended for this purpose.

The two most represented processes of waste management, namely disposal of waste on controlled and uncontrolled landfills, are rather unfavourable for the environment, health of people and animals, as well as economy. Waste disposal on controlled landfills in the period 2009 to 2014 ranged from 76.07 % to 74.02 %, while disposal of waste on uncontrolled landfills was in the range between 23.93 % in 2009 and 15.53 % in 2014. All this indicates pollution of the environment and loss of natural resources. Absence of the processes of reuse, recycling, composting and incineration of waste with energy recovery reflects lack of recognition of the waste as resource and lack recovery of energy and matter contained in the waste.

## Methodology

Types of hazardous waste are determined by the List of waste types. Appropriate treatment and disposal of waste are in accordance with definitions and conditions set in the Law on Waste Management. Collection of data is acquired mainly through surveys, estimates, and administrative data. Reports are in a form of releases of the State Statistical Office (2009, 2010, 2011, 2012, 2013, 2014).

## Policy relevance

### ***List of relevant policy documents:***

Second National Environmental Action Plan of the Republic of Macedonia (2006)

Strategy for Waste Management in the Republic of Macedonia (2008-2020)

National Waste Management Plan (2009-2015) of the Republic of Macedonia

### **Legal grounds**

- Law on Waste Management (2004)
- List of waste types (2005)

## Targets

Establishment of integrated waste management and financially self-sustainable waste management system .

## Reporting obligation


- EUROSTAT

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 057</b>	<b>Final waste management</b>	UNECE	I4a - Final waste disposal: Management of municipal waste	I	A	<ul style="list-style-type: none"> <li>▪ households</li> <li>▪ economy</li> <li>▪ population</li> <li>▪ urbanization</li> <li>▪ waste</li> </ul>	2 - years




# WATER



## MK - NI 018

### USE OF FRESHWATER RESOURCES



#### Definition

The indicator observes the exploitation of freshwater resources according to their use in individual sectors, such as: public water supply, irrigation and electricity production (cooling), losses of water in water supply systems of legal persons registered for water abstraction for manufacturing or distribution of water, as well as the water exploitation index (WEI).

#### Units

- Water exploitation index - WEI expressed in %;
- Quantity of freshwater resources used is expressed in million m<sup>3</sup> per year.

#### Key policy issue

*Is water resources approximation based on water resources sustainability?*

#### Key message

In the period 1990 – 2013, oscillatory trend was tracked in water resources use. Particular rise was recorded in 2012 where the biggest quantities of freshwater resources were used for irrigation. This is due to the fact that 2012 was dry year and distribution of precipitations was such that enabled filling of water accumulations with the required quantities of water for irrigation.

Figure1. Water exploitation index




Figure 2. Water resources use by sectors




Figure 3. Abstracted water




Figure 4. Gross abstracted surface freshwater




Figure 5. Gross abstracted ground freshwater




Figure 6. Freshwater loss in transport





Figure 7. Net distributed freshwater



Data coverage: [excel](#)

Source: State Statistical Office, Water Management Administration, Public Water Supply and Sewerage Enterprise, Water communities

### Assessment

In the period 1990 – 2013, variable trend was tracked in freshwater resources use in the country. Particular rise was recorded in freshwater consumption in 2004 and 2012. Processing industry and irrigation are the main users of fresh surface and ground freshwaters during the analyzed period. In the years from 2000 to 2003, as well as in 2008, 2011 and 2012, the quantities of abstracted freshwater for irrigation exceeded the quantities for processing industry. Throughout the analyzed period, freshwater consumption for electricity production, i.e. power plants cooling, was the lowest.

As far as freshwater abstraction is considered, Figure 3 shows that surface freshwaters recorded the biggest

quantities of freshwater were abstracted in 2012. During the analyzed period, abstraction of ground freshwater resources was without major oscillations except in 2012 when rise was recorded. In the same year (2012) the values of abstracted surface freshwater were the highest, too.

Abstraction of ground freshwater resources is mainly for freshwater supply for households and part of industry, while the share of agriculture, forestry and fishery is negligible.

Diagram number 6 shows that the biggest loss of freshwater in transport occurred in 2004 and 2005, while the lowest losses were recorded in 2007 and years from 1999 to 2003. Contrary to this, in abstraction of ground freshwaters at net distributed freshwater, most of the freshwater is consumed for agriculture, forestry and fishery.

## Methodology

### ▪ Methodology for the indicator calculation

Data is collected and processed by sectors and types of industry.

Water Exploitation Index (wei) is calculated by the mean annual total abstraction of freshwater divided by the mean annual total renewable freshwater resource at the country level.

$$WEI = (\text{totABS}/\text{LTAA}) * 100$$

Where: totABS = mean annual value of total freshwater abstraction for all purposes; LTAA = long term annual mean value of freshwater resources, where data is expressed in average for a period of at least 20 consecutive years. Unit =%

## Policy relevance of the indicator

### List of relevant policy documents:

The National Environmental Action Plan - 2 and Environmental Monitoring Strategy and Data Management Strategy.

The policy for sustainable use of water resources based on the Sixth Environmental Action Programme and Framework Water Directive requirements.

National Strategy for Waters.

### Legal grounds

The Law on Waters prescribes the basic planning documents for protection, maintenance and constant improvement of the disposable water resources and rational use of the available water quantities.

Basic planning and water management development documents in the Republic of Macedonia are:

- The National Water Strategy
- Water Master Plan of the Republic of Macedonia and
- River Basin Management Plans.

The Law specifies that the maintenance and improvement of water regime is carried out on the basis of River Basin Management Plans. Such Plans contain the environmental protection goals, good status of surface water bodies (good quantitative status and chemical status, including good ecological potential) and of the groundwater resources (good quantitative status and chemical status).

Use of water for different purposes is specified under the Decree on Water Classification, according to which water is divided into five different classes based on the level of pollution, while water characteristics are determined on the basis of classes and purposes for which water can be used.

## Targets

No specific targets.

## Reporting obligation


- OECD/EUROSTAT

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MK NI 018</b>	<b>Use of freshwater resources</b>	CSI 018	Use of freshwater resources	<b>P</b>	A	▪ water	annual

## MK - NI 019

# OXYGEN CONSUMING SUBSTANCES IN RIVERS



## Definition

The key indicator for the oxygenation status of water bodies is the biochemical oxygen demand (BOD) which is the demand for oxygen resulting from organisms in water that consume oxidisable organic matter. The indicator illustrates the current situation and trends regarding BOD and concentrations of ammonium ( $\text{NH}_4$ ) in rivers.

## Units

- Annual average BOD after 5 or 7 days incubation ( $\text{BOD}_5/\text{BOD}_7$ ) is expressed in  $\text{mg O}_2/\text{l}$  and annual average total ammonium concentrations in micrograms N/l.

## Key policy issue

***Has pollution of rivers by biochemical oxygen demand ( $\text{BOD}_5$ ) and ammonium not noted increase?***

## Key message

During the assessed period from 2000 to 2013, stable trend in BOD 5 and concentrations of ammonium in rivers was tracked up to 2008, followed by decrease in BOD 5 and ammonium concentrations. As far as ammonium concentrations in rivers are concerned, there are significant variations every year. At some monitoring stations, located on the rivers Crna Reka and Vardar, moderately eutrophic water status related to BOD value was recorded. These results could reflect the status of inefficient treatment of urban and industrial wastewaters in the country, as well as the inadequate protection of river basins.

Adequate protection of rivers, and especially the introduction of regular wastewater treatment in the country, is the top priority of the policy at both national and local levels.

Figure 1. Biochemical oxygen demand ( $\text{BOD}_5$ ) in rivers






Figure 2. Biochemical oxygen demand (BOD5) in rivers by river




Figure 3. Total ammonium in rivers





Figure 4. Total ammonium in rivers by river



Data coverage: [excel](#)

Source: MEPP, HMA, HBI

## Assessment

There was a stable trend in BOD 5 and concentrations of ammonium in the rivers in the Republic of Macedonia in the period 2000 to 2008. Decrease in BOD 5 concentrations and ammonium concentrations were tracked in 2009 and 2010, followed by slight increase of concentrations in the river of Vardar. These results could reflect the status of inefficient treatment of urban and industrial wastewaters in the country, as well as the inadequate protection of river basins.

## Methodology

- Methodology for the indicator calculation

Indicators calculation is based on the methodology established by Eurowaternet, determined by the European Topic Centre for water under the European Environmental Agency.

This process defines the manner of selection of the monitoring stations, the types of parameters to be monitored, as well as the frequency of their collection.

## Policy relevance of the indicator

### List of relevant policy documents:

The National Environmental Action Plan - 2,

The Environmental Monitoring Strategy and Environmental Data Management Strategy,

Strategy for Waters has been developed in order to establish long-term policy that will secure sustainable development of waters by meeting the demands of all water users, protecting waters against pollution and pollution control.

The Law on Waters transposing the following EU Directives into the national legislation:

- Framework Water Directive (FWD) 2000/60/EEC, according to which, by the year of 2015, rivers in EU should achieve good ecological status or good ecological potential.
- Directive on nitrates (91/676/EEC), the goal of which is to reduce nitrates and pollution by

organic matter originating from agricultural lands.

- Directive on urban wastewater treatment (91/271/EEC) aimed at reducing the pollution from sewerage and industrial wastewater treatment plants

The Law on Environment has transposed the Directive on Industrial Pollution Prevention and Control (IPPC) 96/61/EEC is aimed at control and prevention of water resources pollution by industry.

## Legal grounds

The Law on Waters prescribes the main planning documents for water protection, maintenance and permanent improvement of available water resources and sustainable use of available water quantities.

The main planning documents for water management planning and development include:

- The National Strategy for Waters
- Water Management Master Plan of the Republic of Macedonia, and
- River basin management plans.

For the purpose of maintenance and improvement of the quality of water and establishment of the adequacy of water for use for different purposes, the Law on Waters specifies classification of waters and categorization of water bodies, as well as specification of deadline for achievement of the water quality goals for each water category and specification of the minimum standards for water quality and environmental protection goals for all water bodies. According to the Law, management plan will be adopted for each river basin, in order to achieve the environmental protection objectives.

The Decree on categorization of water courses, lakes, accumulations and water resources (1999) specifies the quality of water by specific classes of water in water bodies, lakes, accumulations and groundwater resources. This Decree also establishes five categories of water courses.

## Targets


Reduction and prevention of water pollution and thus achievement of good ecological status or potential of waters. Requirements of the relevant EU Directives (FWD, urban wastewater treatment, nitrates, Directive on hazardous substances, as well as Directives on drinking and bathing waters) have been transposed in the Law on Waters.

## Reporting obligation

- EEA

## General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
MK NI 019	Oxygen consuming substances in rivers	CSI 019	Oxygen consuming substances in rivers	S	A	water	annual



## Definition

Concentrations of orthophosphate and nitrate in rivers, total phosphorus and nitrate in groundwater bodies. The indicator can be used to illustrate geographical variations in current nutrient concentrations and temporal trends.

## Units

- Concentration of nitrate is expressed as mg nitrate (NO<sub>3</sub>)/l, and orthophosphate and total phosphorus as mgP/l.

## Key policy question

*Has the nutrients concentration in water courses shown rising trend?*

## Key message

Despite of the absence of continuous monitoring of the status of groundwaters quality in the Republic of Macedonia during the last years, it can be stated that the concentration of nitrates in drinking water has been in a stable environmental health status.

In the analyzed period, slight drop was recorded in the mean annual concentrations of nitrates and orthophosphates in all three rivers. The River of Vardar was an exception in 2013, when insignificant increase in orthophosphates concentrations was recorded.

Throughout the investigation period, the Lake of Ohrid has sustained its oligotrophic nature as shown on the Table on the concentrations of phosphorus and nitrates. Significantly higher concentration was found in the waters of the Lake of Prespa, thus increasing the risk of Lake's water eutrophication.

Figure 1. Nitrates and orthophosphates in rivers




Figure2. Nitrates in rivers by river




Figure3. Orthophosphates in rivers by river




Figure 4. Total phosphorous in lakes





Figure 5. Total nitrate in lakes



Data coverage: [excel](#)

Source: MEPP, HMA, HBI

## Assessment

Annual mean concentrations of nitrates and orthophosphates have remained relatively stable since the beginning of 1990's. It has been found out that the concentration of these parameters is higher at certain measuring points of VardarRiver.

Analysis of the results from the measurements in the plagian parts of Ohrid Lake throughout the period has confirmed the oligotrophic nature with relatively stable concentrations of phosphorus (below 0.015 mg./l), with higher concentrations of phosphorous (0.030 mg/l) recorded in 2007 in Ohrid Lake and nitrates concentrations within permissible limits (mean annual concentrations below 0.55 mg/l). In 2013, investigations were carried out in Ohrid Lake littoral zone and therefore higher concentrations of nitrates and total phosphorous

were recorded. Concentrations are significantly higher in Prespa Lake, where organic compounds are found at high levels, thus increasing the risk of Lake's water eutrophication.

## Methodology

### ▪ Methodology for the indicator calculation

The calculation of the indicator is based on the methodology established under Eurowaternet, established by the European Topic Centre for waters of the European Environmental Agency.

Under this process, the manner of selection of monitoring stations is defined and the type of monitoring parameters and the frequency of their collection are specified.

## Policy relevance of the indicator

### List of relevant policy documents:

The National Environmental Action Plan - 2 and the Environmental Monitoring Strategy and Environmental Data Management Strategy.

Strategy for Waters has been developed in order to establish long-term policy that will secure sustainable development of waters by meeting the demands of all water users, protecting waters against pollution and providing pollution control.

The Law on Waters transposing the following EU Directives into the national legislation:

- Framework Water Directive (FWD) 2000/60/EEC, according to which, by the year of 2015, rivers in EU should achieve good ecological status or good ecological potential.
- Directive on nitrates (91/676/EEC), the goal of which is to reduce nitrates and pollution by organic matter originating from agricultural lands.
- Directive on urban wastewater treatment (91/271/EEC) aimed at reducing the pollution from sewerage and industrial wastewater treatment plants

The Law on Environment has transposed the Directive on Industrial Pollution Prevention and Control (IPPC) 96/61/EEC is aimed at control and prevention of water resources pollution by industry.

### Legal grounds

The Law on Waters prescribes the main planning documents for water protection, maintenance and permanent improvement of available water resources and sustainable use of available water quantities.

The main planning documents for water management planning and development include:

- The National Strategy for Waters
- Water Management Master Plan of the Republic of Macedonia, and
- River basin management plans.

For the purpose of maintenance and improvement of the quality of water and establishment of the adequacy of water for use for different purposes, the Law on Waters specifies classification of waters and categorization of water bodies, as well as specification of deadline

for achievement of the water quality goals for each water category and specification of the minimum standards for water quality and environmental protection goals for all water bodies. Such plans contain environmental protection objectives in order to achieve good status of surface water bodies (good quantitative and chemical status, including also good ecological potential) and ground water resources (good quantitative status and chemical status).

The Decree on categorization of water courses, lakes, accumulations and water resources (1999) specifies the quality of water by specific classes of water in water bodies, lakes, accumulations and groundwater resources. This Decree also establishes five categories of water courses

Under the Law on Waters, authorities responsible for health protection are obliged to carry out monitoring of waters intended for human consumption and bathing waters, and for undertaking measures for active protection of the population against communicable diseases of high social and health relevance. The competent institutes perform microbiological parasitological, hygienic, toxicological and biochemical analyses within the scope of their activity.

Programme for preventive health protection performs monitoring over the quality of surface waters at all points of health interest, in order to enable timely undertaking of measures for population protection. Waters used as drinking water sources, sports and recreation and primary agricultural production are of highest interest..

## Targets

The indicator is not related directly to the requirements of a single Directive. Ecological quality of surface water requiring reduction of eutrophication and nutrient concentrations is a target specified in several Directives, namely:

- Directive on drinking water (98/83/EC) – maximum permissible concentration of nitrates is 50 mg/l;
- Directive on abstraction of surface water intended for drinking (75/440/EEC) requires nitrates concentration of 25 mg/l.
- Directive on nitrates (91/676/EEC) requires identification of groundwater bodies where the annual concentration exceeds or may exceed 50 mg/l/nitrates.
- Directive on urban wastewater treatment (91/71/EEC) specifies reduction of the pollution caused by organic matter as its objective.

## Reporting obligation

- EEA


## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MKNI 020	Nutrients in freshwaters	CSI 020	Nutrients in freshwater	S	A	Water	Annually



## MK - NI 022

### BATHING WATER QUALITY



### Definition

The indicator describes the changes over time in the quality of designated bathing waters in terms of compliance with standards for microbiological parameters (total coliforms and faecal coliforms) and physicochemical parameters (mineral oils, surface-active substances and phenols) introduced by the EU Bathing Water Directive (76/160/EEC).

### Units

- The data is expressed in a form of percentage of inland bathing waters with mandatory standards and levels specified in guidelines for microbiological and physicochemical parameters.

### Key policy issue

*Has the quality of bathing water improved?*

### Key message

The quality of lake water is at mainly satisfactory level. However, there are rivers which with their entry into the lakes contribute to deterioration of the quality of lake water. The percentage of samples with non-compliant quality is still very high (especially for physical and chemical parameters) Settlements around the three natural lakes are among the rare ones with wastewater treatment plants available in the country.

Approximation of the national legislation and standards in this area with the EU Bathing Water Directive should continue.

Figure 1. Quality of bathing freshwater - lakes





Figure 2. Quality of bathing freshwater – artificial lakes



Data coverage: **excel**

Source: Public Health Institute of the Republic of Macedonia

## Assessment

The greatest proportion of water areas in the country belongs to natural lakes, the shores of which are used for recreation purposes. The quality of water in these lakes is threatened by discharges of wastewater, uncontrolled use of lake waters for agricultural and tourism purposes, as well as by weather conditions. Apart from natural lakes, there are artificial lakes - water accumulations in the Republic of Macedonia, used for both recreation and economic purposes.

The problems of bathing water quality protection in the lakes are closely related to the implementation of one of the highest priorities in the country's environment protection - construction of adequate wastewater treatment facilities.

As international waters, the waters of the biggest natural lakes, i.e. Ohrid and Prespa, are also subject of bilateral and trilateral agreements between the Republic of Macedonia, Republic of Albania and Republic of Greece, respectively.

## Methodology

- Methodology for the indicator calculation

Standard methodology for sampling - annual data.

## Policy relevance of the indicator

### List of relevant policy documents

The National Environmental Action Plan - 2 and the Environmental Monitoring Strategy and Environmental Data Management Strategy.

Bathing Water Directive (76/160/EEC) requires the countries to identify water bodies intended for

bathing and carry out monitoring of their quality during the bathing period. Water bodies identified for bathing are those water bodies designated by the competent authorities and those where bathing has been practiced traditionally by high number of swimmers. The bathing period is determined in accordance with the period during which the highest number of swimmers is present. Qualitative monitoring takes place on daily basis during the bathing season, as well as two weeks before the commencement of the bathing season. 95% of the samples have to comply with mandatory standards.

### Legal grounds

Law on Waters, Decree on categorization of water courses, lakes, accumulations and water resources.

### Targets

It is necessary that all water bodies identified for bathing comply with mandatory values of water quality specified in Bathing Water Directive and the provisions of the Law on Waters.

### Reporting obligation

- WHO

### General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MKNI 022	Bathing water quality	CSI 022	Bathing water quality	S	B	shore water	annually

## **MK - NI 024**

# URBAN WASTE WATER TREATMENT



## Definition

Percentage of population connected to primary, secondary and tertiary wastewater treatment plants. The indicator illustrates:

1. changes in wastewater treatment;
2. conformity in terms of providing primary, secondary and tertiary treatment;
3. levels of urban wastewater treatment in large cities (agglomerations >150 000 p.e.).

## Units

- Percentages of population connected to primary, secondary and tertiary wastewater treatment.

## Key policy issue

***To what extent will the system of urban waste waters collection, removal and treatment improve the status of freshwaters in the Republic of Macedonia?***

## Key message

The requirements of the Directive concerning municipal wastewater treatment have not been implemented in the current Law on Waters.

According to the results on the distribution of the population in the Republic of Macedonia in relation to treated municipal wastewaters involving only mechanical treatment, biological treatment and latest treatment technology, it can be concluded that there is no conformity with the Urban Wastewater Treatment Directive. The percentage of the population covered by municipal wastewater treatment with included biological treatment is very low. Therefore, the introduction of regular treatment of wastewaters in the country is top priority, both at local and national levels.

In the past period, no reduction in BOD 5 and in concentrations of ammonium in rivers (MK NI 019) has been observed in the Republic of Macedonia. At some monitoring stations, located on the rivers Crna Reka and Vardar, eutrophic water status with high BOD value was recorded. These results could reflect the status of inefficient treatment of urban and industrial wastewaters in the country, as well as the inadequate protection of river basins.

Figure 1. Treatment of wastewater from the public sewerage network





Figure 2. Percentage of population with and without public sewerage network



Data coverage: [excel](#)

Source: Republic Institute for Health Protection

## Assessment

According to the results on the distribution of the population in the Republic of Macedonia in relation to treated municipal wastewaters involving only mechanical treatment, biological treatment and application of latest treatment technology, and in relation to public sewerage, it can be concluded that the percentage of such population is very low. Despite of the rising trend, the current state is unsatisfactory with regard to EU requirements.

## Methodology

- Methodology for the indicator calculation

In accordance with the requirements of EUROSTAT

## Policy relevance of the indicator

### List of relevant policy documents

The National Environmental Action Plan (NEAP) 2.

Environmental Monitoring Strategy and Environmental Data Management Strategy.

Under the Urban Wastewater Treatment Directive, the EU Member States are required to provide connection to wastewater collection systems in all agglomerations exceeding 2 000 population equivalent. Secondary (biological) treatment must be provided in all agglomerations exceeding 2 000 population equivalent and discharging wastewater directly into receiving freshwater resources. It provides for specific requirements for different deadlines for compliance achievement depending on the sensitivity of receiving waters with regard to agglomerations exceeding 10 000 population equivalent.

The performance of wastewater treatment is monitored for five different parameters: BOD, COD, total suspended matter, total nitrates and total phosphorous.

In the case of smaller agglomerations and those connected to wastewater collection systems, the treated wastewater at the outlet has to comply with quality targets of the recipient.

### Legal grounds

The Law on Waters prescribes maintenance and improvement of water regime carried out on the basis of river basin management plans. Such Plans contain environmental protection objectives, good ecological status of surface water bodies (good quantitative and chemical status, including good ecological potential) and groundwater resources (good quantitative status and good chemical status).

River Basins Management Plans will be implemented through issuance of permits for water use, permits for extraction of sand, gravel and stone and permits for water discharges specifying quantitative and qualitative requirements in each case individually.

For the purpose of maintenance and improvement of the quality of water and establishment of the adequacy of water for use for different purposes, the Law on Waters specifies classification of waters and categorization of water bodies, as well as specification of deadline for achievement of the water quality goals for each water category and specification of the minimum standards for water quality and environmental protection goals for all water bodies. According to the Law, Programme with measures for environmental protection goals shall be adopted for each river basin.

The Decree on categorization of water courses, lakes, accumulations and water resources specifies the quality of water by specific classes of water in water bodies, lakes, accumulations and groundwater resources. This Decree also establishes five categories of water courses with regard to water quality goals specified therein. In order to maintain the quantity and the quality of the water (water regime), the competent authority issues water management consent in relation to construction, reconstruction, connection or extension of facilities that make impact on waters and water management permit for use of water as resource or as recipient. In circumstances of absence of emission standards for individual polluters and pollutants, the said documents are issued on the basis of specific expert assessment by the competent authority, for each case separately, taking into account the principle of sustainable use of water resources and providing care for the quality of wastewater discharged, in order to prevent the water to exceed the quality standards applied for the recipient

The Law on Public Health Protection specifies that the Regional Institutes for Health Protection,

coordinated by the Republic Institute for Health Protection, are obliged to monitor environmental-health and other conditions of relevance for the protection of drinking water and to undertake measures for active protection of the population against communicable and other diseases of high health and social relevance. These Institutes perform microbiological, parasitological, hygiene, toxicological and biochemical analyses within the scope of their activity.

Monitoring of the surface waters quantity is performed under the Programme for Preventive Public Health Protection, adopted at annual basis and published in the Official Gazette of the Republic of Macedonia. Monitoring of the surface waters quality at all points of health relevance is performed under the Programme for Preventive Public Health Protection, in order to enable undertaking of timely measures for public health protection. Waters used as drinking water sources, for sports and recreation, as well as for primary agriculture production, are of highest interest

The National Strategy for Waters is adopted to cover 30 period. It should provide sustainable development of waters, through meeting the demands of all users, protecting waters against pollution, protecting and improving aquatic ecosystems and providing protection against harmful impacts of waters. The Strategy should be adopted by the Assembly of the Republic of Macedonia.

## Targets

Requirements of the relevant EU Directives, (FWD, on urban wastewater treatment, on nitrates, on hazardous substances, as well as Directives on drinking and bathing waters) have been transposed in the Law on Waters, thus enabling reduction and prevention of water pollution and achievement of good ecological status or potential of waters.

The Urban Wastewater Treatment Directive, aimed at protecting the environment against impacts caused by urban wastewater discharges. In addition to this, compliance with the requirements specified in the Urban Wastewater Treatment Directive and Directive on Integrated Pollution Prevention and Control is incorporated in the goals of the Framework Water Directive, the main goal of which is the achievement of good chemical and biological status of all waters by 2015.

## Reporting obligation


- EUROSTAT

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MK NI 024</b>	<b>Urban wastewater treatment</b>	CSI 024	Urban waste water treatment	<b>P</b>	A	water waste	Annually

## **MK - NI 039**

### **DRINKING WATER QUALITY**



#### **Definition**

This indicator shows the exceedance of limit values set in Drinking Water Directive (80/778/EEC) and its amendment (98/83/EC which entered into force in 2003) and in the Rulebook on drinking water safety (Official Gazette of the Republic of Macedonia No.57/04), as well as the guideline values set for the quality of drinking water by the World Health Organization (WHO, 2004 and 2006).

Exceedance of drinking water quality limit values occurs when the concentration/dose of the pollutant exceeds the limit values specified in the above listed regulations.

Where more than one limit values exist (see the section on Policy goals), the indicator shall adopt the most strict case.

#### **Units**

- Number of aerobic mesophilic bacteria in 1 ml,
- Number of coliform bacteria in 100 ml,
- Number of thermo-tolerant coliform bacteria in 100 ml,
- Concentration of physico-chemical pollutants in mg/l,
- Parameters for radiological safety of drinking water in bekerels/l and total indicative dose in mSV/l.

#### **Key policy issue**


*What progress has been made in reducing the concentrations of pollutants in urban and rural environments in order to reach drinking water limit values specified in the Rulebook?*

#### **Key message**

Access to safe drinking water in the Republic of Macedonia amounts 94% (period from 2001 to 2013) with a note that population in urban areas has 99% access to safe drinking water and 78% of rural population has access to health safe drinking water, while the rest is exposed at occasional risk of bacteriological pollution of drinking water.



Figure 1. Drinking water quality in percentage



Data coverage: [excel](#)

Source: Public Health Institute of the Republic of Macedonia, PHI – 10 Regional

### Assessment

Sanitary and hygienic condition of facilities and health safety of analyzed water samples are, generally, satisfactory, i.e. within the expected limits compared to previous years. In the period 2001-2013, the percentage of unsafe samples based on physical and chemical analysis ranged between 3.4 and 7.5 %, while the percentage of unsafe samples based on microbiological analysis ranged between 0.8 and 2.99%. The most frequent causes of unsafe findings in the physical and chemical include absence of residual chlorine or increased content of iron in raw water and in very few samples it is due to increased content of nitrites from dag or drilled wells of individual users.

With bacteriologically positive findings, the cause is mostly increased number of aerobic mesophile bacteria. Toxic parameters are within the prescribed legal norms.

In the segment of health safe drinking water supply in rural populated places, the deficiencies in terms of undefined sanitary protection zones around drinking water sources, lack of adequate equipment for drinking water filtering and disinfection and inappropriate technical maintenance, have been constantly present. Therefore, there is high percentage of bacteriologically unsafe (9-25%).

### Methodology

- Methodology for the indicator calculation

The 10 regional Public Health Institutes – Skopje, Kumanovo, Veles, Shtip, Kochani, Strumica, Prilep, Bitola, Ohrid and Tetovo with their hygiene-epidemiological stations, in cooperation with the Public Health Institute – Skopje, carry out regular and continuous monitoring of drinking water corresponding with the number of measuring points and schedule specified in the Rulebook on drinking water safety (Official Gazette of the Republic of Macedonia No.57/04). Institutes perform basic physico-chemical and bacteriological analyses of drinking water samples, while the Public Health

Institute of the Republic of Macedonia performs monitoring of periodical physico-chemical analysis, analysis of pesticide residues, analyses of contaminants, parasitological and radiological analysis.

### *Uncertainty*

- **Methodological uncertainty**

Data is, generally, representative for the whole urban area in the Republic of Macedonia. The indicator is subject to modifications from year to year, depending on the introduction of new drinking water treatment plants and in line with the enhanced trend of rural population coverage with safe drinking water supply.

- **Data uncertainty**

Data is, generally, representative for the whole urban area in the Republic of Macedonia. Representativeness of monitoring selection is in accordance with the requirements of Directive 98/83/EC.

## **Policy relevance of the indicator**

### **List of relevant policy documents:**

The National Environmental Action Plan - 2 (2006) sets the improvement of the quality of drinking water through reduction of emissions of the main pollutants into surface and groundwaters as its main objective. The same document specifies the primary measure to be applied: to strengthen the processes of drinking water quality monitoring and assessment.

The 1999 National Environmental Health Action Plan (NEHAP) sets two main objectives:

- Reduction and minimization of health risks for the population through provision of drinking water for every citizen, which is safe from health point of view, sufficient in quantity, with guaranteed microbiological, organoleptical and physico-chemical composition, compliant with national standards and WHO Guidelines, as well as waters intended for sports and recreation and healthy food production;
- Reduction of exposure to toxic chemicals through water originating from agriculture and industry.

The NEHAP also sets the following priorities:

- approximation of the legislation on the quality of ambient and drinking waters with the recommendations of the EU (approximation completed in 2004) and with the WHO Guidelines;
- introduction of disincentive prices for non-earmarked consumption of drinking water by commercial and non-commercial users and restrictive prices for the population in circumstances of draught for the purpose of consumption streamlining (implemented under the Law on Drinking Water Supply and Urban Wastewater Collection);
- establishment of sanitary protection zones around water supply sources in order to prevent contamination of anthropogenic origin (permanent process performed and most of the public utilities have established zones in line with the Elaborates for sanitary protection zones developed by the Public Health Institution RIHP and other authorised vocational institutions);
- completion of the process of construction of municipal and industrial wastewater treatment systems;

- monitoring of the quality of surface and groundwaters, especially at drinking water abstraction, places intended for sports and recreation and points for abstraction of water for irrigation, monitoring of discharged untreated and treated municipal and industrial wastewaters in accordance with EU and WHO Guidelines (monitoring is performed regularly and continuously by the Public Health Institute - Skopje and the 10 Regional public health centers with their local units);
- although the pilot project for fluoridation of milk consumed by pre-school children has been initiated, introduction of drinking water fluoridation as the most efficient, the least costly and socially and medically most fair means for massive caries prophylaxis has remained as public health option.

## Legal grounds

Law on Health Protection, Law on Waters, Programme for preventive health protection in the Republic of Macedonia, Law on Drinking Water Supply and Urban Wastewater Collection, Decree on Water Classification, which in its Article 2, specifies five classes of surface watercourses, lakes and accumulations and ground water resources.

The Law on Food and Foodstuffs and Materials in Contact with Food, in its Article 4 includes drinking water as food.

Rulebook on drinking water safety (sets frequency of drinking water safety control).

In the Law on Nature Protection, one of the main goals defined in Article 4, item 6 of the Law is the securing of the right of citizens to a healthy environment.

The following EU Directives have been transposed in the new legal acts:

Drinking Water Directive (80/778/EEC) and its amendment (98/83/EC which entered into force in 2003).

## Targets

The Rulebook on drinking water safety specifies the limit values for the parameters monitored in drinking water in terms of human health protection.

### ***Limit values of concentrations of certain parameters in drinking water***

- According to the said Rulebook, limit values have been specified for the purpose of human health protection, harmonized with the EU Directive and WHO Guidelines on the quality of drinking water (2004).

## Reporting obligation

European Environmental Agency

- Exchange of data on drinking water quality, based on the Council Decision on the establishment of reciprocal exchange of information and data on drinking water quality (98/83/EC).

World Health Organization - ENHIS


- Drinking water quality, in line with the WHO Guidelines on drinking water quality of 1987 and 2004, respectively.

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MKNI 039</b>	<b>Drinking water quality</b>	WEU13	Drinking water quality	<b>S</b>	<b>A</b>	Water quality	Annually

## MK - NI 040

### IRRIGATED LAND



## Definition

The indicator tracks the trend in irrigated areas in a given time interval on the whole territory of the Republic of Macedonia, as well as total quantities of consumed water on the entire territory and proportion of irrigated land compared to the total cultivable land area.

## Units

- Area of irrigated land (expressed in hectares), quantity of water used for irrigation expressed in cubic meters consumed at annual level, % of irrigated land in the total cultivable land area.

## Key policy issue

*Is the water abstraction based on water sustainability?*

## Key message

An uneven trend in water use for land irrigation was observed in the period between 2001 and 2013, due to weather conditions in the given year, as well as to organizational restructuring of the sector. Particular growth in water use for land irrigation was recorded in 2012.

Data is not part of the official statistics published in the country.

Figure 1. Use of water resources




Figure 2. Total area irrigated





Figure 3. Percentage of irrigated area relative to total cultivated land area



Data coverage: [excel](#)

Source: State Statistical Office

## Assessment

Figure 2 shows the whole irrigated land area for the period 2001-2013, reflecting an uneven trend in water use for land irrigation. There is an evident data that the quantity of water consumed by this sector in 2012 was significantly bigger compared to the entire successive interval. This is due to the fact that 2012 was a dry year, but precipitations were distributed in such a way that enables filling of accumulations with sufficient quantities of water. Figure 3 presents the percentage of irrigated land area compared to the entire cultivable land area in the Republic of Macedonia, showing that the percentage is really low with the average being below 5% for the entire time interval, except in 2012 when higher percentage was recorded, reaching 6.2%.

## Methodology

- Methodology for the indicator calculation

Data is collected and processed by years.

## Policy relevance of the indicator

### List of relevant policy documents:

The National Environmental Action Plan - 2 and Environmental Monitoring Strategy and Data Management Strategy.

The policy for sustainable use of water resources based on the Sixth Environmental Action Programme and Framework Water Directive requirements as transposed in the national Law on Waters.

### Legal grounds

The Law on Waters provides for integrated approach, specifying the conditions and the manner of waters use and allocation, protection against harmful impacts of water, as well as standards and values for water quality and control of pollution, while taking into account integration of measures and activities for water protection in all development, strategic, planning and programme documents.

The main planning documents for water management planning and development include:

- The National Strategy for Waters
- Water Management Master Plan of the Republic of Macedonia, and
- River basin management plans.

The National Strategy for Waters is aimed at establishing long-term policy to ensure sustainable use of water by meeting the demands of all users with adequate quality water in sufficient quantities, rational and cost-effective consumption of waters, water protection against contamination and contamination control.

The Water Master Plan of the Republic of Macedonia provides for integrated planning and implementation of programmes and measures, technical and economic solutions for rational water use, protection of waters against contamination and protection against harmful impacts of water, based on the principles of sustainable development and the timeframe for their implementation.

The River Basins Management Plans enable maintenance and improvement of water regime. Such Plans contain the environmental protection goals, good status of surface water bodies (good quantitative status and chemical status, including good ecological potential) and ground water resources (good quantitative status and chemical status).

Use of water for different purposes is specified under the Decree on Water Classification, according to which water is divided into five different classes based on the level of pollution, while water characteristics are determined on the basis of classes and purposes for which water can be used.

## Targets

No specific targets.

## Reporting obligation

OECD/EUROSTAT

## General metadata


Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MKNI 040</b>	<b>Irrigated land</b>	WQ4	Irrigated land	<b>D</b>	A	Water	Annually



# AGRICULTURE



## MK - NI 08 MINERAL FERTILIZER CONSUMPTION



### Definition

Mineral fertilizers are substances containing chemical elements required for plants growth, especially nitrogen, phosphorus and potassium.

This indicator shows the consumption of mineral fertilizers in the Republic of Macedonia, by presenting total amounts in tonnes consumed substances, and their application per hectare cultivated land area.

### Units

- Tones,kg/ha.

### Key policy issue

**What is the trend in the amount of used mineral fertilizers in agriculture?**

### Key message

Mineral fertilizer consumption in agriculture noted a falling trend in the period from 2000 to 2012. The overall mineral fertilizers consumption reduced by 64.61%. Consumption of nitrogen mineral fertilizers dropped by 63.77%. Consumption of phosphorous fertilizers dropped by 77.01%. Consumption of combined mineral fertilizers dropped by 93.26% between 2000 and 2009, while in the period from 2009 to 2012 there was gradual increase by 11.82%. Consumption of potassium fertilizers showed periodical trends of reduction and increase, so that consumption in 2012 increased by 100% compared to 2011. Consumption of mineral fertilizers on cultivated land area (kg/ha) by agricultural companies and agricultural cooperatives during the observed period showed periodical trends of reduction and increase.

Figure 1. Consumption of mineral fertilizers





Figure 2. Use of mineral fertilizers on cultivated land area (kg/ha) by agricultural cooperatives and agricultural companies)



Data coverage: [excel](#)

Source: Statistical Yearbooks, State Statistical Office

## Assessment

In the observed period, consumption of mineral fertilizers in agriculture dropped from 16.16 tons to 5.809 tons of fertilizers. The quantity of mineral fertilizers used on cultivated land area (of agricultural companies and cooperatives) expressed in kilograms per hectare, during the observed period showed periodical trends of increase and decrease. In 2004, with 77.74 kg/ha, consumption of mineral fertilizers was the lowest, while in 2006 with 147.24 kg/ha it was the highest.

It is difficult to connect the trend in reduction in mineral fertilizers consumption directly with the impact on the quality of the environment. The ultimate effect on the quality of environment depends to a great extent on other factors, such as use of organic fertilizer, yield from crops, soil types, management of agricultural farms, etc.

## Methodology

- Methodology for the indicator calculation

Consumption of individual groups of mineral fertilizers as combined mineral fertilizers, nitrogen fertilizers, phosphorous fertilizers, potassium fertilizers, as well as total mineral fertilizers per hectare utilized agricultural area is obtained by dividing the total quantity of consumed group of mineral fertilizers in kg by the total utilized agricultural area presented in ha.

## Policy relevance of the indicator

### List of relevant policy documents:

The Second National Environmental Action Plan (NEAP 2) specifies the measure for rationale use of natural resources, as well as controlled use of mineral fertilizers. The same document also specifies the measure for establishment of monitoring and information system for soil, to monitor the mineral fertilizers consumption.

### Legal grounds

The Law on Agricultural Land specifies the measures for improved agricultural land fertility through undertaking of agrotechnical measures, one of them being fertilizers application, i.e. use of mineral fertilizers. It is specified that agricultural land protection against pollution and contamination is performed by prohibition, restriction and prevention of direct input of harmful matters in soil, input of harmful matters through water and air and undertaking of other measures for its productivity maintenance and improvement. It is also specified that, for the purpose of agricultural land protection against pollution and contamination, the provisions contained in the regulations on environment and nature protection and improvement shall apply accordingly.

The Law on Fertilizers Use regulates the manner of fertilizers use.

The Law on Nature Protection, in its Article 4, specifies the goals of protection, including preservation and recovery of existing biological and landscape diversity in a state of natural balance and prevention of harmful activities and nature disruption.

## Targets


No specific targets.

## Reporting obligation

No reporting obligation

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MKNI 08	Mineral fertilizer consumption	IRENA 08	Mineral fertilizer consumption	D		Agriculture Soil Water	Annually



### Definition

Plants protection products or pesticides are chemical substances which restrain diseases and pests in plants. This indicator shows the quantities of pesticides used for plants protection, such as fungicides, herbicides, insecticides and category of total including, apart from the mentioned ones, other plant protection products.

### Units

- Total quantities of used substances in tones, share of different groups of pesticides, as well as their application per hectare utilized agricultural area (kg/ha).

### Key policy issue

*Has the use of pesticides in agriculture increased in quantity?*

### Key message

Application of pesticides in agriculture, including all plant protection products, like fungicides, herbicides, insecticides and total quantity showed a trend of reduction in quantity consumed in the period between 2000 and 2005, sharp increase in 2006 and then decrease again by 2012.

The application of fungicides from 2000 to 2006 showed trend of variation of reductions and increases, in the period from 2006 to 2012 it reduced by 77.66%. Application of herbicides also showed a trend of reduction from 2000 to 2006, while from 2006 to 2012 the overall use of pesticides in agriculture reduced by 71.72%.

Figure 1. Use of plant protection products




Figure 2. Share of plant protection products in percentage





Figure 3. Total plant protection products used on the total cultivable land (kg/ha) (from agricultural companies and agricultural cooperatives)



Data coverage: [excel](#)

Source: Statistical Yearbooks, State Statistical Office

### Assessment

In the period 2000 to 2006, use of pesticides in agriculture showed variations of reduction and increase in the period from 2006 to 2012 it dropped from 336 to 95 tons. With regard to the share of plant protection products, in the period 2000 to 2012, fungicides noted highest share. In 2012, fungicides were the most used with 68.42%, then insecticides with 21% and herbicides with

10.52%.

The total amount of plant protection products used on the total cultivable land in agricultural companies and agricultural cooperatives expressed in kg/ha, from 2006 when consumption was the highest, to 2012 reduced from 5.08 to 2.09 kg/ha, which is a reduction by 58.85 %.

## Methodology

- Methodology for the indicator calculation

The share of different pesticide groups as fungicides, herbicides and insecticides is obtained when the quantity of each group is divided by the total quantity of consumed pesticides, and then the value obtained is multiplied by 100. The application of individual group per hectare utilized agricultural area is obtained when the total quantity of consumed pesticides expressed in kg is divided by the total utilized agricultural area (agricultural companies and agricultural cooperatives) in the Republic of Macedonia expressed in ha.

## Policy relevance of the indicator

### List of relevant policy documents:

The Second National Environmental Action Plan (NEAP 2) specifies the measure for rationale use of natural resources, as well as controlled use of pesticides, i.e. plant protection products. The same document also specifies the measure for establishment of monitoring and information system for soil, to monitor the pesticides consumption.

### Legal grounds

The Law on Agricultural Land specifies the measures for improved agricultural land fertility through undertaking of agro-technical measures, hydro-amelioration, agramelioration and anti-erosion measures.

The Law on Nature Protection specifies the goals of protection, including preservation and recovery of existing biological and landscape diversity in a state of natural balance and prevention of harmful activities and nature disruption.

The Law on Plants Protection regulates the protection of plants against diseases, pests and weeds, as well as use of plant protection products.

The Law on Plant Protection Products regulates approval, placement on the market, use and control of active substances that are products; maximum level of residues, equipment for products application; exchange of information related to products, products production, records keeping of legal and natural persons involved in production and placement of products on the market, conditions for authorization by authorities responsible for implementation, monitoring and control of this law.

The Law on Plant Health regulates the health of plants, measures and obligations concerning occurrence of harmful organisms in plants, plant products and other articles and objects, prevention of their inlet and spread, measures for control, biological measures for plant protection, access to and exchange of information and information system, costs and compensations, responsibilities of competent authorities, authorized services, authorities and bodies in the area of plants health and it also regulates other issues in the area of plants health.

## Targets

No specific targets.

## Reporting obligation

None

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MKNI 09</b>	<b>Consumption of pesticides</b>	IRENA 09	Consumption of pesticides	<b>D</b>		Agriculture	Annually





### Definition

The nutrient balance or nitrogen balance establishes the link between nutrients used in agriculture and changes in the quality of the environment, in order to achieve sustainable use of soil nutrients in terms of their input and output.

The indicator estimates the potential surplus of nitrogen on agricultural land. This is done by calculating the balance between nitrogen added to a hectare agricultural land. The indicator accounts for all inputs to and outputs from the farm. The inputs consist of the amount of nitrogen applied via mineral fertilisers and animal manure as well as nitrogen fixation by legumes, deposition from the air, and some other minor sources. Nitrogen output is contained in the harvested crops, or grass and crops eaten by livestock. Uncontrolled escape of nitrogen to the atmosphere, e.g. as N<sub>2</sub>O from agriculture is difficult to estimate and therefore not taken into account.

### Units

- The gross nitrogen balance is expressed in (kgN/year) per hectare (ha).


### Key policy issue

*Has the impact of agriculture on the environment improved?*

### Key message

In the period from 2000 to 2001, significant rise in gross nitrogen balance was tracked, followed by a period of stagnation, to note rapid drop in the period from 2002 to 2003 and further slight fall in the gross nitrogen balance expressed both in thousand tones and kilograms nitrogen per hectare agricultural land (kgN/ha). Constant surplus in nitrogen balance indicates potential environmental problems, while constant deficit indicates potential risk of reduced nutrients in the soil.

Figure 1 Gross nitrogen balance expressed in thousand tones and kgN/ha agricultural land area



Data coverage: **excel**

**Source:** Statistical Yearbook, State Statistical Office, Calculation of the gross nitrogen balance was made by the Faculty of Agricultural Science and Food, University of "St. Cyril and Methodius", Skopje

## Methodology

- Methodology for the indicator calculation

Methodology for the indicator calculation has been taken from OECD/Eurostat national nutrient balances, which takes into account all input and output matters in the farm.

Input nitrogen matters consist of:

1. Total amount of applied fertilizers
  - Inorganic fertilizers
  - Organic fertilizers (manure excluded)
2. Manure
3. Nitrogen fixation by legumes
4. Deposition from the air
5. Other minor sources (semen and other reproductive material)

Output nitrogen matters include:

1. Harvested crops placed on the market, including also forage crops
2. Grass and crops eaten by livestock

Uncontrolled escape of nitrogen to the atmosphere in a form of  $N_2O$  from agriculture is difficult to estimate and therefore not taken into account.

- Source of applied methodology

OECD/Eurostat Gross Nitrogen Balances Handbook (12/2003)

## Uncertainty

- Methodological uncertainty

Data used in the calculation of this indicator has been partially based on estimates by experts, using harmonized methodology which might not always reflect specific circumstances in our country. Certain coefficients used in calculations differ significantly from country to country. Data on nitrogen input is considered more adequate and more comprehensive than on the output. Uncertainty is present with regard to harvested forage crops, as well as grass crops eaten by livestock.

Based on the above, data on gross nitrogen balance in our country should be taken with certain extent of precaution.

- Uncertainty of data sets

Data on the amounts of applied manure are accompanied by certain extent of uncertainty, and statistical data on semen and other reproductive material, as well as data on grass crops eaten by livestock, i.e. those that are not placed on the market, should be taken by certain extent of uncertainty, too.

## Policy relevance of the indicator

The gross nitrogen balance is an issue regulated by the Framework Law on Waters, incorporating the requirements of Nitrates Directive (91/676/EC) and Framework Water

Directive (2000/60/EC). The Nitrates Directive is aimed at reducing and preventing in future the water pollution by nitrates from agricultural sources. This Directive restricts the application of manure at 170 kg N/ha/year. The goal of the Framework Water Directive is the achievement of good ecological status of surface and ground waters in terms of quality of biological communities, hydrological characteristics and chemical characteristics.

## List of relevant policy documents

The National Environmental Action Plan 2 (NEAP 2) specifies the measures for rational use of natural resources, as well as controlled use of pesticides, i.e. plant protection products, as well as the measure for establishment of soil monitoring and information system to monitor the status of pesticides consumption.

## Legal grounds

The Law on Agricultural Land specifies the measures for improved agricultural land fertility through undertaking of agro-technical measures, hydro-amelioration, agramelioration and anti-erosion measures. It also prescribes that for the purpose of protection of agricultural land against pollution and contamination, the provisions contained in the regulations on environment and nature protection and improvement shall apply accordingly.

The Law on Nature Protection, in its Article 4, specifies the goals of protection, including preservation and recovery of existing biological and landscape diversity in a state of natural balance and prevention of harmful activities and nature disruption.

The Law on Environment, in its Article 8, promotes the principle of sustainable development, meaning that, when undertaking or performing any activity, rationale and sustainable use of natural resources shall be taken into account, thus meeting the needs for healthy environment, as well as social and economic needs of present generations, without jeopardizing the rights of future generations to meet their own needs.

## Targets

No specific targets.

## Reporting obligation


EEA

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 025</b>	<b>Gross nitrogen balance</b>	CSI 025	Gross nutrient balance	<b>P</b>	A	<ul style="list-style-type: none"> <li>▪ agriculture</li> <li>▪ water</li> </ul>	3 - annually

## MK - NI 026

### AREAS UNDER ORGANIC FARMING



### Definition

The indicator is calculated as share (percentage) of area under organic farming (sum of existing areas under organic farming and areas in a process of conversion for organic farming) in the total area or total cultivable land area.

### Units

The indicator is presented as sum of area under organic farming and area being converted for organic farming, measured in ha. Share of organic farming is given as a percentage of total utilized agricultural area.

### Key policy issue

***Whether the share of organic cultivable area in the total cultivable area is in increase?***

### Key message

In the period from 2005 to 2011, the organic agricultural production area and the number of organic operators is in constant increase, but in 2012 and 2013 the trend decreased.

In the period from 2005 to 2011, the organic agricultural production area increased from 592.54 to 6.580,92 ha, in 2012 there was drop to 4.663.08 hectares, and in 2013 it dropped to 3168.16 hectares.

The share of organic farming in the overall cultivable land area had significant increase, namely from 0.109% in 2005 raised to 1.288% in 2011. The share noted drop in 2012 and 2013 by 51.7% compared to 2011 and in 2013 it amounted 0.622%.

The number of certified organic operators increased from 50 in 2005 to 780 in 2011, and in 2012 and 2013 it dropped to 576 and 400 operators, respectively.

Figure 1. Area with organic agricultural production




Figure 2. Share of area with organic agricultural production in cultivable and total agricultural area.




Figure 3. Organic vegetable production in ha





Figure 4. Ratio between the number of operators and the area under organic agricultural production



Data coverage: [excel](#)

Source: Statistical Yearbook, State Statistical Office, Ministry of Agriculture, Forestry and Water Economy, Division of Organic farming.

## Assessment

In the period from 2005 to 2011, the production areas with organic farming have increased from 266 ha to 1007.26 ha subsequently. There is also an increase in areas under conversion from 327 to 5.573.66 ha, the trend in 2012 showed reduction compared to 2011 in areas under conversion from 5.573.66 hectares to 2.817.02 hectares, and in 2013 to 3.168.16 hectares.

The share of organic farming areas in the total cultivable area grew from 0.109% in 2005 to 1.288% in 2011, while in 2012 the share dropped and amounted 0.914%; negative trend continued in 2013 and the share of organic farming areas in the total cultivable area was 0.622% which was a drop by 51.7% compared to 2011 when the share was highest. .

The share of organic farming areas in the total cultivable area was insignificant given that, according to the National Plan for Organic Production 2013-2020, the target of 4 % share of organic cultivable area should be achieved in Macedonia by 2020.

It can be noted from Figure 3 that cereals were the main organic crop in Macedonia in 2013, with share of 49.17%, followed by forage with 21.82 % share, while the aromatic/medicinal crops had the lowest share with 0,69 % of the overall certified areas.

The number of certified organic operators in the period from 2005 to 2011 grew proportionally with the growth of the area under organic agricultural farming (figure 4), and dropped in 2012 and 2013.

## Methodology

The method of the European Environmental Agency.

The indicator is presented as sum of area under organic farming and area being converted

for organic farming, divided by the total cultivable land area or total agricultural area. This value is multiplied by 100 in order to present the value in percentage.

## Policy relevance of the indicator

### List of relevant political documents

- National strategy with action plan for development of organic farming in the Republic of Macedonia 2008-2011
- Strategy for agriculture and rural development – condition and need to establish agro-environmental policy.
- The Strategy for compliance of the Macedonian agricultural and food sector with the EU Common agricultural policy – indicates the need to identify the regions suitable for organic farming and to comply with the European regulations.
- The Treaty with the World Trade Organization – improvement of the international food safety standards
- The Second National Environmental Action Plan – strives to include environmental issues directly in the agricultural development policy and to maintain the natural resources needed for sustainable development on high level. In accordance with this, a control and certification body for organic agricultural production, recognized by the EU has been defined and established.
- National strategy for biodiversity with Action Plan
- National strategy for sustainable development

### Legal grounds

The framework for the organic farming is established by the Law on Organic Farming and regulations which are in a process of adoption; the Law on Stimulating Agriculture Development, Law on Environment and Law on Nature Protection.

The provisions of this Law have been harmonized with international and European ones, especially with EU Regulation No.2092/91, which is of particular importance in the context of future development of trade exchange in organic products with European countries.

### Targets

In 2011, the organic cultivable land has a 2% share in the total cultivable land in Macedonia. In 2011, while the wild collection area has a share of 5% in the total cultivable area in Macedonia.

### Reporting responsibilities

- Annual report for quality of the environment in the Republic of Macedonia
- Environmental statistics
- European Environment Agency


### General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MKNI 026</b>	<b>Area under organic farming</b>	CSI 026 IRENA 07	Area under organic farming	<b>R</b>	A	agriculture biological diversity	Annually

# ENERGY







### Definition

Final energy consumption is energy supplied to meet the demand of the final consumers and is calculated as the sum of final energy consumption from all sectors, namely industry, transport, agriculture, households, and other sectors.

The indicator "Final energy consumption by sector" is expressed in thousand tonnes of oil equivalent (ktoe) and in percentage as a ratio between final energy consumption by each sector and final energy consumption by all sectors.

### Units

- thousand tonnes of oil equivalent (ktoe)
- percentage (%)

### Key question

Is final energy consumption increasing and in which sector it is the highest?

### Key message

Policies in energy sector should favour measures aimed at rational and efficient energy consumption, especially by households and industry.

During the period between 2000 and 2013, final energy consumption in the Republic of Macedonia increased by 12.73%, with an annual average rate of 1.29%. Non-energy consumption is the sector with the most rapid growth in energy consumption noting an increase by 1119.36%, followed by other sectors with 37.49%. During the same period, final energy consumption in transport increased by around 35.47%, while final energy consumption in industry increased by 5.24%. Significant fall in final energy consumption was recorded in the sector agriculture (-61.23%) and households (-6.45%).

The highest share in the total final energy consumption was recorded in the sectors industry with 31% and transport with 27.5%.

Figure 1. Final energy consumption by sector





Figure 2. Share of individual sectors in final energy consumption



Data coverage: [excel](#)

Source: State Statistical Office

## Assessment

Final energy consumption in the Republic of Macedonia correlates closely with the dynamics of industrial production, due to the high share of industry in the total energy consumption.

In the period between 2000 and 2013, final energy consumption in the Republic of Macedonia increased by 12.73%, with an annual average rate of 1.29%. Non-energy consumption is the sector with the most rapid growth in energy consumption noting an increase by 1119.36%, followed by other sectors with 37.49%. Analysis of data on final energy consumption in 2009 indicates fall in industry as a result of economic crisis, and the biggest increase in industry sector was recorded in 2007. During the same period, final energy consumption in transport noted constant growth and increased by around 35.47%. Final energy consumption in households decreased by 6.45% in the period 2000 to 2013, but significant increase was recorded 2008 to 2011. Significant fall in final energy consumption was recorded in the sector agriculture (-61.23%), with highest consumption in 2004, and then in the period 2005 to 2011 there was constant fall in consumption.

The highest share in the total final energy consumption was recorded in the sectors industry with 31% and transport with 27.5%, followed by households (25%) and other sectors (12.2%), non-energy consumption (3.1%) and agriculture (1.2%).

## Methodology

- Methodology for the indicator calculation

Statistical methodology for calculation:

- Regulation on Energy Statistics of the European Parliament and of the Council (Regulation no.1099/2008).
- "Energy Statistics Methodology Eurostat F4, 1998"
- National classification of activities, Rev.2 (Official Gazette of the Republic of Macedonia no. 147/2008)

## Policy relevance of the indicator

- Strategy for Energy Efficiency Promotion in the Republic of Macedonia by 2020<sup>1</sup>
- Strategy for Energy Development in the Republic of Macedonia by 2030.<sup>2</sup>

## Legal grounds

Law on Energy; Energy Balance of the Republic of Macedonia - annual planning document defining the demands for energy and the possibility for their supply .

## Reporting obligation

- Eurostat
- ECE/UN
- IEA/OECD

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MK NI 027</b>	<b>Final energy consumption by sector</b>	CSI 027 EE 18	Final energy consumption by sector	<b>D</b>	A	Energy	annually

<sup>1</sup> <http://www.konkurentnost.mk/StrateskiDokumenti/StrategijazaunapreduvanjenaEERMdo2020godina.pdf>

<sup>2</sup> <http://www.build.mk/docs/users/cloverstack/Strategija%20za%20razvoj%20na%20energetikata%202008-2020%20so%20vizija%20do%202030.pdf>

## **MK - NI 028**

### **TOTAL ENERGY INTENSITY**



#### **Definition**

Total energy intensity is the ratio between the total energy demand (or total energy consumption) and the Gross Domestic Product.

The total energy consumption is calculated as the sum of the total energy demand from solid fuels, oil, natural gas and renewable sources.

The Gross Domestic Product (GDP) is converted by the Price Adjusted Rate of Exchange (PARE) method, applying the OUN Methodology (2000 database).

The total energy demanded (or total energy consumption) is expressed in thousand tonnes oil equivalent, and Gross Domestic Product in million EUR.

The indicator "Total energy intensity" is expressed in kilograms oil equivalent per 1000 EUR (kgoe/1000 EUR).

The indicator is also calculated in indexes with 2000 as base year (2000=100).

#### **Units**

- million EUR
- thousand tonnes oil equivalent (ktoe)
- kilograms oil equivalent (kgoe)
- indexes (2000=100)


#### **Key question**

What will be the dynamics of strategic targets implementation and achievement?

#### **Key message**

The trend in Energy intensity in the Republic of Macedonia recorded decline by 50.3% in 2013 compared to 2000, mainly due to the trend of growth in GDP for the same past period amounting 98%. The biggest decline in energy intensity compared to previous year, amounting 14.5% was recorded in 2013 relative to 2012, and the biggest increase of 5.5% was noted in 2002 relative to 2001.

Figure 1. Total energy intensity



Data coverage: [excel](#)

Source: State Statistical Office

## Assessment

Constant change in the trend of total energy intensity was notable in the period from 2000 to 2013 at an average rate of decline of 5.11%. The trend of decline by 50.3% of total energy intensity in 2013 compared to 2000 was specific, mainly due to the growth of GDP for the same past period reaching 98%.

The biggest decline in energy intensity compared to previous year, by 14.4%, was recorded in 2013 relative to 2012, and the biggest growth by 5.51% was noted in 2002 relative to 2001.

Comparative analysis of energy consumption relative to GDP, so called indicator of energy intensity, has shown that the Republic of Macedonia belongs to the group of countries with relatively high energy consumption levels, due to the high energy intensity of the facilities that lead the economic growth. Also, due to the long-lasting treatment of the electricity price as a social category, the residential sector uses significant quantities of electricity for heating.

## Methodology

- Methodology for the indicator calculation

Statistical methodology for calculation:

- Regulation on Energy Statistics of the European Parliament and of the Council (Regulation no.1099/2008),
- „Energy Statistics Methodology Eurostat F4, 1998"
- National classification of activities NCA Rev.2 (Official Gazette of the Republic of Macedonia no. 147/2008).

## Policy relevance of the indicator

### List of relevant policy documents:

- Strategy for Energy Efficiency Promotion in the Republic of Macedonia by 2020<sup>1</sup>
- Strategy for Energy Development in the Republic of Macedonia by 2030<sup>2</sup>

### Legal grounds

Law on Energy; Energy Balance of the Republic of Macedonia - annual planning document defining the demands for energy and the possibility for their supply (Article 16 of the Law on Energy).

## Targets

Target to be achieved in EU is spending tonnes oil equivalent per 1.000 US\$ GDP, while the target in the Republic of Macedonia is 0,75 tonnes oil equivalent. The implementation of measures under the Strategy for Energy Efficiency Promotion, this should drop down to 0,45 to 0,49 into 2020.

## Reporting obligation

- Eurostat
- ECE/UN
- IEA/OECD

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 028	<b>Total energy intensity</b>	CSI 028 EE 23	Total energy intensity	R	B	▪ energy	annually

<sup>1</sup> <http://www.konkurentnost.mk/StrateskiDokumenti/StrategijazaunapreduvanjenaEERMdo2020godina.pdf>

<sup>2</sup> <http://www.build.mk/docs/users/cloverstack/Strategija%20za%20razvoj%20na%20energetikata%202008-2020%20so%20vizija%20do%202030.pdf>



**Definition**

Total energy consumption or gross inland consumption represents the total quantity of energy necessary to satisfy the total national needs for energy for energy transformations, all types of consumption by energy sector and final energy and non-energy consumption.

The total energy consumption is calculated as sum of the total energy consumption originating from solid fuels, oil, natural gas and renewable sources.

The indicator "Total energy consumption by fuel" is expressed in thousand tonnes of oil equivalent (ktoe) and in percentage as ratio between the total energy consumption per fuel and the total energy consumption of all fuels.

**Units**

- thousand tonnes of oil equivalent (ktoe)
- percentage (%)

**Key question**

What are the trends concerning the share of fuels in the total energy consumption?

**Key message**

In the period between 2000 and 2013, the total energy consumption by fuel decreased by 1.56%. The highest share in the total energy consumption belongs to solid fuels ranging from 43% to 58%. For the period 2000 to 2013, there was significant increase in the amounts of natural gas used as one of the fuels with cleaner ecological footprint, noting increase by 140.5%. Use of oil shows obviously constant values, except in 2002 (1173 ktoe) and 2007 (1042 ktoe), when increase in the use of oil was recorded.

Significant change in the trend of solid fuels use and gradual structural substitution with cleaner or renewable energy sources is necessary.

Figure 1. Total energy consumption by fuel





Figure 2. Share by fuel in the total energy consumption



Data coverage: [excel](#)

Source: State Statistical Office

## Assessment

In the past period, dominant energy sources in the Republic of Macedonia included solid fuels, oil and oil derivatives. The share of solid fuels in the total energy consumption for the period between 2000 and 2013 was significantly higher compared to other energy sources ranging from 43% to 58%. Also, the share of liquid fuels in the total energy consumption was rather significant and ranged between 29.5% and 40.6%. Reduction of ecological footprint of electricity production in thermal power plants fueled by low calorific coal – lignite is among the greatest challenges in the attempt to reduce the effects of gas emissions in the production process.

Renewable energy sources have negligible share in the total energy consumption ranging from 7.8% to 12.6%, except for wood which is mostly used in individual households without adequate filters for combustion gas emissions.

Increase in the share of renewable energy sources in total energy consumption would be at the same time a possibility for economy restructuring in several sectors, energy efficiency in households and industry, as well as jobs creation.

## Methodology

- Methodology for the indicator calculation

Statistical methodology for calculation:

- Regulation on Energy Statistics of the European Parliament and of the Council (Regulation no.1099/2008),
- "Energy Statistics Methodology Eurostat F4, 1998"

## Policy relevance of the indicator

- National Strategy for Sustainable Development in the Republic of Macedonia 2009 – 2030<sup>1</sup>
- Strategy for Energy Efficiency in the Republic of Macedonia by 2020<sup>2</sup>

<sup>1</sup> <http://www.moepp.gov.mk/wp-content/uploads/2014/12/Nacionalna-Strategija-za-Odrzliv-Razvoj-vo-RM-NSSD-Del-1.pdf>

<sup>2</sup> <http://www.konkurentnost.mk/StrateskiDokumenti/StrategijazaunapredovanjenaEERMdo2020godina.pdf>



- Strategy for Energy Development in the Republic of Macedonia by 2030<sup>3</sup>

## Legal grounds

Law on Energy; Energy Balance of the Republic of Macedonia - annual planning document defining the demands for energy and the possibility for their supply (Article 16 of the Law on Energy).

## Targets

- Reduction in the dependence on imported fuels and reduction in inefficient energy consumption;
- Modernization of energy infrastructure and diversification of energy supply (extension of the network for natural gas is an important essential element for the implementation of all measures envisaged towards energy efficiency);
- Participation in of regional cooperation and compliance with the legislation of the Energy Community.


## Reporting obligation

- Eurostat
- ECE/UN
- IEA/OECD

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 029</b>	<b>Total energy consumption by fuel</b>	CSI 029 EE 24	Total energy consumption by fuel	<b>D</b>	A	energy	annually

<sup>3</sup> <http://www.build.mk/docs/users/cloverstack/Strategija%20za%20razvoj%20na%20energetikata%202008-2020%20so%20vizija%20do%202030.pdf>



**Definition**

Renewable energy sources are defined as renewable non-fossil energy sources: hydropower, geothermal, solar and wind power; solid biomass; biogas, liquid biofuels, etc.

The indicator "Renewable energy consumption" is expressed as ratio of total renewable energy consumption and the total energy consumption originating from all fuels (in percentage).

**Units**

- thousand tonnes of oil equivalent (ktoe)
- percentage (%)

**Key question**

How fast is the share of renewable energy in the total energy consumption?

**Key message**

Policies in energy sector should favour measures for greater use of renewable energy sources.

Relatively low share of renewable energy in the total energy consumption (10.33% at an average) indicates dominant use of fossile fuels which is unfavourable in terms of both depletion of energy resources and environmental pollution.

Biomass has the highest share of renewable energy in the total energy consumption and ranges from 4.6% to 7.7%, while the lowest share belongs to solar electric energy ranging between 0.003% and 0.029%. Hydro electricity has a share in the range between 2 and 7.3%.

Figure 1. Share of renewable energy in the total energy consumption by energy source (%)





Figure 2. Total share of renewable energy in the total energy consumption (%)



Data coverage: [excel](#)

Source: State Statistical Office

## Assessment

Relatively low share of renewable energy in the total energy consumption (10.33% at an average) indicates the dominant use of fossil fuels which is unfavorable in terms of both energy resources depletion and environment pollution. The highest share of renewable energy in the total energy consumption was recorded in 2010 with 14.6%, and the lowest one of 7.8% in 2012 and 2007.

In the period from 2000 to 2002, there was a drop by 34.6%, and from 2002 to 2006 there was a growth by 39.6% in the share of renewable energy in the total energy consumption, followed by decline in 2007 by 28%, and then growth again in the period from 2008 to 2010 of 73.5%, and significant decline in the period from 2011 to 2013 amounting 26.1%.

Biomass has the highest share of renewable energy share in the total energy consumption and ranges from 4.6% to 7.7%, while the lowest share belongs to solar electric energy ranging from 0.003% to 0.029%. Hydro electricity has a share in the range between 2% and 7.3%.

The minimal share of renewable energy sources in the share of the total energy production and consumption in the Republic of Macedonia indicates that the available resources (e.g. geothermal, hydro, solar power) are insufficiently utilized, but also the aspects of energy security, in terms of all steps that need to be undertaken by the state to prevent threats in relation to planned demands for energy by the national economy. Energy security or threat to economy and social welfare the factors of which are minimized with reduction of dependence on energy and energy resources import, indicate the importance of social resources streamlining towards maximum utilization of natural renewable sources.

## Methodology

- Methodology for the indicator calculation

Statistical methodology for calculation:

- Regulation on Energy Statistics of the European Parliament and of the Council (Regulation no.1099/2008),
- "Energy Statistics Methodology Eurostat F4, 1998"

## Policy relevance of the indicator

- Strategy for Energy Efficiency in the Republic of Macedonia by 2020<sup>1</sup>
- Strategy for Energy Development in the Republic of Macedonia by 2030<sup>2</sup>
- Strategy for Utilization of Renewable Energy Sources (RES) in the Republic of Macedonia by 2020<sup>3</sup>

## Legal grounds

Law on Energy; Energy Balance of the Republic of Macedonia - annual planning document defining the demands for energy and the possibility for their supply (Article 16 of the Law on Energy).

## Targets

Reduction in the dependence on imported fuels and reduction in inefficient energy consumption; Modernization of energy infrastructure and diversification of energy supply; Participation in of regional cooperation and compliance with the legislation of the Energy Community.

## Reporting obligation

- Eurostat
- ECE/UN
- IEA/OECD

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 030	Renewable energy consumption	CSI 030 EE 26	Renewable energy consumption	R	B	energy	annually


<sup>1</sup> <http://www.konkurentnost.mk/StrateskiDokumenti/StrategijazaunapreduvanjenaEERMdo2020godina.pdf>

<sup>2</sup> <http://www.build.mk/docs/users/cloverstack/Strategija%20za%20razvoj%20na%20energetikata%202008-2020%20so%20vizija%20do%202030.pdf>

<sup>3</sup> <http://www.economy.gov.mk/dokumenti/strategii/3102.html>

## MK - NI 031

### RENEWABLE ELECTRICITY



#### Definition

Renewable electricity sources are defined as renewable non-fossil sources of energy, such as: hydropower, geothermal, solar and wind power; solid biomass; biogas, liquid biofuels, etc.

The indicator "Renewable electricity" measures the share of electricity produced from renewable sources in gross national electricity production (in percentage).

The gross national electricity consumption is a sum of the total gross production and import of electricity minus electricity exported.

#### Units

- GWh
- percentage

#### Key question

What is the share of electricity originating from renewable sources in the gross electricity consumption in the Republic of Macedonia?

#### Key message

The share of electricity originating from renewable sources in the gross electricity consumption in the Republic of Macedonia is rather low. It makes relatively high annual fluctuation depending on hydrological conditions, considering that so far only hydro and solar power from among renewable sources contribute to electricity production from renewable sources. Efforts should be made to utilize other renewable sources for electricity production.

There was a variable trend in the rate of renewably energy sources utilization during the analyzed period. In 2010, resulting from favourable hydrological conditions, the share of renewable electricity in the total gross electricity consumption was the highest amounting 28%, while the lowest share was recorded in 2001 amounting 15%. After the fall by 58% in the period 2010 to 2012, it is positive that 2013 recorded growth in the share of renewable energy in the gross electricity consumption in the Republic of Macedonia by 60%.

Figure 1. Share of renewable electricity in gross domestic electricity consumption





Figure 2. Trend in electricity production from renewable sources (%)



Data coverage: [excel](#)

Source: State Statistical Office

## Assessment

The production of electricity from renewable sources in the Republic of Macedonia is based on the hydropower, while minor contribution has been noted by solar electricity for the last four years. Production in large hydropower plants is predominant.

At present, the share of renewable electricity in the overall electricity consumption is very important and depends on hydrological conditions during the year. Variable hydrological conditions result in variations in the production of hydro energy due to lower quantities of precipitation. This indicates the fact that there is a need for investments in installations to enable utilization of other renewable energy sources as well, such as geothermal, solar and wind power and increase in the share of the total electricity production.

During the observed period, there was a variable trend of utilization of energy from renewable sources. In 2010, resulting from favourable hydrological conditions, the share of renewable electricity in the total gross electricity consumption was the highest amounting 28%, while the lowest share was recorded in 2001 amounting 9%. After the fall by 58% in the period 2010 to 2012, it is positive that 2013 recorded growth in the share of renewable energy in the gross electricity consumption in the Republic of Macedonia by 60%.

## Methodology

- Methodology for the indicator calculation

Statistical methodology for calculation:

- Common surveys for coal, oil, natural gas, electricity and heat, renewable energy for 2005 by Eurostat, ECE/UN and IEA/OECD.
- "Energy Statistics Methodology Eurostat F4, 1998"

## Policy relevance of the indicator

- Strategy for Renewable Energy Sources Utilization in the Republic of Macedonia by 2020<sup>1</sup>
- National Strategy for Sustainable Development in the Republic of Macedonia 2009 – 2030<sup>2</sup>
- Strategy for Energy Efficiency Promotion in the Republic of Macedonia by 2020<sup>3</sup>
- Strategy for Energy Development in the Republic of Macedonia by 2030<sup>4</sup>

## Legal grounds

Law on Energy; Energy Balance of the Republic of Macedonia - annual planning document defining the demands for energy and the possibility for their supply (Article 16 of the Law on Energy).

## Targets

To achieve the required increase in order to reach the EU indicative target of 21% share by 2020.

## Reporting obligation

- Eurostat
- ECE/UN
- IEA/OECD

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 031	Renewable electricity	CSI 031 EE 27	Renewable electricity	R	B	energy	annually

<sup>1</sup> <http://www.economy.gov.mk/dokumenti/strategii/3102.html>

<sup>2</sup> <http://www.moep.gov.mk/wp-content/uploads/2014/12/Nacionalna-Strategija-za-Odrziv-Razvoj-vo-RM-NSSD-DeI-1.pdf>

<sup>3</sup> <http://www.konkurentnost.mk/StrateskiDokumenti/StrategijazaunapredovanjenaEERMdo2020godina.pdf>

<sup>4</sup> <http://www.build.mk/docs/users/cloverstack/Strategija%20za%20razvoj%20na%20energetikata%202008-2020%20so%20vizija%20do%202030.pdf>

# FISHERY







## Definition

The indicator shows the number of freshwater species living in rivers and lakes in the Republic of Macedonia and the fish species represented in fishponds that are subject to aquicultural production.

At present, the indicator shows the status of:

- catch of two economically significant fish species in fishing waters;
- total catch of other fish species in fishing waters;
- total aquicultural fish production in fishponds.

## Units

Number of fish species, kilograms (tones) fish catch.

## Key policy issue

How sustainable is the fish catch in the Republic of Macedonia?

## Key message

The average fish catch in the Republic of Macedonia is 1.364 tons of different fish species. The overall fish catch had periodical trend of reduction and increase, from 2003 to 2006 the catch reduced because individual fishing companies, business entities and concessionaires lost their permits for fishing activities in individual aquatic basins and high number of sports fishing clubs were terminated.

The greatest fish cat was noted in 2000 amounting 1834, and the lowest one was in 2006 and amounted 813 tons. Carp is the leading species in lowland waters with a catch of 194 tons in 2012, and trout in highland waters with a catch of 1080 tons in 2012. Trout had the highest share in the total fish catch in 2012 with 73.92%, followed by carp with 13.29%, other fish with 10.19% and catfish had lowest share with 2.6%.

Figure 1. Total fish catch




Figure 2. Catch of the main fish species





Figure 3. Total fish catch by fish species



Data coverage: [excel](#)

Source: State Statistical Office

### Assessment

During the observed period, with regard to total fish catch (production of commercial fish and fishing by sports fishermen) in the Republic of Macedonia, the average catch is 1.364 tons of different fish species. In the period from 1995 to 2003, the overall fish catch had periodical trend of reduction and increase, from 2003 to 2006 the catch decreased because some fishing companies, business entities and concessionaires lost their licenses for fishing activities in

certain water basins, and significant number of sports fishing clubs were terminated.

Trend of increase by 104.6% in the total fish catch was noted in the period 2006 to 2010 compared to 2006, i.e. it increased from 813 to 1664 tons of fish. Then, in 2011 and 2012, there was drop in fish catch again compared to 2010 by 12.2%.

Figure 3 shows that the carp is predominant fish species in lowland waters with a catch of 194 tons in 2012, while trout is leading in highland waters with a catch of 1080 tons in 2012. In the total fish catch in 2012, trout had the highest share with 73.92%, followed by carp with 13.29%, other fish species with 10.19% and catfish with 2.6%.

Fishing and fish stock exploitation in fishponds and artificial water accumulations in the Republic of Macedonia is under permanent supervision, with constant care for the fish stocks and regular stocking with economically important fish species. In this way, sustainable development and exploitation of fish as an important economic resource is provided, as well as for sports fishing. Exploitation of fish stocks from natural lakes has been coping with permanent problems for a longer period, including over-fishing and uncontrolled fish catch in those aquatic ecosystems. These activities affect particularly the endemic fish species, such as Ohrid trout (*Salmo letnica* Kar.), as well as other endemic species represented by small populations in certain aquatic ecosystems.

## Methodology

### Methodology for the indicator calculation

The source of data on the characteristics of fish stocks in the Republic of Macedonia is the State Statistical Office and their methods are used for data processing.

### Uncertainty

Uncertainty derives from the assumed incomplete data on fish catch in rivers and lakes. The uncertainty increases further because of the limited number of literature data on genetic structure of fish populations in natural aquatic ecosystems.

## Policy relevance of the indicator

### List of relevant policy documents

The Study on the State of Biological Diversity in the Republic of Macedonia and the National Strategy for Biological Diversity Protection with Action Plan establish integrated approach to the protection and sustainable use of biological diversity components including fishery.

Spatial Plan of the Republic of Macedonia.

### Legal grounds

The Law on Fishery and Aquiculture regulates the management, planning, commercial management and aquiculture of fish in fishing waters, fish ponds, semi fish ponds, cages and other fish breeding resources.

## Targets

The overall fish production in the Republic of Macedonia has been envisaged to grow by 2.300 tons by 2020. The main mass in this grow will consist of trout fish (1.435 tons or 62% of the total catch) mostly from fishponds.<sup>1</sup>

## Reporting obligation

- FAO – Fisheries and Aquaculture Department

---

<sup>1</sup>Spatial Plan of the Republic of Macedonia

## General metadata


Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MKNI 041</b>	<b>Fish stock characteristics</b>	<b>FISH 3</b>	Fish stock characteristics	<b>S</b>		Water Biodiversity Tourism	annual

# TRANSPORT



## MK - NI 035

### PASSENGER TRANSPORT DEMAND



#### Definition

The indicator "**passenger transport demand**" will be presented in two different ways:

1) To measure decoupling of passenger transport demand from economic growth, the volume of passenger transport relative to GDP will be used. The decoupling indicator is defined as the ratio between pkm (inland modes) and GDP (Gross Domestic Product in constant 2000 EUR). In the presentation of this indicator in the Republic of Macedonia, the index 2000=100 is taken as baseline year.

2) Share of passenger transport: This indicator is defined as the percentage share of each mode of transport in the total inland transport. The unit used is the passenger-kilometre (pkm), which represents one passenger traveling a distance of one kilometre. It is based on transport by passenger cars, buses and coaches and trains.

All data should be based on movements on national territory, regardless of the nationality of the vehicle. However, data collection methodology should be harmonised at the EU level.

#### Units

The unit used is the passenger-kilometre (pkm), which represents one passenger travelling a distance of one kilometre. It is based on transport by passenger cars, buses and coaches, and trains.

Passenger transport demand and GDP are shown as an index 2000=100.

#### Key policy question

***Is the passenger transport in road transport reduced compared to other transport modes?***

***Is the passenger transport demand decoupled from economic growth?***

#### Key message

The overall passenger transport demand has grown by 18.87% during the analyzed period.

The share of road transport in the total passenger transport was 99% in 2013, reflecting an increase by 4.54 % compared to 1990, for railroad transport it was 1% in 2013 or decline by 81.1 % relative to 1990.

The share of the private cars in the total passenger transport is the highest. The share in 2013 was 93.3%, being at the same time the highest share during the analyzed period. Increased demand for cars has direct reflection on the structure of passenger inland transport, negative impacts on environment and health, especially for the fact that high number of vehicles transports low number of passengers. Data on the share of private vehicles and public road transport in cities has been estimated.

Demand for passenger transport per capita was the lowest in 2006 – 2.69 km, and the highest in 2013 with 3.89 km.

Figure 1 Passenger kilometers of individual modes of passenger transport in the total passenger transport




Figure 2 Share of individual modes of passenger transport in the total passenger transport




Figure 3: Demand for passenger transport per capita





Figure 4: Total passenger transport demand/GDP ratio



**Data coverage:** excel

**Data source:** State Statistical Office

### Assessment

The total passenger transport demand during the analyzed period ranged between 5492 and 8024 passenger kilometers, noting increase by 18.87%. The share of road transport in the total passenger transport was 99% in 2013 reflecting increase by 4.54% compared to 1990, and it was 1% in railroad transport in 2013 or decrease by 81.1% compared to 1990.

The share of private cars in the total passenger transport was the highest. In 2013, it had share of 93.3% representing also the highest share during the analyzed period. The lowest share of 57.7% was recorded in 1990. In the period from 2000 to 2006, there was a falling trend by 17.97%, followed by growing trend by 2013 by 46.1%. This indicates that the reduction of environmental pollution requires reduction in the share of road transport in the



share of passenger transport in favour of other transport modes. This would also result in avoided costs for the expensive liquid fuel which while combusting pollutes the environment.

The road public transport in cities throughout the analyzed period noted significant trend of reduction amounting 51.69%,

Inter-city public road transport during the analyzed period noted variable trend with the biggest reduction of 81.1% recorded for railway transport meaning that this transport mode was used at very low rate.

Data on the share of private cars and public road transport in cities was estimated.

The passenger transport demand per capita was the lowest in 2006 with 2.69 km per capita, and the highest in 2013 with 3.89 km per capita.

The trend of the passenger kilometer/GDP ratio from 2000 to 2013 was mainly in decrease. It is indexed for the year 2000=100 in order to monitor the changes in the intensity of passenger transport demand relative to economic growth presented through GDP.

## Methodology

### ■ Methodology for the indicator calculation

In order to measure the decoupling of passenger transport demand from economic growth, the volume (i.e. intensity) of passenger transport relative to GDP is calculated. Relative decoupling occurs when the passenger transport demand rises at rate lower than the one of the GDP. Absolute decoupling occurs when the passenger transport demand falls while GDP rises or remains constant.

The unit used is passenger-kilometre (pkm), which represents one passenger traveling a distance of one kilometre.

With regard to EU Member States, according to Regulation on road transport and Regulation (EC) No 91/2003 on railroad transport statistics, data is based on all movements of passenger transport on the national territory.

### Source of used methodology

Structural indicators of Eurostat on transport

State Statistical Office

### ■ Methodological uncertainty and data uncertainty

All data is based on the movements on the national territory, regardless of the nationality of the vehicle. The methodology of data collection has been harmonised at EU level, but estimated data has been used for the purposes of data calculation on the transport by passenger car. Sources include EUROSTAT, National statistical offices, ECMT, UNECE, UIC, DG TREN.

In order to answer the question whether the passenger transport demand is decoupling from economic growth, the intensity of passenger transport relative to changes in real GDP is considered.

### ■ Uncertainty of data sets

In order to obtain full picture of passenger transport demand and corresponding problems in the environment, it would be very useful to supplement data with data on the number of vehicle-kilometers.

## Policy relevance of the indicator

### List of relevant policy documents

**The National Strategy for Transport** prepared and adopted by the Government of the Republic of Macedonia determines the main directions of the transport policy development in the Republic of Macedonia through identification of goals and development strategy for road, rail and air transport sectors.

### Legal grounds

The road transport is regulated by the Law on Road Transport. It regulates the conditions and the manner of performing transport of passengers and goods in internal and international road transport.

Transportation of dangerous goods is regulated by the Law on Dangerous Goods Transportation in Road and Railroad Transport, regulating the conditions under which transport of dangerous goods shall be performed (preparation of matter, loading, transport, on road procedures, unloading, safety in transportation, vehicles equipment and staff training).

Railroad transportation is regulated by the Law on Railroads, Law on Agreements on Transportation in Railroad Traffic, Law on Agency Regulating Railroad Transport Services Market and Law on Railroad Transport Safety.

## Reporting obligation

- EUROSTAT

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 035</b>	<b>Passenger transport demand</b>	CSI 035	Passenger transport demand	<b>P</b>	<b>B</b>	<ul style="list-style-type: none"><li>▪ Ttransport</li><li>▪ GDP</li></ul>	Monthly Periodically Annually

## MK - NI 036

### FREIGHT TRANSPORT DEMAND



#### Definition

The indicator "**freight transport demand**" will be presented in two different ways:

1) To measure decoupling of freight transport demand from economic growth, the volume of freight transport relative to GDP will be used. The decoupling indicator is defined as the ratio between tkm (inland transport) and GDP (Gross Domestic Product in constant 2000 EUR). Presentation of this indicator in the Republic of Macedonia will be based on the baseline year 2000=100.

2) Modal split share of freight transport: This indicator is defined as the percentage share of each mode of freight transport in total inland transport. The unit used is tonne-kilometre (tkm), which represents movement of one tonne of goods over a distance of one kilometre. It includes road and railroad inland transport. Railroad transport is based on movements on national territory, regardless of the nationality of the vehicle. Road freight transport is based on all movements of vehicles registered in the reporting country.

#### Units

The unit used is the tonne-kilometre (tkm), which represents the movement of one tonne of goods over a distance of one kilometre. It includes transport by road and rail.

Freight transport demand and GDP are shown as an index (2000=100).

#### Key policy question

*Is the freight transport in road transport reduced compared to other transport modes?*

*Is the freight transport demand decoupled from economic growth?*

#### Key message

The freight transport demand noted variable trend of increase and decrease during the analyzed period. In the period from 1990 to 2000, there was a decline in freight transport demand, followed by rapid increase from 2000 and 2006. In the period 2006 to 2009, there was a notable trend of decline and then, in 2010, 2011 and 2012 there was a trend of increase again to be followed by another decline in 2013. Railroad freight transport also showed variable trend of increase and decrease during the analyzed period.

Freight kilometers/GDP ratio in the period 2000 to 2013 noted continuous trend of increase.

Figure 1: Freight transport by modes in tone kilometers (in million km)




Figure 2: Share in percentages of tone kilometers of individual freight transport modes in the total freight transport





Figure 3: Total freight transport demand/GDP ratio



**Data coverage:** excel

**Data source:** State Statistical Office

## Assessment

The freight transport demand noted variable trend of increase and decrease during the analyzed period. Figure 2 shows that the highest share in freight transport belongs to road transport ranging between 59.6% in 2000 and 93.6% in 2003, while railroad freight transport had a share of 6.4% in 2003 to 40.4 % in 2000 of the total freight transport demand. The share of railroad in the total freight transport noted positive trend in 2013 reaching 7.6%, which mitigated the negative impact of transport on the environment.

The above is sufficient indication that the indicator should endeavour towards reduction of road transport in favour of other freight transport modes. The reason for this is that the road freight transport uses costly liquid fuels which during combustion on roads cause significant pollution of the environment, air, soil, biodiversity, etc.

Figure 3 shows that the trend of the index of the total freight transport demand relative to GDP had variable trend of increase and decrease, with domination of growing trend in the period from 2000 to 2006, followed by falling trend domination in the period 2006 to 2013. It is indexed on baseline year 2000=100 in order to monitor the changes in the intensity of freight transport demand relative to economic growth expressed through GDP.

## Methodology

### ■ Methodology for the indicator calculation

In order to measure the decoupling of freight transport demand from economic growth, the volume (i.e. intensity) of freight transport relative to GDP is calculated. Relative decoupling occurs when the freight transport demand rises at rate lower than the one of the GDP. Absolute decoupling occurs when the freight transport demand falls while GDP rises or remains constant.

The unit used is the tonne-kilometre (tkm), which represents movement of one tonne over a distance of one kilometre.

With regard to EU Member States, according to Regulation on road transport and Regulation (EC) No 91/2003 on railroad transport statistics, data is based on all movements of passenger transport on the national territory.

According to Regulation (EC) No 1172/98, data on freight road transport is based on all

movements of vehicles registered in the reporting country. All other data on transport refer mainly to movements on national territory, regardless of the nationality of the vehicle.

#### ■ Sources of used methodology

Structural indicators of EUROSTAT on transport

State Statistical Office.

#### ■ Methodological uncertainty and data uncertainty

All data is based on movements on national territory, regardless of the nationality of the vehicle. Methodology of data collection has been harmonized at EU level. Sources include EUROSTAT, National statistical offices, ECMT, UNECE, UIC, DG TREN.

In order to answer the question whether the freight transport demand is decoupling from economic growth, the intensity of freight transport relative to changes in real GDP is considered.

#### ***Uncertainty of data sets***

In order to obtain full picture of transport demand and corresponding problems in the environment, it would be very useful to supplement data with data on the number of vehicle-kilometers.

## **Policy relevance of the indicator**

#### ***List of relevant policy documents***

**The National Strategy for Transport** prepared and adopted by the Government of the Republic of Macedonia determines the main directions of the transport policy development in the Republic of Macedonia through identification of goals and development strategy for road, rail and air transport sectors.

#### ***Legal grounds***

The road transport is regulated by the Law on Road Transport. It regulates the conditions and the manner of performing transport of passengers and goods in internal and international road transport.

Transportation of dangerous goods is regulated by the Law on Dangerous Goods Transportation in Road and Railroad Transport, regulating the conditions under which transport of dangerous goods shall be performed (preparation of matter, loading, transport, on road procedures, unloading, safety in transportation, vehicles equipment and staff training).

Railroad transportation is regulated by the Law on Railroads, Law on Agreements on Transportation in Railroad Traffic, Law on Agency Regulating Railroad Transport Services Market and Law on Railroad Transport Safety.

## **Targets**

The fourth goal of the National Strategy for Transport is securing sustainable protection of the environment.

## **Reporting obligation**


- EUROSTAT

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MK NI 036</b>	<b>Freight transport demand</b>	CSI 036	Freight transport demand	<b>P</b>	<b>B</b>	<ul style="list-style-type: none"><li>▪ Transport</li><li>▪ GDP</li></ul>	Monthly Periodically Annualy

## MK - NI 054

### ROAD MOTOR VEHICLES BY FUEL TYPE



#### Definition

This indicator defines the number of motor vehicles broken down by road motor vehicle type (passenger cars, motorcycles, motor coaches and buses, commercial vehicles, operational vehicles, trailers and tractors) and fuel type (gasoline, diesel, mixture, gas oil, electricity) on national level.

#### Units

- percentage (%).

#### Key policy question

***What is the share of road motor vehicles by fuel type in the total number of road motor vehicles by vehicle type?***

#### Key message

Emissions of polluting substances originate from almost all economic and social activities, and especially emissions from transport, contribute greatly to overall air emissions. Utilization of alternative energy sources, renewable sources, biofuel and natural gas are all primary processes for air quality improvement.

Use of gasoline by cars is the highest, though with falling trend. At the account of decline in the use of gasoline, use of diesel has recorded increasing trend from 2.3% in 1995 to 35.3% in 2013, which does not assume improvement in terms of environment protection and reduction of air emissions. Other types of fuel have very low share with growing trend and range from 0.5% in 1995 to 3.0% in 2013.

With other types of vehicles – motor coaches and buses, trucks and trailers - diesel is dominant type of fuel with growing trend, followed by gasoline. Other types of fuel have very low share with growing trend.

Figure 1. Share of passenger cars by fuel type in the total number of passenger cars






Figure 2. Share of motor coaches and buses by fuel type in the total number of motor coaches and buses




Figure 3. Share of trucks by fuel type in the total number of trucks





Figure 4. Share of trailers by fuel type in the total number of trailers



Data coverage: [excel](#)

Data source: State Statistical Office, Ministry of Interior

## Assessment

Emissions of polluting substances originate from almost all economic and social activities, and especially emissions from transport, contribute greatly to overall air emissions. Policies and strategies for air pollution reduction are aimed primarily at reducing emissions from transport (combustion of fossil fuels in vehicles). Utilization of alternative energy sources, renewable sources, biofuel and natural gas are all primary processes for air quality improvement.

Data during the reporting period indicate that use of gasoline in cars was the biggest, though with falling trend. In 2005, 97.2% of the cars used gasoline to fall at 61.7% of the cars in 2013. At the account of decline in the use of gasoline, use of diesel has recorded increasing trend from 2.3% in 1995 to 35.3% in 2013, which does not assume improvement in terms of environment protection and reduction of air emissions. Other types of fuel have very low share with growing trend and range from 0.5% in 1995 to 3.0% in 2013.

With other types of vehicles – motor coaches and buses, trucks and trailers - diesel is dominant type of fuel with growing trend, followed by gasoline.

Use of diesel in motor coaches and buses recorded increase from 80.7% in 1995 to 88.1% in 2013, and gasoline recorded drop from 17.3% in 1995 to 9.9% in 2013. Other types of fuel have very low share and range between 1.9% and 4.0%.

As far as trucks are concerned, use of diesel has almost constant share during the whole period, ranging between 67.2% and 72.3%; the same accounts for the use of gasoline with a range between 67.2% and 72.3%. Other types of fuel have very low share and range between 2.0% and 10.1%.

With trailers, as with motor coaches and buses, use of diesel was the biggest with increase from 81.1% in 1995 to 86.6% in 2013, while gasoline recorded decrease by 18.4% in 1995 to 10.1% in 2013. Other types of fuel have very low share and range between 0.6% and 4.1%.

## Methodology

- Methodology for the indicator calculation

Data for the indicator is obtained from the number of road motor vehicles by fuel type and total number of road motor vehicles by type of motor vehicles and is calculated as share of the number of road motor vehicles by fuel type in the total number of road motor vehicles by type of motor

vehicles. The sum of the shares (%) of all types of road motor vehicles by fuel should amount 100 (%), relative to total number of road motor vehicles by type of motor vehicles.

## Policy relevance of the indicator

### List of relevant policy documents

**National Strategy for Transport** prepared and adopted by the Government of the Republic of Macedonia determines the main directions of transport policy development in the Republic of Macedonia through identification of targets and strategy for development of road, railroad and air transport sector.

### Legal grounds

- Law on Road Transport (Official Gazette of RM no. 68/04, 127/06, 114/09, 83/10, 140/10, 17/11, 6/12, 23/13 and 120/13)
- Law on Transport of Hazardous Matters in Road and Railroad Transport (Official Gazette of RM no. 92/07, 161/09, 17/11, 54/11, 13/13, 163/13 and 38/14)
- Law on State Statistics (Official Gazette of RM no. 54/97, 21/07 , 51/11 , 104/13 and 42/14)
- Law on Road Transport Safety (Official Gazette of RM no. 54/07, 86/08, 98/08, 64/2009, 161/09, 36/11, 51/11, 114/12 and 27/14)

## Target

The fourth target of the National Strategy for Transport is securing of sustainable protection of the environment.

## Reporting obligation


- EUROSTAT

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 054	Road motor vehicles by fuel type			D	A	Transport Energy	Monthly Periodically Annually

## MK-NI 055

# AVERAGE AGE OF ROAD MOTOR VEHICLES



## Definition

This indicator classifies road motor vehicles by motor vehicle type (passenger car, motor coaches and buses, commercial vehicles and trailers) and average age on country level.

## Units

- percentage (%).

## Key policy question

*Has the vehicle fleet resulted in decreased average age of vehicles?*

## Key message

Data for the reporting period on all vehicle categories indicates that vehicles aged above 10 years have the highest share in the overall number of vehicles.

The average age of passenger cars has increasing trend ranging between 14.43 and 17.2 years, motor coaches and buses have variable trend of decrease and increase ranging between 20 and 15.3 years. The average age of trucks has decreasing trend and ranges between 16 and 14.1 years. For trailers, the average age has significant decreasing trend from 2001 by 20%, ranging at 16,12 and 12.01 years.

Figure 1. Share of passenger cars by average vehicle age in the overall number of passenger vehicles




Figure 2. Share of motor coaches and buses by average vehicle age in the overall number of motor coaches and buses




Figure 3. Share of trucks by average vehicle age in the overall number of trucks




Figure 4. Share of trailers by average vehicle age in the overall number of trailers





Figure 5. Average vehicle age



Data coverage: [excel](#)

Data source: State Statistical Office, Ministry of Interior

### Assessment

Emissions of transport have great contribution to overall air emissions. Therefore, it is very important to renew vehicle fleet. Data for the reporting period on all vehicle categories indicates that vehicles aged above 10 years have the highest share in the overall number of vehicles.

The average age of passenger cars has increasing trend ranging between 14.43 and 17.2 years, motor coaches and buses have variable trend of decrease and increase ranging between 20 and 15.3 years. The average age of trucks has decreasing trend and ranges between 16 and 14.1 years. For trailers, the average age has significant decreasing trend from 2001 by 20%, ranging at 16,12 and 12.01 years

With regard to passenger cars, the share in the overall number of vehicles in 2013, compared to 1990, aged above 10 years had increasing trend by 24.9%, vehicles aged from 5 to 10 years had decreasing trend by 29.9%, vehicles aged from 2 to 5 years were in decrease by 21.6% and vehicles aged up to 2 years had a trend of decrease by 78.2%.

The share in the overall number of motor coaches and buses in 2013 compared to 1990, of motor coaches and buses aged above 10 years had decreasing trend by 5.4%, motor coaches and buses aged between 5 and 10 years had decreasing trend by 73%, motor coaches and buses aged between 2 and 5 years had variable trend ranging between 0% and 17.4% and vehicles aged up to 2 years had variable trend ranging between 0% and 11.1%.

With trucks during the reporting period, the share in the overall number of trucks aged above 10 years had decreasing trend by 2%, trucks aged between 5 and 10 years had trend ranging between 16.1% and 25.4%, trucks aged between 2 and 5 years had share ranging between 7.6% and 15.1% and vehicles aged up to 2 years had decreasing trend by 78.2%.

The share in the overall number of trailers in 2013 compared to 1990, of trailers aged above 10 years had decreasing trend by 29%, trailers aged between 5 and 10 years had variable trend and ranged between 21.4% and 48.6%, trailers aged between 2 and 5 years had decreasing trend by 84.2% and vehicles aged up to 2 years had variable trend ranging between 0% and 13%.

## Methodology

- Methodology for the indicator calculation

Data for the indicator is obtained from the number of road motor vehicles by type and year of manufacturing, prepared by age groups and is calculated as share of the number of road motor vehicles by type. The sum of the shares (%) of all types of road motor vehicles by age groups should amount 100 (%), relative to total number of road motor vehicles by type of motor vehicles by age groups.

## Policy relevance of the indicator

### List of relevant policy documents

**National Strategy for Transport** prepared and adopted by the Government of the Republic of Macedonia determines the main directions of transport policy development in the Republic of Macedonia through identification of targets and strategy for development of road, railroad and air transport sector.

### Legal grounds

- Law on Road Transport (Official Gazette of RM no. 68/04, 127/06, 114/09, 83/10, 140/10, 17/11, 6/12, 23/13 and 120/13)
- Law on Transport of Hazardous Matters in Road and Railroad Transport (Official Gazette of RM no. 92/07, 161/09, 17/11, 54/11, 13/13, 163/13 and 38/14)
- Law on State Statistics (Official Gazette of RM no. 54/97, 21/07, 51/11, 104/13 and 42/14)
- Law on Road Transport Safety (Official Gazette of RM no. 54/07, 86/08, 98/08, 64/2009, 161/09, 36/11, 51/11, 114/12 and 27/14)

## Target

The fourth target of the National Strategy for Transport is securing of sustainable protection of the environment.

## Reporting obligation


- EUROSTAT

### General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MK NI 055</b>	<b>Average age of road motor vehicles</b>	TERM 033	Average age of the vehicle fleet	<b>D</b>	<b>A</b>	Transport	Monthly Periodically Annually




HEALTH



## **MK - HI 042**

# **AIR POLLUTION AND LEAD IN BLOOD – LEAD LEVEL IN CHILDREN’S BLOOD**



## **Definition**

This indicator tracks the exceedances of Guideline values of lead level in children’s blood in urban areas. The level of lead in the blood of children is expressed as average value of individual concentrations of lead in the blood in micrograms/decilitres ( $\mu\text{g}/\text{dl}$ ).

Excess in air quality limit values occurs when the concentration of pollutant exceeds the limit values specified for lead in the Decree on the limit values of levels and types of polluting substances in ambient air (2005).

Where there are several limit values (see the section on Policy goals), the indicator uses the most stringent case:

- Lead (Pb): annual limit value in ambient air
- Lead (Pb): level of lead in blood

## **Units**

- Lead (Pb) concentrations in ambient air are expressed in micrograms/ $\text{m}^3$ .
- Lead level in the blood is expressed in micrograms/ $\text{dl}$ .

## **Key policy issue**

**What progress has been made towards reduction of pollutants concentrations in urban environments in order to achieve the limit values for ambient air in urban environments defined in the Decree?**

What progress has been made towards reduction of lead level in blood of vulnerable segment of the population, such as children in urban environments defined in the WHO Guidelines?

## **Key message**

There has been a trend of reduction in the lead concentrations below the limit values in the period presented.

Figure 1. Lead concentrations in air in Skopje and Veles





Figure 2. Level of lead in children's blood in Veles



Data coverage: [excel](#)

Source: Public Health Institute of the Republic of Macedonia

## Assessment

### Lead

The lead in the air most frequently originates from lead and zinc ores smelting. The occurrence of high lead concentrations in the ambient air in Veles in the period before 2003 resulted from the emissions released by Lead and Zinc Smelter in Veles.

High difference in lead concentrations in Veles measured in the period 2001-2002 was due to the emissions released by Lead and Zinc Smelter, while lead concentrations in ambient air decreased between 2003 and 2005 as a result of the terminated operation of the Smelter.

The level of lead in the blood of school aged children between 10 and 14 has been falling along with the improvement of the ambient air quality with regard to lead as air pollutant. In the period from 2001 to 2003, the level of blood with children in Veles was in excess of the levels recommended by the WHO. In 2004, when the Lead and Zinc Smelter was out of operation, the recorded levels of lead in school aged children's blood was below the guideline value of WHO. In 2005, there was another increase in the level of lead in children's blood.

The graphical presentation shows the falling trend in lead concentrations in the ambient air in the period between 2001 and 2009.

The graphical presentation shows the falling trend in the levels of lead in children's blood in the period between 2001 and 2005. The latest measurement of lead concentrations in blood was conducted in 2005.

## Methodology

- Methodology for the indicator calculation

### Lead in ambient air – Pb $\mu\text{g}/\text{m}^3$

For each station located in urban environment, annual lead concentration in ambient air (annual limit value of 0,5  $\mu\text{g}/\text{m}^3$ ) is calculated using available 24 hourly data during entire year. Selected urban stations include stations of the following type: stations measuring the pollution from the traffic, stations measuring pollution from industry and so called urban background stations. The average concentration is obtained by averaging the results from all stations located in Veles. In Skopje, there is only one monitoring station with discontinued monitoring.

### Lead in children's blood – Pb/blood $\mu\text{g}/\text{dl}$

In 2005, analysis was made on vein blood taken from school aged children between 10 and 14 in Veles and the individual levels of blood recorded have provided the average level of lead in the blood (Guideline value of WHO is below 10  $\mu\text{g}/\text{dl}$ ). Analysis is made to estimate the level of current exposure of children to lead in urban environment resulting from historical pollution (up to 2003) and pollution caused by traffic.

## Uncertainty

- Methodological uncertainty

Generally, data is not representative for the whole urban environment in the Republic of Macedonia. After the closure of the Lead and Zinc Smelter in Veles and elimination of lead from petrol, concentrations of lead in ambient air have dropped significantly. Compared to the defined methodology of the European Environmental Agency, where the indicator is calculated using data only from the so called urban background stations, in our indicator calculation we have used data on ambient air quality with regard to lead content produced by one measuring station in the Municipality of Centre in the City of Skopje, with monitoring of discontinuous type. Such monitoring has not been established in other cities of our country.

- Uncertainty of data

Data is generally not representative for the whole urban environment in the Republic of Macedonia. According to the defined methodology of the European Environmental Agency, only data series from monitoring stations with at least 75% coverage in the course of one year shall be taken into account in the indicator calculation (or more than 274 valid daily data during one calendar year). In our case, this was not observed in the use of data from the monitoring station in Skopje. Representativeness

of choice in the case of monitoring stations in Veles on ambient air quality with regard to lead content complies with the requirements of the EU Directive 1999/30/EC.

## Policy relevance of the indicator

List of relevant policy documents:

The Second National Environmental Action Plan –2 sets the improvement of air quality through reduction of basic pollutants emission as its main priority. The same document specifies two basic measures that need to be taken: to prepare national plan for ambient air protection and strengthening the ambient air quality monitoring and assessment process.

The 1999 National Environmental Health Action Plan identifies the goal of implementation of targeted epidemiological investigations of the health status primarily with vulnerable population groups in health risky areas (Lead and Zinc Smelter - Veles) in terms of air pollution (non-ferrous metallurgy).

Legal grounds

The Law on Environment regulates areas that have direct impact on the quality of air, i.e. should contribute to air emissions reduction. Thus, the Law regulates the issues of IPPC, EIA/SEA, local environmental action plans and climate change.

The Law on Ambient Air Quality - The goals of this Law include: avoidance, prevention and reduction of harmful effects on human health and environment as a whole, prevention and reduction of pollutions leading to climate change, as well as provision of adequate information on the ambient air quality.

The Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term targets (2005).

The Law and the Decree transpose the requirements of the following EU Directives:

- Framework Air Quality Directive 96/62/EC on ambient air quality assessment and management
- Directive 1999/30/EC on limit values for sulphur dioxide, nitrogen dioxide and nitrogen oxides, suspended particulate matters and lead in ambient air.

The level of lead in ambient air and lead in the blood of exposed population are defined in the World Health Organization Air Quality Guidelines of 1987 and 2000.

## Targets

The Decree on the limit values of levels and types of polluting substances in ambient air and on the alert thresholds, deadlines for the limit values achievement, margins of limit value tolerance, target values and long-term targets specifies the limit values for lead. The WHO Air Quality Guidelines define the guideline values for the lead concentration in ambient air and lead level in the blood of exposed population.

Regular monitoring enables the values of lead concentrations to be maintained below the limit values, namely:

Annual limit value of 0.5 µg/m<sup>3</sup> in ambient air.

Lead level in children's blood below 10 µg/dl.

## Reporting obligation

European Environmental Agency

- Exchange of air quality data based on the Council Decision on the establishment of reciprocal exchange of information and data from all networks and individual ambient air quality measuring stations (97/101/EC).

World Health Organization - ENHIS

- Levels of lead in children's blood, according to WHO Air Quality Guidelines, 1987 and 2000.

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 004</b>	<b>Excess in air quality limit values in urban areas</b>	CSI 004 AP14	Exceedance of air quality limit values in urban areas	<b>S</b>	<b>A</b>	Air Air quality	annually
<b>MK NI 042</b>	<b>Level of lead in children's blood</b>	ENHIS RP G4_Chem_Ex1	Blood levels in children	<b>S</b>		Health	annually

## **MK – NI 043**

### **MORTALITY DUE TO RESPIRATORY DISEASES (J00 - J99) WITH INFANTS**



#### **Definition**

The indicator shows mortality rate of respiratory diseases (J00 - J99) with infants. In this context, infants are alive born children aged 1 to 12 months, respiratory diseases are acute or chronic condition of respiratory systems, including acute respiratory infections, bronchitis, pneumonia or influenza (J00—J99). According to data available from epidemiological studies, infants' mortality could be in correlation with the level of air pollution in the environment, along with series of other factors that can influence respiratory diseases mortality and morbidity. Therefore, the indicator is considered useful in the assessment of the load of diseases attributed to ambient air quality.

#### **Units**

- Number of deaths per 1000 alive born infants aged between 1 and 12 months.

#### **Key policy issue**

**What steps have been taken to reduce or prevent the increase in the mortality rate from respiratory diseases (J00 - J99) with infants?**


**What cross-sectoral policies have been implemented so far in order to reduce respiratory morbidity and mortality with the general population and with infants in particular?**

#### **Key message**

Respiratory diseases with infants are not attributable only to air pollution, but also to biological and genetic factors of infants, presence of allergens, infective agents, nutrition, socio-economic factors and parents' educational background.

The analysis of the trend indicates variable conditions in the reporting period with an evident drop in 2005, which then raised again in the next year. Mortality rate is also dependent on the effectiveness of the health system and availability of health care services.

Figure 1. Mortality rate from respiratory diseases (J00-J99)



Data coverage: [excel](#)

Source: State Statistical Office

## Assessment

The mortality rate from **respiratory diseases (J00 - J99)** with infants, in the period 2002 – 2008, ranged between 0.18 and 0.52. The increased mortality rate with infants from respiratory diseases can be attributed to exposure at pollutants in the ambient air both indoors and outdoors, while the falling of the rate results from improved conditions and air quality. However, such significant changes in the levels of air pollution were not observed. Different criteria in diagnosis setting and reporting of death and causes leading to it may have had influence on this variable trend.

Respiratory diseases as cause of death in the general population are ranked at fifth position in the structure of causes for death in general population.

## Methodology

- Methodology for the indicator calculation

The indicator of mortality from respiratory diseases (J00 - J99) with infants is calculated as a rate of infants that died from respiratory diseases (J00 - J99) per 1000 infants aged from 1 to 12 months.

## Policy relevance of the indicator

### List of relevant policy documents

**The Second National Environmental Action Plan –2** sets the improvement of air quality through reduction of basic pollutants emission as its main priority. The same document specifies two basic measures that need to be taken: to prepare national plan for ambient air protection and strengthening the ambient air quality monitoring and assessment process.

**The 1999 National Environmental Health Action Plan** identifies the goal of implementation of targeted epidemiological investigations of the health status primarily with vulnerable population groups in health risky areas (Lead and Zinc Smelter - Veles) in terms of air pollution (non-ferrous



metallurgy).

**The National Children’s Environment and Health Action Plan** in which the current children’s health profile in the country is presented, identifies the current environmental health risks to children from air pollution, indoors primarily passive tobacco smoking and outdoors, originating from the processes of heating, traffic and industry.

**The Environmental Health Action Plan for Europe and the Handbook – Planning to protect Children against hazards** – The Regional Priority Goal III reads: We aim to prevent and reduce respiratory diseases due to outdoor and indoor air pollution, thereby contributing to a reduction in the frequency of asthmatic attacks in order to ensure that children can live in an environment with clean air. *We aim to achieve a substantial reduction in the morbidity and mortality from acute and chronic and respiratory disorders in children and adolescents.”*

## Legal grounds

The **Law on Environment** regulates areas that have direct impact on the quality of air, i.e. should contribute to air emissions reduction. Thus, the Law regulates the issues of IPPC, EIA/SEA, local environmental action plans and climate change.

The **Law on Ambient Air Quality** – the principle of precautionary and accountable behaviour stipulates that: In the course of activities that may affect the quality of ambient air, everyone shall be obliged to behave in a careful and responsible manner in order to avoid and prevent ambient air pollution and harmful effects on human health and environment as a whole.

## Targets

To reduce the mortality rate from respiratory diseases (J00 - J99) with infants, or prevent the mortality rate increase.

## Reporting obligation

- Every doctor stating death is obliged to fill in the reporting list on the death event, stating the causes for the death and then such data is collected in the national database of the State Statistical Office

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 043	Mortality due to respiratory diseases (J00—J99) with infants	ENHIS Air_E2	Mortality due to respiratory diseases	S		Air Health Traffic Local self-government	annually



## Definition

Incidence of leukemia is the rate of new diagnosed cases of leukemia defined by ICD-10 codes C90-95 in children aged 0 to 14 years.

## Units

- Number of diagnosed cases of leukaemia per 100 000 persons aged 0 to 14 years.


## Key policy issue

**What progress has been made to reduce the incidence of childhood leukaemia?**

## Key message

The incidence of childhood leukaemia shows a rising trend in the reporting period 2002-2007, with a peak of the incidence in 2004 and especially significant increase in 2007.

Figure 1. Incidence of leukemia among children aged 0-14 years



Data coverage: **excel**

Source: National Register of malignant diseases

## Assessment

The indicator uses data from the national register of malignant diseases for the period 2002 - 2007. The indicator is based on data on children aged 0 - 14 years.

Considering the fact that causal factors of leukaemia are not fully clear, policies to reduce incidence

are difficult to formulate or have limited effects. For example, policies to reduce exposure to ionising or electromagnetic radiation potentially prevent only a small portion of leukaemia cases. Therefore, it is necessary to undertake further coordinated research into environmental impacts on leukaemia and environmental/genetic interactions. It is particularly important to monitor childhood leukaemia. National registers with continual follow-up that employ standardized or comparable methods and should include environmental health indicators should be universal.

In the context of public health, leukaemia related mortality is an important co-indicator for assessment of the quality of health care system.

## Methodology

- Methodology for the indicator calculation

National estimates of incidence rates are standardized to world standard population aged 0 – 14 years: number of new cases per 100000 person-years.

## Policy relevance of the indicator

### List of relevant policy documents

Leukaemia is often discussed when environmental issues and childhood diseases are considered. The causes of the majority of cases are, however, unknown. As a result, there is a lack both of policies aiming directly at reducing the incidence of leukaemia and of major programmes fostering research into the potential risk factors for leukaemia in Europe.

The new Regulation of the European Parliament and the Council concerning the Registration, Evaluation, Authorization and Restriction of Chemicals (REACH) is of relevance. It considers that the carcinogenicity, mutagenic and reproductive toxicity of chemical industrial substances are priority criteria when they are submitted to security constraints and declarations authorizing their use. The target of REACH is to substitute progressively substances that are known to be safer for most carcinogenic, mutagenic and toxic industrial substances. Of further relevance is Council Directive 97/43/EURATOM, which aims to protect patients from excessive exposure to radiation for medical use and ensure that there is minimum exposure during pregnancy and early childhood.

**The Second National Environmental Action Plan - 2 (2006)**, as its main goal, sets the achievement of environmental quality by which concentrations of pollutants will not lead to significant impacts or risks to human health, establishment of effective system of prevention, control and assessment of health risks in accordance with the requirements of the national and EU legislation, as well as with WHO recommendations, through taking the specific measure of NEHAP review and further implementation, with particular accent on the risks to children's health.

**The 1999 National Environmental Health Action Plan**, in its section on ionising and non-ionising radiation, specifies the priorities and specific activities that need to be undertaken to reduce negative impacts from these radiations on human health.

### Legal grounds

The **Law on Chemicals**, the basic principle of the Law is the precaution principle, placing human health safety and environment protection on top.

**The Law on Ionising Radiation** incorporates provisions concerning the protection of the population against ionising radiation. The legal person causing ionising radiation, in case of caused release of radioactive substances in the environment, and consequently adverse impacts on human health, shall compensate the damage.

## Targets

To reduce leukaemia incidence in children aged 0 to 14.

## Reporting obligation


- Every doctor establishing diagnosis of leukaemia is obliged to fill in the reporting list on the malignant disease, and then such data is delivered to Regional public health centres, further delivered to the national Public Health Institute

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by <b>DPSIR</b>	Type	Linkage with area	Frequency of publication
<b>MK NI 044</b>	Incidence of childhood leukaemia	ENHIS RPG4_Rad_E1	Incidence of childhood leukaemia	<b>S</b>		Health Waste Consumption of radioactive preparations	Annually

## ***MK – HI 045***

# **INCIDENCE OF MELANOMA IN PEOPLE AGED UNDER 55 YEARS**



## **Definition**

The incidence of melanoma in people aged under 55 years is the number of detected cases during one year expressed per 100 000 residents of selected population.

The incidence is obtained by calculating the number of cases on the mean population size during the period considered. The age-standardized rate is calculated using the age groups. The age group considered here are those aged 0–54 years.

## **Units**

- Number of cases per 100 000 persons-year.

## **Key policy issue**


**What steps have been taken to prevent the rising trend of melanoma?**

**What cross-sectoral policies have been implemented so far to reduce the exposure of the general population and especially children, as it seems that the latter population is the most sensitive and exposure during this period of life leads to consequences in later age?**

## **Key message**

Melanoma is one of cancers with the most rapid rate of growth among people in Europe, which is also the case with us. Melanoma rate trends differ in different regions: in North Europe, where rates are high, it seems that they have been stopped as of the 1990s, especially among people aged below 55. This seems to be a result of changed pattern of stay on sun and protective behavior towards UV radiation among younger generation. Contrary to this, the rates in South and Eastern Europe, where the Republic of Macedonia belongs, still note significant rise in all age groups.

Figure 1. Melanoma incidence rate in the population aged 0-54 years



Data coverage: **excel**

Source: National Register of malignant diseases, State Statistical Office

## Assessment

The incidence of melanoma ranges between rates of 0.9 in 2001, retained at 1.4 during the follow-up years, which means that it tracks an increasing trend.

The analysis of the incidence of melanoma over the period 2001-2007 shows significant rising trend in the 2001-2002 period, which is then retained in the next years, but with increased incidence rate. The rising trend is an alarm for taking preventive measures, i.e. protecting the population against UV radiation.

National policies for reduction of exposure at artificial UV radiation, including regulations on sunbeds use by children and teenagers should be implemented in many countries of the WHO European Region.

Excessive exposure at UV radiation can be best prevented through regional and local campaigns for public awareness increase and information, especially in educational institutions. The goal is to encourage school age children to undertake self-protection measures against sun.

The above policy is supported by the experiences from West European countries. Stagnation as of 1990s following the precedent rising trend in North Europe among people aged below 55 supports the conclusion specific protection activities against UV radiation in these countries are effective.

## Methodology

- Methodology for the indicator calculation

The incidence of melanoma in people aged under 55 years is calculated as the number of new diagnosed cases of melanoma during one year per 100 000 residents.

## Policy relevance of the indicator

## List of relevant policy documents

There are at present few official regulations in most European countries on policies to reduce excessive exposure of children to UV radiation. Melanoma is strongly linked with exposure to UV radiation during childhood and is therefore largely preventable. WHO has launched the INTERSUN Global UV Project to stress the importance of increasing awareness and knowledge about the potential negative health effects of exposure to UV radiation, especially during childhood. This information should be readily available through various channels such as television, radio, campaigns, meteorological websites and in schools. A UV radiation index can help to identify appropriate action based on the measured UV radiation levels. Furthermore, the use of sunbeds by children should be strongly discouraged, if not forbidden. The INTERSUN Project recommendations can serve as a framework for an action plan to reduce exposure to UV radiation.

**The Second National Environmental Action Plan - 2 (2006)**, in its part on non-ionising radiation, specifies the main targets and measures towards the establishment of effective system of environment protection and control against harmful effects of non-ionising radiation in the Republic of Macedonia.

**The 1999 National Environmental Health Action Plan**, in its section on non-ionising radiation, specifies the priorities and specific activities that need to be undertaken to reduce negative impacts from this radiation on human health.

**The National Strategy for Climate Change**, in its part 6.6. Health, provides an overview of climate change impacts on human health.

## Targets

To reduce the incidence of melanoma through implementation of cross-sectoral policies aimed at public awareness increase and improved education of the population.

## Reporting obligation


- Every doctor establishing diagnosis of melanoma is obliged to fill in the reporting list on the malignant disease, and then such data is delivered to Regional public health centres, further delivered to the national Public Health Institute

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
MK NI 045	Incidence of melanoma in people aged under 55 years	ENHIS RPG4_Uvrd_E1	Incidence of melanoma in people aged under 55 years	S		Health Climate change Tourism	Annually

## MK – HI 046

# MORTALITY FROM TRAFFIC ACCIDENTS WITH CHILDREN AND YOUNG PEOPLE



## Definition

The indicator shows the mortality rate from traffic accidents for children aged 0 - 14 years and young people aged 15 - 24, the trend in a given period and comparison of data on European Region and policy relevance related thereto.

## Units

- Number of deaths per 100 000 children aged 0 - 14 years or per 100 000 young people aged 15 - 24 years, respectively.

## Key policy issue


**What steps have been taken to reduce or prevent the increase of the mortality rate from traffic accidents with vulnerable groups like children and young people?**

**What cross-sectoral policies have been implemented to reduce the number of traffic accidents for the general population and especially children population, as it appears that this population is the most sensitive and exposure in childhood results in consequences at later age?**

## Key message

The mortality rate for children and young people is lower in our country compared to specific mortality rates in other European countries like Greece, Spain, France, Germany. Nevertheless, this rate is sufficient to indicate the need for introduction of intervention programmes as part of cross-sectoral policies.

Figure 1. Mortality from traffic accidents, rate at 100 000, age 0-14 and 15-24 years



Data coverage: [excel](#)

Source: State Statistical Office



## Assessment

The mortality rate for children aged 0-14 years and young people aged 15-24 has been relatively stable in the period 2002 to 2004, with significant rise in 2005. The rise in the rate in 2005 could as well reflect the improved system of causes of death reporting. The period 2005 to 2009 was characterized with variable trend of mortality rate reduction and increase.

## Methodology

- Methodology for the indicator calculation

The indicator mortality from traffic accidents (800 and 848) is calculated as rate of deaths attributable to traffic accidents involving persons aged 0 - 14 years and aged 15 - 24 years per 100 000 residents from among the said age groups.

## Policy relevance of the indicator

### List of relevant policy documents

**The National Children and Environment Protection Action Plan** which presents the current health profile of children in the country defines the existing environmental health risks for children for the purpose of their protection against accidents and trauma.

**The Gudiebook - Planning of Children's Protection against Hazards – Regional Priority Goal II** reads: "We aim to prevent and significantly reduce health consequences from accidents and injuries and pursue a decrease in morbidity from lack of adequate physical activity, by promoting safe, secure, and supportive human settlements for all children".

### Legal grounds

**The Law on Traffic Safety at Roads** - This Law regulates the safety and protection on roads; traffic rules on roads and the system of traffic signs and equipment; obligations in case of traffic accident; conditions for acquiring the right to drive a vehicle; candidates training for drivers; driving examination taking and checking of driver's ability; checking of vehicles, devices and equipment that are compulsory for the vehicles; dimensions, total mass; axial loading of vehicles and conditions that shall be met by vehicles in traffic; technical check-ups of vehicles; special safety measures; organization and tasks of the safety at roads councils, as well as misdemeanour sanctions and misdemeanour procedure administered with regard to misdemeanours in the field of traffic at road.

## Targets

To reduce the mortality rate from traffic accidents with children and young people populations through appropriate intervention programmes.

## Reporting obligation

- Every doctor stating death is obliged to fill in the reporting list on the death event, stating the causes for the death and then such data is collected in the national database of the

## General metadata

Code	Title of the indicator	Compliance with CSI/EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 046</b>	Mortality from traffic accidents in children and young people	ENHIS Traf_E1	Mortality from traffic accidents	<b>S</b>	A	Health Transport Local self-government Physical planning	Annually

# TOURISM



**MK – NI 047-1**

## TOURISM INTENSITY IN THE REPUBLIC OF MACEDONIA



### 1 International tourist intensity

#### Definition

The indicator shows the total number of foreign tourists by years at country level and by statistical regions and structure of visitors by country of origin.

#### Units

- Number

#### Key policy issue

**Does the number of tourists in the Republic of Macedonia have development dimension?**

#### Key message

With regard to international tourist visits, the total number of foreign tourists during the analyzed period has had development nature with a rising trend of 229%.

Greece contributes significantly to the number of tourists in Macedonia with 429.475 tourists during the observed period. By statistical regions, the highest number of foreign tourists was recorded in Skopje and Southwestern regions. With regard to foreign tourists arrivals by types of resorts, the highest the highest number of tourists was recorded in Skopje amounting 1.153.147, and the lowest in spa resorts with 39.995 tourists.

Considering that tourism is organized activity, it is necessary to monitor these parameters and activate organizations structures in environment protection and improvement through timely interventions and planning activities.

Figure 1. Total number of foreign tourists




Figure 2. Total number of foreign tourists by country of origin in the reporting period




Figure 3. Countries with significant share in the number of foreign tourists




Figure 4. Foreign tourists arrivals by statistical regions





Figure 5. Foreign tourists arrivals by types of resorts in the period from 2003 to 2013



**Data coverage:** excel

**Data source:** State Statistical Office

## Assessment

Data in the Figure indicates that the Republic of Macedonia is visited by a high number of countries. Tourists from Europe, Northern America, Asia and Australia prevail. The structure of visits during the observed period is dominated by tourists from the immediate neighbourhood. Leading position among the first twenty counties making significant share in the overall number of foreign tourists belongs to tourists from Greece, Serbia and Montenegro, Bulgaria, Turkey, Albania and Serbia, while the highest number of tourists from among other continents comes from United States of America and Russia. Development of attractive and receptive base of the Republic of Macedonia should enable greater presence of tourists from Western European countries having longer tourism tradition and thus higher tourist culture level. The number of foreign tourists has notable rising trend from 121.337 tourists in 1997 to 399.680 tourists in 2013. In the reporting period, Greece has

had the highest share with 11.57% or 429.475 tourists, while South Africa has had the lowest share with 0.006% or 234 tourists of the total number of foreign tourists in Macedonia.

According to regional distribution, Skopje region with 1.411.421 tourists and Southwestern region with 1.048.208 tourists area leading regional centers for tourists, representing two differentiated regions of different characteristics. Southwestern region is dominated by resource attractiveness, while Skopje region by business activity possibilities. Other regions possess alternative possibilities deriving from different environments and therefore it is important to monitor the intensity of foreign tourist visits with a view to redistribute the intensity of visits.

With reference to foreign tourists arrivals by types of resorts, Skopje has had the highest share in the total number of tourists with 40.79%, followed by other tourist resorts with 34.43%, other resorts with a share of 19.55%, mountain resorts with a share of 3.8% and the lowest share belongs to spa resorts with 1.41% of the total number of tourists.

## Methodology

- Methodology for the indicator calculation

The trend in tourism development through dynamics and intensity of tourist industry.

The scale and the intensity, as well as the share of individual countries in the total number of foreign tourists arrivals and overnights, national tourists arrivals and overnights, regional distribution and average number of days of stay.

## Policy relevance

### List of relevant policy documents

- **National Strategy for Tourism Development 2009 - 2013**
- **National Environmental Action Plan - 2** - in Section 4.2.6. Tourism, describes the main challenge for sustainable tourism development, implementation of economic potential with minimum possible impact on the environment.
- **Spatial Plan of the Republic of Macedonia** – in its Chapter 5.4. "Tourism development and organization of tourist areas", defines the status, objectives and planning determinations for tourism development.
- **National Strategy for Sustainable Development of the Republic of Macedonia** – in the section on tourism, presents the directions for sustainable development of tourism, within short, medium, and long-term frames, up to 2030.
- **Strategy for Biological Diversity Protection in the Republic of Macedonia with Action Plan** – under measure C.5 "Stimulation of traditional use of biological diversity and eco-tourism", defines the action for identification of sites suitable for eco-tourism.

### Legal grounds

**The Law on Tourist Activity** specifies the conditions and the manner of performing tourist activity (Chapter 15 Services in rural, ethno and eco-tourism), Law on Catering Activity.

**The Law on Environment, the Law on Nature Protection, the Law on Waste Management, the Law on Ambient Air Quality** and the **Law on Waters** regulate partially the requirements for environmental protection in tourist activity.

## Targets

- Integration of the principles of sustainable development and environmental considerations in tourist sector
- Identification of areas of priority importance for tourism development
- Encouragement of exchange of best practices between public and private tourist interests
- Protection of natural heritage and biological diversity in tourist resorts
- Adoption and implementation of legislation in the area of tourism to regulate the protection of the environment
- Promotion of organic farming, healthy food production and especially traditional production of certain products (e.g. cheese, wine), production of honey, herbs growing, etc.

Promotion of certain types of tourism such as wine tourism, hunting tourism, birds observation tourism, etc.

## Reporting obligation

- Yearly to EUROSTAT
- World Tourist Organization (WTO)
- Annual tourist review of tourism and other services
- Five-year interview of foreign tourists in accommodation establishments

### ▪ General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 047 - 1</b>	Tourism intensity in the Republic of Macedonia	TOUR 12	Tourism intensity	<b>D, P</b>	A	Biological diversity Nature Policies Waste Water Air Transport Soil	Yearly  Every five years



# MK – NI 047-2

## TOURISM INTENSITY IN THE REPUBLIC OF MACEDONIA



### 2 Overnights of foreign tourists

#### Definition

The indicator shows the total number of overnights of foreign tourists by years at country level and by statistical regions. Also, it shows the structure of overnights and average stay of tourists by country of origin.

#### Units

- Number

#### Key policy issue

### What is the trend of overnights of foreign tourists in the Republic of Macedonia?

#### Key message

With regard to international tourist visits, the overnights of foreign tourists during the analyzed period have had rising trend of 232%.

The Netherlands had significant share in the overnights of foreign tourists in Macedonia during the analyzed period with 4.09 days average stay of tourists. By statistical regions, the highest number of overnights was recorded in Southwestern and Skopje regions. With regard to overnights of foreign tourists by types of resorts, the highest number of overnights was recorded in other tourist resorts with 2.602.955 overnights, and the lowest number in spa resorts with 233.471 overnights.

Figure 1. Total number of overnights by foreign tourists



Figure 2. Total number of overnights and average stay by foreign tourists by country of origin




Figure 3. Countries with significant share in the average stay of foreign tourists




Figure 4. Overnights by foreign tourists by statistical regions





Figure 5. Overnights by foreign tourists by types of resorts in the period from 2003 to 2013



**Data coverage:** excel

**Data source:** State Statistical Office

## Assessment

Overnights parallel the intensity of visits by foreign tourists in the Republic of Macedonia. Summarized results indicated that the highest number overnights was accomplished by tourists from European countries, with The Netherlands having the highest share in the average stay of tourists with 4.09 days. From among other continents, tourists from Russian Federation had the longest average stay with 3.08 days. Observation of the trends enables us to follow the extent to which the attractiveness of the environment is adjusted to the demands of these visitors.

Regional distribution of accomplished foreign overnights indicates that tourists in Southwestern region have accomplished highest number of overnights, followed by Skopje region, which is

unfavourable circumstance in the context of foreign tourist visits as the visits take place mainly in urban environments. Observation of relations within regional distribution of overnights accomplished by foreign tourists enables considerations of measures undertaken towards accomplishment of overnights in other regions with specific values.

The average number of foreign tourists provides the opportunity to monitor the level of prevalence of the characteristics of environment. The Figure indicates that foreign tourists stay for relatively short time in the Republic of Macedonia. This duration is around 2.26 days at an average during the analyzed period which reflects significant lagging behind the average stay by national tourists which is 4.7 days.

With regard to overnights of foreign tourists by types of resorts, the highest share in the overall number of overnights is recorded in other tourist resorts with 41.39%, followed by Skopje with 34.51%, other resorts with a share of 16.43%, mountain resorts with a share of 3.95% and the lowest share belongs to spa resorts with 3.71% in the total number of overnights.

## Methodology

- Methodology for the indicator calculation

The trend in tourism development through dynamics and intensity of tourist industry.

The scale and the intensity, as well as the share of individual countries in the total number of foreign tourists arrivals and overnights, national tourists arrivals and overnights, regional distribution and average number of days of stay.

## Policy relevance

### List of relevant policy documents

- **National Strategy for Tourism Development 2009 - 2013**
- **National Environmental Action Plan - 2** - in Section 4.2.6. Tourism, describes the main challenge for sustainable tourism development, implementation of economic potential with minimum possible impact on the environment.
- **Spatial Plan of the Republic of Macedonia** – in its Chapter 5.4. "Tourism development and organization of tourist areas", defines the status, objectives and planning determinations for tourism development.
- **National Strategy for Sustainable Development of the Republic of Macedonia** – in the section on tourism, presents the directions for sustainable development of tourism, within short, medium, and long-term frames, up to 2030.
- **Strategy for Biological Diversity Protection in the Republic of Macedonia with Action Plan** – under measure C.5 "Stimulation of traditional use of biological diversity and eco-tourism", defines the action for identification of sites suitable for eco-tourism.

### Legal grounds

**The Law on Tourist Activity** specifies the conditions and the manner of performing tourist activity (Chapter 15 Services in rural, ethno and eco-tourism), Law on Catering Activity.

**The Law on Environment, the Law on Nature Protection, the Law on Waste Management, the Law on Ambient Air Quality** and the **Law on Waters** regulate partially the requirements for environmental protection in tourist activity.

## Targets

- Integration of the principles of sustainable development and environmental considerations in tourist sector
- Identification of areas of priority importance for tourism development
- Encouragement of exchange of best practices between public and private tourist interests
- Protection of natural heritage and biological diversity in tourist destinations
- Adoption and implementation of legislation in the area of tourism to regulate the protection of the environment
- Promotion of organic farming, healthy food production and especially traditional production of certain products (e.g. cheese, wine), production of honey, herbs growing, etc.

Promotion of certain types of tourism such as wine tourism, hunting tourism, birds observation tourism, etc.

## Reporting obligation

- Yearly to EUROSTAT
- World Tourist Organization (WTO)
- Annual tourist review of tourism and other services
- Five-year interview of foreign tourists in accommodation establishments

### ▪ General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 047 - 2</b>	Tourism intensity in the Republic of Macedonia	TOUR 12	Tourism Intensity	<b>D, P</b>	A	Biological diversity Nature Policies Waste Water Air Transport Soil	Yearly  Every five years

## MK – NI 047-3

# TOURISM INTENSITY IN THE REPUBLIC OF MACEDONIA



## 3 National tourists intensity

### Definition

The indicator shows the total number of overnights of national tourists by years at country level and by statistical regions and average stay of tourists.

### Units

- Number

### Key policy issue

**Does the number of national tourists, overnights and average stay have development dimension?**

### Key message

With regard to national tourist visits, the overall number of tourists during the analyzed period has had falling trend of 26%. Furthermore, it is obvious that the accomplished overnights parallel tourist visits and have had falling trend of 35.2% during the analyzed period. The average stay of national tourists of 4.75 days in 2000 dropped to 4.22 days in 2013.

By statistical regions, the highest number and overnights of national tourists were recorded in Southwestern region and the lowest in Northeastern region.

With regard to national tourist arrivals by types of resorts, the highest number of national tourists was recorded in other tourist resorts with 2.105.414 tourists, and the lowest in spa resorts with 216.242 tourists. As for overnights of national tourists by types of resorts, the highest number of overnights was recorded in other tourist resorts with 12.545.945 overnights and the lowest number on Skopje with 333.334 overnights.

Figure 1. Total number of national tourists and number of overnights




Figure 2. Average stay of national tourists




Figure 3. Number of national tourists and number of overnights by statistical regions




Figure 4. Arrivals of national tourists by types of resorts in the period 2003 to 2013





Figure 5. Overnights of national tourists by types of resorts in the period from 2003 to 2013



**Data coverage:** excel

**Data source:** State Statistical Office

**Assessment**



The number of national tourists has had falling trend of 26%; namely, the number of national tourists in 2000 was the highest, and it was the lowest in 2001. Similar characteristics have been observed with overnights accomplished by national tourists, i.e. the highest number of overnights was accomplished in 2000 and the lowest number of overnights in 2001. Also, the results reflect these relations with regard to average stay, showing significant fall in the number of overnights from 2007 to 2013. The shortest average stay of 4.22 days was recorded in 2013 and the longest one in 2003 amounting 5.1 days.

Southwestern region holds prevalence with regard to national tourists distribution, which can be assessed both as favourable aspect and uneven distribution. Observation of these indicators should indicate the extent to which the number of national tourists will increase in other regions as a result of the promotion of environmental elements in touristic offer on the national tourist market.

Furthermore, we may conclude that accomplished overnights follow tourist visits as a result of the attractiveness of the environment and thus the highest number of overnights is recorded in Southwestern region. Observation of overnights will show the extent to which regions will improve attractive base as factor in accomplishing higher number of overnights.

With regard to national tourist arrivals by types of arrivals, the highest share in the overall number of tourists was recorded in other tourist resorts with 60.52%, followed by mountain resorts with 13.48%, other resorts with a share of 13.34%, Skopje with a share of 6.45% and the lowest share was recorded for spa resorts with 6.22% in the total number of tourists.

With regard to overnights of national tourists by types of resorts, the highest share in the total number of overnights was recorded in other tourist resorts with 60,52%, followed by spa resorts with 13.48%, mountain resorts with a share of 13.34%, other resorts with share of 6.45% and Skopje had highest share of 6.22% in the total number of overnights.

## Methodology

- Methodology for the indicator calculation

The trend in tourism development through dynamics and intensity of tourist industry.

The scale and the intensity, as well as the share of individual countries in the total number of foreign tourists arrivals and overnights, national tourists arrivals and overnights, regional distribution and average number of days of stay.

## Policy relevance

### List of relevant policy documents

- **National Strategy for Tourism Development 2009 - 2013**
- **National Environmental Action Plan - 2** - in Section 4.2.6. Tourism, describes the main challenge for sustainable tourism development, implementation of economic potential with minimum possible impact on the environment.
- **Spatial Plan of the Republic of Macedonia** – in its Chapter 5.4. "Tourism development and organization of tourist areas", defines the status, objectives and planning determinations for tourism development.
- **National Strategy for Sustainable Development of the Republic of Macedonia** – in the section on tourism, presents the directions for sustainable development of tourism, within short, medium, and long-term frames, up to 2030.

- **Strategy for Biological Diversity Protection in the Republic of Macedonia with Action Plan** – under measure C.5 "Stimulation of traditional use of biological diversity and eco-tourism", defines the action for identification of sites suitable for eco-tourism.

## Legal grounds

**The Law on Tourist Activity** specifies the conditions and the manner of performing tourist activity (Chapter 15 Services in rural, ethno and eco-tourism), Law on Catering Activity.

**The Law on Environment, the Law on Nature Protection, the Law on Waste Management, the Law on Ambient Air Quality** and the **Law on Waters** regulate partially the requirements for environmental protection in tourist activity.

## Targets

- Integration of the principles of sustainable development and environmental considerations in tourist sector
- Identification of areas of priority importance for tourism development
- Encouragement of exchange of best practices between public and private tourist interests
- Protection of natural heritage and biological diversity in tourist destinations
- Adoption and implementation of legislation in the area of tourism to regulate the protection of the environment
- Promotion of organic farming, healthy food production and especially traditional production of certain products (e.g. cheese, wine), production of honey, herbs growing, etc.

Promotion of certain types of tourism such as wine tourism, hunting tourism, birds observation tourism, etc.

## Reporting obligation

- Yearly to EUROSTAT
- World Tourist Organization (WTO)
- Annual tourist review of tourism and other services
- Five-year interview of foreign tourists in accommodation establishments

### General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 047</b> - 3	Tourism intensity in the Republic of Macedonia - National tourists intensity	TOUR 12	Tourism Intensity	<b>D, P</b>	A	Biological diversity Nature Policies Waste Water Air Transport Soil	Yearly  Every five years



## Definition

Accommodation establishments are supra-structural facilities facilitating visits and stay of tourists in a given environment. Their observation enables the assessment of regional development. The indicator shows the number of accommodation facilities, rooms and beds.

## Units

- Number of facilities, number of rooms and number of beds.

## Key policy issue

**What is the impact of accommodation establishments on the environment?**

## Key message

The number of accommodation units may have both positive and negative impacts. Positive impacts are related to proper utilization of the space for facilities location, and negative impacts are made when the space is occupied by accommodation establishments in inadequate manner.

The total number of accommodation establishments – facilities in the period 2000 to 2013 has had a trend of increase by 70.7%. Relative to the number of establishments, the number of rooms and beds has noted falling trend by 0.9% for rooms and 4.7% for beds, which is due to improved standard of accommodation facilities.

With regard to the structure of accommodation facilities, it is important to underline that increases occur for facilities of hotel nature, while decreases are primarily in the sphere of workers' resorts and uncategorized facilities for accommodation. The number of hotel facilities in 2013 compared to 2008 increased by 31.14, number of holiday houses, apartments and rooms for rent in the same period decreased by 45.7%, and boarding houses by 44.4%.

Figure 1. Total number of accommodation establishments-facilities




Figure 2. Total number of accommodation establishments – rooms and beds




Figure 3. Accommodation establishments – structure of facilities




Figure 4. Accommodation establishments – rooms structure





Figure 5. Accommodation establishments – beds structure



Data coverage: *excel*

Data source: State Statistical Office

## Assessment

In 2008, statistical monitoring of accommodation establishments by stars was introduced for the first time. Categorization of establishments by stars cannot be automatically linked with the categorization by which establishments were monitored earlier and this caused interruption in annual data series.

The total number of accommodation establishments – facilities in the period 2000 to 2013 had a trend of increase by 70.7%. Compared to the number of facilities, the number of rooms and beds had falling trend of 0.9% for rooms and 4.7% for beds, which is due to improved standard in accommodation facilities. With regard to the structure of facilities, it is important to underline that increases are present with facilities of hotel nature, while decreases are observed primarily in the area of workers' resorts and uncategorized accommodation facilities. The number of hotel facilities in 2013 compared to 2008 increased by 31.14%, number of vacation houses, apartments and rooms for rent in the same period reduced by 45.7% and boarding houses by 44.4%. This can be assessed as positive trend because it is a result of the transformation of non-commercial into commercial sector which will have to be more observant of environmental protection standards.

In the category of motels, the number of rooms and beds had highest increase by 174.7% (rooms) and 128.89% (beds), respectively, and the number of rooms and beds in the category of sleeping cars recorded the greatest drop by 68% (rooms) and 77.53% (beds).

## Methodology

- Methodology for the indicator calculation

Development trend of accommodation units.

Share of individual types of accommodation establishments in the total number.

## Policy relevance

### List of relevant policy documents

- **National Strategy for Tourism Development 2009 - 2013**
- **National Environmental Action Plan - 2** - in Section 4.2.6. Tourism, describes the main challenge for sustainable tourism development, implementation of economic potential with minimum possible impact on the environment.
- **Spatial Plan of the Republic of Macedonia** – in its Chapter 5.4. "Tourism development and organization of tourist areas", defines the objectives and planning determinations for tourism development.
- **National Strategy for Sustainable Development of the Republic of Macedonia** – in the section on tourism, presents the directions for sustainable development of tourism, within short, medium, and long-term frames, up to 2030.
- **Strategy for Biological Diversity Protection in the Republic of Macedonia with Action Plan** – under measure C.5 "Stimulation of traditional use of biological diversity and eco-tourism", defines the action for identification of sites suitable for eco-tourism.

### Legal grounds

The Law on Tourist Activity specifies the conditions and the manner of performing tourist activity (Chapter 15 Services in rural, ethno and eco-tourism); Law on Catering Activity.

The Law on Environment, the Law on Nature Protection, the Law on Waste Management, the Law on Ambient Air Quality and the Law on Waters regulate partially the requirements for environmental protection in tourist activity.

### Targets

- Integration of the principles of sustainable development and environmental considerations in tourist sector
- Identification of areas of priority importance for tourism development
- Encouragement of exchange of best practices between public and private tourist interests
- Protection of natural heritage and biological diversity in tourist destinations
- Adoption and implementation of legislation in the area of tourism to regulate the protection of the environment
- Promotion of organic farming, healthy food production and especially traditional production of certain products (e.g. cheese, wine), production of honey, herbs growing, etc.
- Promotion of certain types of tourism such as wine tourism, hunting tourism, birds observation tourism, etc.


### Reporting obligation

- Statistical Yearbook
- WTO
- EUROSTAT

## General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 048</b>	Tourism density and facilities dynamics	TOUR 14	Tourism density	<b>S, P</b>	A	Biodiversity Nature Policies Waste Water Air Transport	Annually





## Definition

Share in BDP is the share of the gross value added, in percentage, in the area of tourism in the total Gross Domestic Product on national level.

## Units

- %.


## Key policy issue

**What is the impact of tourism relying on the characteristics of the environment on the economic development of the Republic of Macedonia?**

## Key message

Share of tourism in the economic development of the Republic of Macedonia is very low. Its share was the highest in 2003 reaching 1.9%, and it was the lowest in 2013 amounting 1.02%. Share of tourism in the overall gross domestic product notes constant trend of decrease.

Figure 1. Share of tourism in GDP in %



**Data coverage:** excel

**Data source:** State Statistical Office

## Assessment

The Figure shows that the share of tourism in GDP is relatively low with falling trend. I.e. with no improvement. Its share was the highest in 2003 reaching 1.9%, and it was the lowest in 2013 amounting 1.02%. Observation should enable insight in the extent in which effects from tourist development on the environment improve.

Revenues acquired on the basis of tourist fee and taxes will enable environment planning and protection and also offer the opportunity to legal and natural persons to improve their living and working conditions.

## Methodology

- **Methodology for the indicator calculation**

Share of value added from tourism in GDP.

## Policy relevance

### List of relevant policy documents

- **National Strategy for Tourism Development 2009 - 2013**
- **National Environmental Action Plan - 2** - in Section 4.2.6. Tourism, describes the main challenge for sustainable tourism development, implementation of economic potential with minimum possible impact on the environment.
- **Spatial Plan of the Republic of Macedonia** – in its Chapter 5.4. "Tourism development and organization of tourist areas", defines the objectives and planning determinations for tourism development.
- **National Strategy for Sustainable Development of the Republic of Macedonia** – in the section on tourism, presents the directions for sustainable development of tourism, within short, medium, and long-term frames, up to 2030.
- **Strategy for Biological Diversity Protection in the Republic of Macedonia with Action Plan** – under measure C.5 "Stimulation of traditional use of biological diversity and eco-tourism", defines the action for identification of sites suitable for eco-tourism.

## Legal ground

**The Law on Tourist Activity** specifies the conditions and the manner of performing tourist activity; **Law on Catering Activity**.

**The Law on Environment, the Law on Nature Protection, the Law on Waste Management, the Law on Ambient Air Quality** and the **Law on Waters** regulate partially the requirements for environmental protection in tourist activity.

## Targets

- Integration of the principles of sustainable development and environmental considerations in tourist sector
- Identification of areas of priority importance for tourism development
- Encouragement of exchange of best practices between public and private tourist interests
- Protection of natural heritage and biological diversity in tourist destinations

- Adoption and implementation of legislation in the area of tourism to regulate the protection of the environment
- Promotion of organic farming, healthy food production and especially traditional production of certain products (e.g. cheese, wine), production of honey, herbs growing, etc.
- Promotion of certain types of tourism such as wine tourism, hunting tourism, birds observation tourism, etc.

## Reporting obligation

- Yearly publication on GDP
- Statistical Yearbook

## General metadata

Code	Title of the indicator	Compliance with CSI EEA or other indicators		Classification by DPSIR	Type	Linkage with area	Frequency of publication
<b>MK NI 049</b>	<b>Economic value of tourism industry</b>	TOUR35	Economic value of tourism industry	<b>D</b>	B	Biodiversity Nature Policies Waste Water Air Transport	Annually